

PLEASE NOTE

This image contains more than one label approved for this product on this date.

When tank-mixing herbicides, ~~always read and follow~~ the most restrictive label recommendations.

FEB 1 - 2000

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under . . .

(See reverse side)

241-337

02/01/2000

PROWL[®]

3.3 EC herbicide

Supplemental Labeling

EPA Reg. No. 241-337

FOR POSTEMERGENCE USE IN FIELD CORN

POSTEMERGENCE TANK MIX APPLICATIONS OF PROWL 3.3 EC PLUS OTHER HERBICIDES

OBSERVE ALL PRECAUTIONARY STATEMENTS ON THE PROWL 3.3 EC LEAFLET LABEL BEFORE USING. SEE THE PROWL 3.3 EC LABEL FOR MIXING AND APPLICATION INFORMATION, WEED SPECIES CONTROLLED, AND FOLLOW CROP RESTRICTIONS. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF HERBICIDE APPLICATION.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of herbicide application.


PROWL 3.3 EC may be tank mixed with any registered herbicide(s) labeled for postemergence use in field corn. PROWL 3.3 EC can be applied based on the timing indicated by the companion label or until corn is 30 inches tall or 8 visible leaf collars (V8) which ever is more restrictive. Use drop nozzles and apply as a directed spray if the corn canopy density prevents applications from reaching the soil and target weeds.

Potential PROWL 3.3 EC tank mix options include but are not limited to the following herbicides.

2,4-D	Buctril ⁴	Laddok S-12 ²	Salvo ⁸
Accent ¹	Buctril+atrazine	Liberty ⁶	Shotgun ⁸
Accent Gold	Celebrity ²	Liberty ATZ	Spirit ³
Atrazine	Clarity ²	LIGHTNING ⁹	Sterling ⁹
Banvel ²	Distinct ²	Marksman ²	Sterling Plus
Basis Gold ³	Exceed ³	NorthStar ³	Tough 5EC ³
Beacon ³	Hornet ⁵	Roundup Ultra ⁷	

Refer to the tank-mix partner product(s) label(s) for application rates and timing by soil types, weed species, and weed or crop growth stage; restrictions, precautions, adjuvant recommendations, weeds controlled and other information. Follow all precautions and restrictions including state and local use restrictions that may apply to specific products.

When tank-mixing herbicides, always read and follow the most restrictive label recommendations.



(See reverse side)

WEEDS CONTROLLED

PROWL 3.3 EC applied in postemergence tank-mixes provides RESIDUAL control of the following weeds when they germinate after the application.

Barnyardgrass	Panicum, Fall
Crabgrass	Pigweeds (Amaranthus), Palmer, Smooth & Redroot
Foxtail – giant, green & yellow	Sandbur, Field
Johnsongrass, Seedling	Waterhemp, Common
Kochia	Waterhemp, Tall
Lambsquarter, Common	

WEEDS SUPPRESSED

PROWL 3.3 EC applied in postemergence tankmixes provides RESIDUAL suppression of the following weeds. Effective control of these weeds may require a program of herbicide use and/or mechanical cultivation to control weed escapes.

Shattercane	Wild proso millet
Smartweed, Pennsylvania	Woolly cupgrass
Velvetleaf	

PROWL 3.3 EC herbicide will not control emerged or established weeds. Adequate soil moisture is required to provide residual weed control or suppression.

Refer to the PROWL 3.3 EC herbicide leaflet label for additional weeds controlled by PROWL 3.3 EC alone.

**Broadcast Rate of PROWL 3.3 EC in Postemergence Tank-Mixes
(Pints per Acre)**

Soil Texture	<1.5% Organic Matter	1.5-3.0% Organic Matter	>3.0% Organic Matter
Coarse	1.8 to 2.4 pts	2.4 to 3.6 pts	2.4 to 3.6 pts
Medium	2.4 to 3.6 pts	2.4 to 3.6 pts	2.4 to 4.8 pts
Fine	2.4 to 3.6 pts	2.4 to 4.8 pts	2.4 to 4.8 pts

Refer to the PROWL 3.3 EC label for additional information regarding WEEDS CONTROLLED, MIXING AND SPRAYING INSTRUCTIONS



- ®Registered Trademark of American Cyanamid Company
- ¹Registered Trademark of E. I. duPont de Nemours and Company
- ²Registered Trademarks of BASF Corporation
- ³Registered Trademarks of Novartis Crop Protection, Inc.
- ⁴Registered Trademark of Rhone Poulenc Ag Company
- ⁵Registered Trademark of Dow AgroSciences LLC
- ⁶Registered Trademark of AgrEvo USA Company
- ⁷Registered Trademark of Monsanto Company
- ⁸Registered Trademarks of Uniroyal Chemical Company
- ⁹Registered Trademark of Terra International, Inc.

NEXT

LABEL

PROWL

3.3 EC herbicide

Supplemental Labeling

EPA Reg. No. 241-337

FOR POSTEMERGENCE USE IN SEED CORN POSTEMERGENCE TANK MIX APPLICATIONS OF PROWL 3.3 EC PLUS OTHER HERBICIDES

OBSERVE ALL PRECAUTIONARY STATEMENTS ON THE PROWL 3.3 EC LEAFLET LABEL BEFORE USING. SEE THE PROWL 3.3 EC LABEL FOR MIXING AND APPLICATION INFORMATION, WEED SPECIES CONTROLLED, AND FOLLOW CROP RESTRICTIONS. THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF HERBICIDE APPLICATION.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PROWL 3.3 EC may be tank mixed with any registered herbicide(s) labeled for postemergence use in seed corn production fields. PROWL 3.3 EC can be applied based on the timing indicated by the companion label or until corn is 20-24 inches tall or 8 visible leaf collars (V8) which ever is more restrictive. Use drop nozzles and apply as a directed spray if the corn canopy density prevents applications from reaching the soil and target weeds.

PROWL may be tank mixed with registered herbicides(s) for postemergence use in seed corn production fields. In these tank-mixes, always read and follow the most restrictive label recommendations.

Refer to the tank-mix partner product(s) label(s) for application rates and timing by soil types, weed species, and weed or crop growth stage; restrictions, precautions, adjuvant recommendations, weeds controlled and other information. Follow all precautions and restrictions including state and local use restrictions that may apply to specific products.

WEEDS CONTROLLED

PROWL 3.3 EC applied in postemergence tank-mixes provides RESIDUAL control of the following weeds when they germinate after the application.

Barnyardgrass	Panicum, Fall
Crabgrass	Pigweeds (Amaranthus), Palmer, Smooth & Redroot
Foxtail - giant, green & yellow	Sandbur, Field
Johnsongrass, Seedling	Waterhemp, Common
Kochia	Waterhemp, Tall
Lambsquarter, Common	

ACCEPTED
FEB 1 2000
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No. 241-337

(See other side)

WEEDS SUPPRESSED

PROWL 3.3 EC applied in postemergence tankmixes provides RESIDUAL suppression of the following weeds. Effective control of these weeds may require a program of herbicide use and/or mechanical cultivation to control weed escapes.

Shattercane	Wild proso millet
Smartweed, Pennsylvania	Woolly cupgrass
Velvetleaf	

PROWL 3.3 EC herbicide will not control emerged or established weeds. Adequate soil moisture is required to provide residual weed control or suppression.

Refer to the PROWL 3.3 EC herbicide leaflet label for additional weeds controlled by PROWL 3.3 EC alone.

**Broadcast Rate of PROWL 3.3 EC in Postemergence Tank-Mixes
(Pints per Acre)**

Soil Texture	<1.5% Organic Matter	1.5-3.0% Organic Matter	>3.0% Organic Matter
Coarse	1.8 to 2.4 pts	2.4 to 3.6 pts	2.4 to 3.6 pts
Medium	2.4 to 3.6 pts	2.4 to 3.6 pts	2.4 to 4.8 pts
Fine	2.4 to 3.6 pts	2.4 to 4.8 pts	2.4 to 4.8 pts

Refer to the PROWL 3.3 EC label for additional information regarding WEEDS CONTROLLED, MIXING AND SPRAYING INSTRUCTIONS.

RESTRICTIONS FOR POSTEMERGENCE APPLICATIONS

To avoid serious crop injury with postemergence applications observe the following restrictions:

Plant into a seedbed that is firm. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed. **PLANT CORN AT LEAST 1 1/2 INCHES DEEP. CORN SEED MUST BE COMPLETELY COVERED WITH SOIL.**

- **DO NOT APPLY PREPLANT INCORPORATED or serious corn injury can result.**
- **DO NOT apply following PROWL 3.3 EC preemergence applications.**
- **DO NOT apply with liquid fertilizer as the carrier.**



® Registered Trademark of American Cyanamid Company.

NEXT

LABEL

PROWL[®]
3.3 EC herbicide

1/2
**Supplemental
Labeling**

EPA Reg. No. 241-337

FOR POSTEMERGENCE USE IN SWEET CORN

OBSERVE ALL PRECAUTIONARY STATEMENTS ON THE PROWL 3.3 EC LEAFLET LABEL BEFORE USING. SEE THE PROWL 3.3 EC LABEL FOR MIXING AND APPLICATION INFORMATION, WEED SPECIES CONTROLLED, AND FOLLOW CROP RESTRICTIONS. THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF HERBICIDE APPLICATION.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INSTRUCTIONS

PROWL 3.3 EC may be applied in sweet corn from early postemergence until corn is 20-24 inches tall or has 8 visible leaf collars (V8) which ever is more restrictive. Use drop nozzles and apply as a directed spray if the corn canopy density prevents applications from reaching the soil and target weeds.


PROWL 3.3 EC alone will not control emerged weeds. PROWL 3.3 EC can be applied as a tank mix with atrazine or other herbicides labeled for use in sweet corn. When using tank mixtures with PROWL 3.3 EC, always read the companion product label(s) to determine the specific use rates by soil types, weed species, and weed or crop growth stage. **Always follow the most restrictive label.**

In addition, follow precautions and restrictions including state and local use restrictions that may apply to specific products

PROWL 3.3 EC or PROWL 3.3 EC tank-mix combination treatments are most effective in controlling weeds when mechanically incorporated or incorporated by adequate rainfall or overhead irrigation within 7 days after application. If cultivation is necessary because of soil crusting, soil compaction or weed germination before rain or irrigation, use shallow tillage.

When PROWL 3.3 EC alone is used, if corn loss occurs due to weather conditions, any crop registered for PROWL 3.3 EC preplant incorporated use can be replanted the same year without adverse effects.

PROWL 3.3 EC is not recommended for use on peat or muck soils.



(See other side)

2p

PROWL 3.3 EC USE RATES FOR SWEET CORN

Recommended use rates for PROWL 3.3 EC alone and in tank mix applications are given in the following table.

**Broadcast Rate¹ per Acre of PROWL 3.3 EC Postemergence
(Pints per Acre)**

Soil Texture	<1.5% Organic Matter	1.5-3.0% Organic Matter	>3.0% Organic Matter
Coarse	1.8 to 2.4 pts	2.4 to 3.6 pts	3.6 pts
Medium	2.4 to 3.6 pts	3.6 pts	3.6 to 4.8 pts
Fine	2.4 to 3.6 pts	3.6 to 4.8 pts	3.6 to 4.8 pts

¹USE THE HIGH RATE FOR EACH SOIL CLASSIFICATION WHEN USING PROWL 3.3 EC ALONE.

RESTRICTIONS FOR POSTEMERGENCE APPLICATIONS

To avoid serious crop injury with postemergence applications observe the following restrictions:

Plant into a seedbed that is firm. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed. **PLANT CORN AT LEAST 1 1/2 INCHES DEEP. CORN SEED MUST BE COMPLETELY COVERED WITH SOIL.**

- **DO NOT APPLY PREPLANT INCORPORATED** or serious corn injury can result.
- **DO NOT** apply following PROWL 3.3 EC preemergence applications.
- **DO NOT** apply with liquid fertilizer as the carrier.



®Registered Trademark of American Cyanamid Company

PE-XXXX 12/16/99