

241-337

02/14/2000

1/2

PROWL[®]

3.3 EC herbicide

Supplemental Labeling

EPA Reg. No. 241-337

FOR POSTEMERGENCE USE IN POPCORN

OBSERVE ALL PRECAUTIONARY STATEMENTS, MIXING AND APPLICATION INSTRUCTIONS, AND ROTATIONAL CROP RESTRICTIONS IN THE PROWL 3.3 EC LEAFLET LABEL FORE USING.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system. This labeling must be in the possession of the user at the time of herbicide application.

GENERAL INSTRUCTIONS

DO NOT APPLY PREPLANT INCORPORATED or serious corn injury can result.

Plant into a seedbed that is firm and free of clods and trash. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed. **PLANT CORN AT LEAST 1 1/2 INCHES DEEP. CORN SEED MUST BE COMPLETELY COVERED WITH SOIL.**

PROWL 3.3 EC or PROWL 3.3 EC tank-mix combination treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application. If cultivation is necessary because of soil crusting, soil compaction or weed germination before rain or irrigation, use shallow tillage (such as rotary hoe), and make certain corn seeds are below the tilled area.

When PROWL 3.3 EC alone is used, if corn loss occurs due to weather conditions, corn or any crop registered for PROWL 3.3 EC preplant incorporated use can be replanted the same year without adverse effects. If corn is replanted, seeding depth must be below retilled area or crop injury may occur.

PROWL 3.3 EC is not recommended for use on peat or muck soils.

Use Method and Timing

Early Postemergence - Apply PROWL 3.3 EC tank mixes postemergence according to corn height and weed size specified on the tank mix partner label. PROWL 3.3 EC alone will not control emerged weeds. For maximum effectiveness, wait at least 7-11 days before cultivating early postemergence treatments.

ACCEPTED
FEB 14 2000
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-337

(See reverse side)

PROWL 3.3 EC Use Rates in Popcorn

Recommended use rates for PROWL 3.3 EC alone and in tank mix applications are given in the following table.

**Broadcast Rate¹ per Acre of PROWL 3.3 Early Postemergence in Popcorn
(Pints per Acre)**

Soil Texture	<1.5 % Organic Matter	1.5-3.0% Organic Matter	>3.0% Organic Matter
Coarse	1.8 to 2.4 pts	2.4 to 3.6 pts	3.6 pts
Medium	2.4 to 3.6 pts	3.6 pts	3.6 to 4.8 pts
Fine	2.4 to 3.6 pts	3.6 to 4.8 pts	3.6 to 4.8 pts

¹USE THE HIGH RATE FOR EACH SOIL CLASSIFICATION WHEN USING PROWL 3.3 EC ALONE.

PROWL 3.3 EC Tank Mixes in Popcorn

PROWL 3.3 EC may be applied in a tank mix with atrazine, Bicep¹, Bicep II, Dual¹, Dual II, Lariat² or Bullet², Lasso² or any other product labeled for use in popcorn. Refer to the companion label for weeds controlled in addition to PROWL 3.3 EC alone.


When using tank mixtures with PROWL 3.3 EC, always read the companion product label(s) to determine the specific use rates by soil types, weed species, and weed or crop growth stage. In addition, follow precautions and restrictions including state and local use restrictions that may apply to specific products. Always follow the most restrictive label.

RESTRICTIONS FOR EARLY POSTEMERGENCE APPLICATIONS

To avoid serious crop injury with early postemergence applications observe the following restrictions:

Plant into a seedbed that is firm. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed. **PLANT CORN AT LEAST 1 1/2 INCHES DEEP. CORN SEED MUST BE COMPLETELY COVERED WITH SOIL.**

- **DO NOT APPLY PREPLANT INCORPORATED** or serious corn injury can result.
- **DO NOT** apply following PROWL 3.3 EC preemergence applications.
- **DO NOT** apply in liquid fertilizer as the carrier.



®Registered trademark of American Cyanamid Company
¹ Trademark of Ciba Corporation
² Trademark of Monsanto Agricultural Products Company