241-337

2-28-2001

Form Approved, OMB No. 2070-0060

United States

	Registration
	Amendment Other
X	Other

OPP Identifier Number

\$EPA	Environmental Protection Agei Washington, DC 20460			X	Amenda Other		261557
	Applicat	ion for F	esticide -	Section			
1. Company/Product Number 241-337			2. EPA Product Manager Mr. James Tompkins 3. Proposed Classification				
4. Company/Product (Name)			PM#	.0		X	None Restricted
PROWL 3.3 E				25			
5. Name and Address of Applicant (Include ZIP Code) BASF Corporation P.O. Box 400 Princeton, NJ 08543-0400			6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No				
Check if th	is is a new address	i	Product Name				
		Sec	tion - II				
Amendment - Explain Resubmission in res	Final printed labels in response to Agency letter dated "Me Too" Application. Other - Explain below.						
Explanation: Use additional page(s) if necessary. (For section I and Section II.) Notification of removal of self-imposed expiration date from attached supplemental label per PR Notice 98-10. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.							
		Sect	ion - III				
1. Material This Product W	ill Be Packaged In:						
Child-Resistant Packaging Yes* No * Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt. No. per	If "Yes		o. per ontainer	2. Type of	Container Metal Plastic Glass Paper Other (S	pecify)
3. Location of Net Contents Information 4. Size(s) Retail Container 5. Loc Label Container					cation of Lab On Label On Label		ns panying product
6. Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled							
		Sect	ion - IV				
1. Contact Point (Complet	e items directly below for identifica	tion of indivi	dual to be con	tacted, if nec	essary, to pro	cess this	application.)
Name Frederick L. Groya, Ph.D. Title Senior G			Telephone No. (Include Area Code) obal Regulatory Affairs Manager 509ะได้-3130				
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. 6. Date Application Received (Stamped)							
2. Signature / 2. Thuya 3			ilor Global Regulatory Affairs Manager				
4. Typed Name Frede	rick L. Groya, Ph.D	5. Date	pruary	21, 20	001		**************************************



Supplemental Labeling

NOTIFICATION

EPA Reg. No. 241-337

FEB 2 8 2001

FOR USE IN SOYBEANS

FOR DISTRIBUTION AND USE ONLY IN INDIANA, MICHIGAN AND OHIO

USE OF PROWL 3.3 EC HERBICIDE ALONE OR IN TANK MIXTURES AS PREEMERGENCE APPLICATIONS NORTH OF INTERSTATE 80

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PROWL 3.3 EC AND TANK MIX PARTNER LEAFLET LABELS BEFORE USING. SEE THE PROWL 3.3 EC AND TANK MIX PARTNER LEAFLET LABELS FOR PRECAUTIONS, GENERAL INFORMATION, MIXING AND APPLICATION AND SPRAYING INSTRUCTIONS USE RATES, USES WITH OTHER PRODUCTS, WEEDS CONTROLLED, ROTATIONAL CROP RESTRICTIONS AND DISCLAIMERS. OBSERVE THE MOST RESTRICTIVE INSTRUCTIONS ON ANY PRODUCT LABEL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

CHEMIGATION: PROWL 3.3 EC may be applied through sprinkler irrigation systems in soybeans. Refer to the PROWL 3.3 EC and tank mix partner labels for use directions for chemigation. **DO NOT** apply tank mixture through any type of irrigation system unless the label instructions on chemigation of all products are followed.

PROWL 3.3 EC is not recommended for use on peat or muck soils.

GENERAL INSTRUCTIONS

PROWL 3.3 EC herbicide controls labeled annual grasses and certain broadleaf weeds as they germinate, but it will not control established weeds.

Destroy existing weeds before applying PROWL 3.3 EC. Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

Observe all use directions given in the PROWL 3.3 EC and tank mix partner label sections on precautions, general information, mixing and application instructions, uses with other products, weeds controlled, rotational crop restrictions and disclaimers. Always follow the most restrictive instructions on any product label.

Preemergence General Instructions for North of I-80

Use the following instructions when applying the recommended preemergence soil applications of PROWL 3.3 EC alone or in labeled tank mixtures in soybeans grown north of Interstate Highway 80 (I-80) in Indiana, Michigan and Ohio. **DO NOT** use this label in any other state.

A preemergence application of PROWL 3.3 EC may be applied at planting or up to 2 days after planting of soybeans. DO NOT APPLY POSTEMERGENCE to soybeans. Apply to a seedbed which is firm and free of clods. PROWL 3.3 EC may be applied in conventional, minimum and no-tillage systems. Observe all other instructions in the PROWL 3.3 EC leaflet label in the section on PROWL 3.3 EC TANK MIXTURES IN MINIMUM AND NO-TILL SYSTEMS.

Preemergence treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application. If moisture is insufficient to activate the herbicides, a shallow cultivation (preferably with a rotary hoe) should be made after emergence of soybeans, but while weeds are small enough to be controlled by mechanical cultivation, otherwise a postemergent herbicide treatment may be required.

A shallow, thorough mixing of soil after herbicide application with a rotary hoe or similar implement will improve consistency of performance and reduce dependency on rainfall to activate herbicides. If at least I inch of rainfall or mechanical incorporation does not occur before soybean emergence, crop injury may occur.

(See reverse side)

DO NOT overlap spray by aerial or ground equipment. Always check tank-mix compatibility with tank mix partners prior to filling spray tank.

Tank Mixtures and Sequential Applications

PROWL 3.3 EC may be preemergence applied north of I-80 using the rates below in tank mixture with the following products: 2,4-DB (Butyrac¹ brand 200), Roundup², Gramoxone Extra³, SCEPTER®, SCEPTER® 70DG, PURSUIT®, PURSUIT® DG, Sencor⁴/Lexone⁵,

Lorox⁵, Lasso², Dual⁶, Canopy⁵, and Preview⁵. Refer to the PROWL 3.3 EC leaflet label and tank-mix partner labels for all other use directions, restrictions and additional weeds controlled for these mixtures.

PROWL 3.3 EC may be preemergence applied north of I-80 using the rates below in sequential application followed by a postemergence product such as SCEPTER, SCEPTER 70DG, PURSUIT, or PURSUIT DG herbicides. Refer to the PROWL 3.3 EC leaflet label and sequential product label for all other use directions, restrictions and additional weeds controlled.

PROWL 3.3 EC Application Rates

Use the following rate table for all PROWL 3.3 EC preemergence treatments applied north of I-80. **Limitation:** Note that the maximum PROWL 3.3 EC application rate is 2.4 pints per acre for medium and fine texture soils.

Broadcast Rate per Acre of PROWL 3.3 EC Herbicide Alone or in Tank Mixture Applied Preemergence in Soybeans Grown North of I-80 in Indiana, Michigan and Ohio

Soil Texture	PROWL 3.3 EC (pints) Up to 3.0% Organic Matter*	PROWL 3.3 EC (pints) More than 3.0% Organic Matter*
COURSE	1.2 to 1.8	1.8
MEDIUM	1.8 to 2.4	1.8 to 2.4
FINE	1.8 to 2.4	2.4

^{*}NOTES: Use the higher rate of PROWL 3.3 EC if heavy weed populations are anticipated, or if hard-to-control weed species will be present, or if extensive crop residues were present following seedbed preparation (minimum or no-till systems), or if soil organic matter falls in the high end of the range. Refer to the PROWL 3.3 EC leaflet label section on Soil Texture for soil type classifications. PROWL 3.3 EC is not recommended for use on peat or muck soils.

BASF Corporation

26 Davis Drive Research Triangle Park, NC 27709



[®]Registered Trademarks of BASF

¹Butyrac is a trademark of Rhone-Poulenc Ag Company

²Lasso and Roundup are trademarks of Monsanto Company

³Gramoxone Extra is a trademark of Zeneca

⁴Sencor is a trademark of Mobay Corporation

⁵Canopy, Lexone, Lorox and Preview are trademarks of E. I. DuPont

De Nemours & Co.

⁶Dual is a trademark of CIBA

Rev. 1, 01/15/01

NVA 2001-04-129-0020

©2001 BASF Corporation All rights reserved