

USE FOR WEED CONTROL IN LAWNS AND TURF

OBSERVE ALL PRECAUTIONARY STATEMENTS IN
THE STOMP 3.3 EC LEAFLET LABEL BEFORE USING

ACCEPTED

AUG 23 1991

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 241-341

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at time of pesticide application. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

Use only on well-established turfgrass with a dense and uniform stand.

STOMP 3.3 EC treatments will not control established weeds. Applications must be made prior to germination of weeds.

Three applications per year may be made. Allow at least 2 months between applications except where indicated under RATE AND APPLICATION CHART.

MIXING AND APPLICATION INSTRUCTIONS

Add STOMP 3.3 EC herbicide slowly to partially-filled tank (1/2 to 3/4 full) of water while agitating and then fill the remainder of the tank with water. MAINTAIN CONTINUOUS AGITATION WHILE ADDING STOMP 3.3 EC AND UNTIL SPRAYING IS COMPLETED. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying as needed.

Apply with properly calibrated equipment in sufficient water per acre to provide uniform spray distribution. Low pressure (25-50 psi) sprayers are recommended. Avoid application when winds may cause drift.

•Registered Trademark of American Cyanamid Company.

COMPATIBILITY

STOMP 3.3 EC is compatible with most commonly used herbicides. When STOMP 3.3 EC is used in tank-mixture with another herbicide, refer to each label for rates, methods of application, proper timing, weeds controlled, limitations and precautions. Always use in accordance with the more restrictive label limitations and precautions. When tank mixing be sure to add STOMP 3.3 EC to the partially filled tank first, mix thoroughly, and then add combination products to the mixture. DO NOT apply tank mix combinations unless previous experience indicates the mixture is effective and will not result in application problems or plant injury. DO NOT attempt to mix STOMP 3.3 EC directly with liquid fertilizer. Premix one part of STOMP 3.3 EC with two parts water and add this diluted mixture slowly into the tank while agitating.

PRECAUTIONS

Not for use on bentgrass or Poa annua (putting greens and tees) or on dichondra, where these are desired species.

This product may cause yellowing and thinning of cool season grasses in winter overseeded turf.

Delay seeding by 4 months and sprigging by 5 months after application. Prior to seeding or sprigging, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before applying product.

On turf that has been severely thinned due to winter damage, excessive soil moisture, low temperature--below 40°F., scalping, insects, disease, etc., wait until turf has filled in and rooting of stolons is complete before applying.

This product may stain, thus avoid contact with areas such as sidewalks, driveways, etc. If contact with the spray mixture occurs, promptly rinse with water.

WEED CONTROL IN LAWNS AND TURF

STOMP 3.3 EC can be used for weed control in lawns and turf.

TURFGRASS TYPES:

STOMP 3.3 EC should only be applied to well established lawns and turf.

STOMP 3.3 EC can be used on the following turfgrasses:

Bahiagrass	Fine fescue	St. Augustinegrass
Bermudagrass	Kentucky bluegrass	Tall fescue
Centipede grass	Perennial ryegrass	Zoysiagrass

WEEDS SPECIES CONTROLLED:

When applied as directed, STOMP 3.3 EC will control the following weed species:

GRASSES CONTROLLED:

<u>Common Name</u>	<u>Scientific Name</u>
Annual bluegrass	<u>Poa annua</u>
Barnyardgrass	<u>Echinochloa crus-galli</u>
Crabgrass	<u>Digitaria</u> spp.
Fall panicum	<u>Panicum dichotomiflorum</u>
Foxtail	<u>Setaria</u> spp.
Goosegrass	<u>Eleusine indica</u>

BROADLEAF WEEDS CONTROLLED:

<u>Common Name</u>	<u>Scientific Name</u>
Common chickweed	<u>Stellaria media</u>
Cudweed	<u>Gnaphalium</u> spp.
Hop clover	<u>Trifolium procumbens</u>
Henbit	<u>Lamium amplexicaule</u>
Knotweed	<u>Polygonum aviculare</u>
Mouseear chickweed	<u>Cerastium vulgatum</u>
Prostrate spurge	<u>Euphorbia supina</u>
Purslane	<u>Portulaca oleracea</u>
Oxalis	<u>Oxalis</u> spp.
Evening primrose	<u>Oenothera biennis</u>

474

RATE AND APPLICATION CHART

<u>TURF SPECIES</u>	<u>WEEDS CONTROLLED</u>	<u>RATES OF STOMP 3.3 EC</u>	<u>COMMENTS</u>
<u>NORTHERN GRASSES:</u>			
Fine fescue Kentucky bluegrass Perennial ryegrass Tall fescue	Barnyardgrass Crabgrass Fall panicum Foxtail Hop clover Prostrate spurge Yellow woodsorrel Cudweed	3.6 to 4.8 pts per Acre or 1.3 to 1.8 oz per 1000 sq.ft.	Apply in spring before weeds germinate
	Goosegrass Evening primrose	7.2 pts per Acre or 2.6 oz. per 1000 sq.ft.	Apply in spring before weeds germinate. Repeat application if necessary in 8 weeks at 3.6 pts per Acre or 1.3 oz. per 1000 sq.ft.
	Annual bluegrass Common chickweed Mousetear chickweed Henbit	3.6 pts to 4.8 pts per Acre or 1.3 to 1.8 oz. per 1000 sq.ft.	Apply in late summer to early fall before weeds germinate
<u>SOUTHERN GRASSES:</u>			
Bahia Bermudagrass Centipedegrass St. Augustinegrass Tall fescue Zoysiagrass	Barnyardgrass Crabgrass Fall panicum Foxtail Prostrate spurge Yellow woodsorrel Cudweed	4.2 to 4.8 pts per Acre or 1.6 to 1.8 oz. per 1000 sq.ft.	Apply in spring before weeds germinate. Repeat application if necessary in 8 wks. using 3.6 pts per acre.
	Goosegrass Hop clover Evening primrose	6.4 to 7.9 pts per Acre or 2.4 to 2.9 oz. per 1000 sq.ft.	
	Henbit Annual bluegrass Common chickweed Mousetear chickweed	3.6 to 4.8 pts per Acre or 1.3 to 1.8 oz. per 1000 sq.ft.	Apply in late summer to early fall before weeds germinate