

PM 25

241-341

1 of 8

STOMP® 3.3 EC
herbicide

FOR CONTROL OF MOST ANNUAL GRASSES AND CERTAIN BROADLEAF
WEEDS IN NONCROPLAND AREAS, ORNAMENTALS, AND CHRISTMAS TREES

ACTIVE INGREDIENT:

pendimethalin, N-(1-ethylpropyl)-3,4-dimethyl-2,
6-dinitrobenzenamine..... 37.4%

INERT INGREDIENTS:..... 62.6%

TOTAL..... 100.0%
(1 gallon contains 3.3 lbs. of pendimethalin)

EPA Reg. No. 241-341

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCION

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto
hasta que la etiqueta le haya sido explicada ampliamente.

FIRST AID


If in eyes: Hold eyelids open and flush with steady, gentle stream of
water for 15 minutes. Call a physician if eye irritation
persists.

If swallowed: DO NOT induce vomiting. Call a physician or Poison Control
Center immediately.

For skin contact: Wash thoroughly with soap and water. Call a physician
if skin irritation persists.

NOTE TO PHYSICIAN: Because of increased risk of chemical pneumonia or
pulmonary edema caused by aspiration of the hydrocarbon solvent, vomiting
should be induced only under professional supervision.

AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
VEGETATION AND PEST CONTROL
WAYNE, NJ 07470 ©1991



Net Contents:

11/91

®Registered Trademark of American Cyanamid Company

298

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION**

Causes moderate eye irritation. Harmful if swallowed or absorbed through the skin. Wear goggles or face shield. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Use with adequate ventilation.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water when disposing of equipment washwaters.

DISCLAIMER

The label instructions for use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risk referred to above.

Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages.

American Cyanamid Company makes no other express or implied warranty, including other express or implied warranty of FITNESS or of MERCHANTABILITY.

389

GENERAL INFORMATION

STOMP 3.3 EC herbicide is recommended for preemergence control of grasses and certain broadleaf weed species as they germinate on noncropland areas, ornamentals, and established Christmas trees. STOMP 3.3 EC herbicide treatments are most effective in controlling weeds when adequate rainfall is received within 30 days after application.

STOMP 3.3 EC herbicide will not control established weeds. Therefore, areas to be treated should be free of established weeds at the time of treatment, or STOMP 3.3 EC herbicide may be used in conjunction with herbicides registered for postemergence use in noncropland areas, ornamentals, and established Christmas trees. Consult the labels of those herbicides for suggested treatments, rates to be used and precautions or restrictions for use in noncropland areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at time of herbicide application. DO NOT apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

STORAGE: DO NOT STORE BELOW 40°F. Extended storage at temperatures below 40°F can result in the formation of crystals on the bottom of container. If crystallization does occur, store the container on its side at room temperature (70°F) and rock occasionally until crystals redissolve.

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE

DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER

DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

4 of 8

Observe all cautions and limitations in this label and the labels of products used in combination with STOMP 3.3 EC herbicide. The use of STOMP 3.3 EC herbicide not consistent with this label can result in injury to crops, animals, or persons. Keep containers closed to avoid spills and contamination.

MIXING INSTRUCTIONS

1. Fill tank one-half to three-quarters full with clean water.
2. Add STOMP 3.3 EC herbicide to the partially filled tank while agitating and then fill the remainder of the tank with water.
3. MAINTAIN CONTINUOUS AGITATION WHILE ADDING STOMP 3.3 EC HERBICIDE AND UNTIL SPRAYING IS COMPLETED. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying.
4. If STOMP 3.3 EC herbicide is to be used in tank mixtures with other registered herbicides, then follow directions on the labels of those products which recommend tank mixing.

SPRAYING INSTRUCTIONS

Uniformly apply with properly calibrated ground equipment in sufficient water per acre to uniformly treat the area with a spray pressure of 25 to 50 psi. Maintain continuous agitation during spraying with good mechanical or bypass agitation.

NONCROPLAND WEED CONTROL

STOMP 3.3 EC herbicide is recommended for preemergence control of most annual grasses and certain broadleaf weeds as they germinate on noncropland areas such as railroad, utility and pipeline rights-of-way, highway guardrails, delineators, and sign posts, utility substations, petroleum tank farms, pumping installations, fence rows, storage areas, windbreaks, shelterbelts, and other similar areas.

For postemergence control of weeds, tank-mix combinations of STOMP 3.3 EC herbicide plus ARSENAL® herbicide are recommended. DO NOT tank mix with ARSENAL in California. Recommended rates for the tank-mix compounds should be determined from the product labels of both STOMP 3.3 EC herbicide and ARSENAL prior to use.

®Registered Trademark of American Cyanamid Company

548

WEED CONTROL IN AND AROUND ORNAMENTAL PLANTINGS

STOMP 3.3 EC herbicide is recommended for use on the following established field grown and container ornamentals, plants & trees, and established ground covers. Care must be taken that soil or planting mixes have settled firmly following transplanting and that there are no cracks that would allow direct contact of STOMP 3.3 EC herbicide and roots. Directed sprays where soil and media surfaces are uniformly covered will result in best weed control and ornamental tolerance. STOMP 3.3 EC herbicide sprays can be safely applied over-the-top of those plants listed below.

FIELD AND CONTAINER ORNAMENTALS

<u>Common Name</u>	<u>Scientific Name</u>
Azalea 'Coral Bells'	<u>Rhododendron</u>
Bamboo, Heavenly	<u>Nandina domestica</u>
Birch, River	<u>Betula nigra</u>
Bottle Brush	<u>Callistemon citrinus</u>
Boxwood, Common	<u>Buxus sempervirens</u>
Bush Honeysuckle	<u>Diervilla lonicera</u>
Common Oleander	<u>Nerium oleander</u>
Dogwood	<u>Cornus cardinal</u>
European Fan Palm	<u>Chamaerops humilis</u>
Fir, Balsam	<u>Abies balsamea</u>
Firethorn	<u>Pyracantha coccinea</u>
Holly, Dwarf Burfordi	<u>Ilex cornutanana</u>
Holly, Japanese Compacta	<u>Ilex crenata compacta</u>
Holly 'Savannah'	<u>Ilex attenuata</u>
Hosta	<u>Hosta lancifolia</u>
Juniper, Andorra	<u>Juniperus horizontalis plumosa</u>
Juniper 'Pfitzer'	<u>Juniperus chinensis</u>
Juniper 'Sea Green'	<u>Juniperus chinensis media</u>
Juniper 'Blue Chip'	<u>Juniperus horizontalis</u>
Juniper, Savins 'Broadmoor'	<u>Juniperus sabina</u>
Liriope, Green	<u>Liriope spicata</u>
Liriope, 'Monroe White'	<u>Liriope musari</u>
Liriope, Variegated	<u>Liriope muscari 'Variegata'</u>
Maple, Red	<u>Acer rubrum</u>
Mondo Grass	<u>Ophiopogon japonicum</u>
Mountain Laurel	<u>Kalmia latifolia</u>
Natal Plum	<u>Carissa grandiflora</u>
Pittosporum	<u>Pittosporum tobira</u>
Privet, Japanese	<u>Ligustrum japonicum</u>
Spirea, Bridal Wreath	<u>Spirea vanhouttei</u>

GROUND COVER ORNAMENTALS

STOMP 3.3 EC herbicide may be used for the preemergence control of the weed species listed in and around the following established ground cover plants:

<u>Common Name</u>	<u>Scientific Name</u>
Capeweed	<u>Arctotheca calendula</u>
Daisy, Trailing African	<u>Osteospermum fruticosum</u>
Gazania	<u>Gazania splendens</u>
Rose-of-Sharon	<u>Hypericum calycinum</u>
Iceplant, Large Leaf	<u>Carpobrotus edule</u>
Ivy, English	<u>Helix hedera</u>
Ivy, Geranium	<u>Pelargonium peltatum</u>
Myoporum	<u>Myoporum parviflorum</u>
Pachysandra	<u>Pachysandra terminalis</u>
Potentilla	<u>Potentilla jackmanii</u>
Potentilla	<u>Potentilla verna</u>

PRECAUTION!

Apply only to established plantings. DO NOT apply STOMP 3.3 EC herbicide to seedbeds, transplant beds, or liner beds.

WEED CONTROL IN CHRISTMAS TREE PLANTINGS

STOMP 3.3 EC herbicide is recommended for preemergence control of the weed species listed on this label as they germinate in areas planted with the following Christmas trees:

*Austrian Pine	<u>Pinus nigra</u>
*Balsam Fir	<u>Abies balsamae</u>
*Colorado Blue Spruce	<u>Picea pungens</u>
Douglas Fir	<u>Pseudotsuga menziesii</u>
*Scotch Pine	<u>Pinus sylvestris</u>
*Virginia Pine	<u>Pinus virginiana</u>
*White Fir	<u>Abies concolor</u>
*White Spruce	<u>Picea glauca</u>

*Do not use in California.

DO NOT apply STOMP 3.3 EC herbicide to Christmas tree seed or nursery beds. Apply STOMP 3.3 EC herbicide only to established Christmas tree plantings. STOMP 3.3 EC herbicide can be applied directly over the top of established Christmas tree plantings.

STOMP 3.3 EC herbicide will not control established weeds. Therefore, areas to be treated should be free of established weeds at the time of treatment, or STOMP 3.3 EC herbicide may be used in conjunction with herbicides registered for postemergence use in Christmas trees. Consult the labels of those herbicides for suggested treatments, rates to be used and precautions or restrictions for use in Christmas trees.

7-84

WEED SPECIES CONTROLLED

For preemergence control of the weed species listed, apply STOMP 3.3 EC herbicide at the following rates:

<u>Length of Control</u>	<u>STOMP 3.3 EC Rate</u>
Short Term Control (2-4 months)	2.4 Quarts/Acre
Long Term Control (6-8 months)	4.8 Quarts/Acre

STOMP 3.3 EC herbicide will not control established weeds.

The efficacy of STOMP 3.3 EC herbicide will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation.

The following grass and broadleaf weeds are controlled by preemergence treatments of STOMP 3.3 EC herbicide at the rates recommended:

GRASSES CONTROLLED

<u>Common Name</u>	<u>Scientific Name</u>
Barnyardgrass	<u>Echinochloa crus-galli</u>
Bluegrass, Annual	<u>Poa annua</u>
Crabgrass	<u>Digitaria</u> spp.
Crowfootgrass	<u>Dactyloctenium aegyptium</u>
Foxtail, Giant	<u>Setaria faberi</u>
Foxtail, Green	<u>Setaria viridis</u>
Foxtail, Yellow	<u>Setaria lutescens</u>
Goosegrass	<u>Eleusine indica</u>
Itchgrass	<u>Rottboellia exaltata</u>
Johnsongrass (from seed)	<u>Sorghum halepense</u>
Junglerice	<u>Echinochloa colonum</u>
Lovegrass	<u>Eragrostis</u> spp.
Panicum, Browntop	<u>Panicum fasciculatum</u>
Panicum, Fall	<u>Panicum dichotomiflorum</u>
Panicum, Texas	<u>Panicum texanum</u>
Sandbur, Field	<u>Cenchrus incertus</u>
Signalgrass	<u>Brachiaria platyphylla</u>
Sprangletop, Mexican	<u>Leptochloa uninervia</u>
Sprangletop, Red	<u>Leptochloa filiformis</u>
Witchgrass	<u>Panicum capillare</u>
Woolly Cupgrass	<u>Eriochloa villosa</u>

BROADLEAF WEEDS CONTROLLED

<u>Common Name</u>	<u>Scientific Name</u>
Carpetweed	<u>Mollugo verticillata</u>
Chickweed, Common	<u>Stellaria media</u>
Chickweed, Mouseear	<u>Cerastium vulgatum</u>

Common Name

Scientific Name

Clover, Hop
 Cudweed
 Fiddleneck
 Filaree
 Henbit
 Knotweed (prostrate)
 Kochia
 Lambsquarters
 Pigweed
 Puncturevine
 Purslane
 Pusley, Florida
 Rocket, London
 Sheperdspurse
 Smartweed, Pennsylvania
 Spurge, Annual
 Spurge, Prostrate
 Woodsorrel, Yellow
 Velvetleaf (Buttonweed)

Trifolium procumbens
Gnaphalium spp.
Amsinckia intermedia
Erodium spp.
Lamium amplexicaule
Polygonum aviculare
Kochia scoparia
Chenopodium album
Amaranthus spp.
Tribulus terrestris
Portulaca oleracea
Richardia scabra
Sisymbrium irio
Capsella bursa-pastoris
Polygonum pensylvanicum
Euphorbia spp.
Euphorbia matriculata
Oxalis stricta
Abutilon theophrasti