

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS**

CAUTION

Causes moderate eye irritation. Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Use with adequate ventilation.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water or to areas where surface water is present. Do not apply to intertidal areas below the mean high water mark. Oil and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water when disposing of equipment wash waters.

DISCLAIMER

The label instructions for use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein, described when used in accordance with the directions for use, subject to the risk referred to above. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages.

American Cyanamid Company makes no other express or implied warranty, including other express or implied warranty of FITNESS or MERCHANTABILITY.

GENERAL INFORMATION

STOMP 33 EC herbicide is recommended for pre-emergence control of grasses and certain broadleaf weeds species as they germinate on non-cropland areas and in established Christmas tree plantations. STOMP 33 EC herbicide treatments are most effective in controlling weeds when adequate rainfall is received within 30 days after application.

STOMP 33 EC herbicide will not control established weeds. Therefore, areas to be treated should be free of established weeds at the time of treatment. STOMP 33 EC herbicide may be used in conjunction with herbicides registered for postemergence use in non-cropland areas and in established Christmas tree plantations. Consult the labels of those herbicides for suggested treatments, rates to be used and precautions or restrictions for use in non-cropland areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at time of herbicide application. DO NOT apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

STORAGE: DO NOT STORE BELOW 40 F. Extended storage at temperatures below 40 F can result in the formation of crystals on the bottom of container. If crystallization does occur, store the container on its side at room temperature (70 F) and rock occasionally until crystals redissolve.

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or residue is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of as a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Observe all cautions and limitations in this label and the labels of products used in combination with STOMP 33 EC herbicide. The use of STOMP 33 EC herbicide not consistent with this label can result in injury to crops, animals, or persons. Keep containers closed to avoid spills and contamination.

MIXING INSTRUCTIONS

1. Fill tank one-half to three-quarters full with clean water.
2. Add STOMP 33 EC herbicide to the partially filled tank while agitating and then fill the remainder of the tank with water.
3. MAINTAIN CONTINUOUS AGITATION WHILE ADDING STOMP 33 EC HERBICIDE AND UNTIL SPRAYING IS COMPLETED. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying.
4. If STOMP 33 EC herbicide is to be used in tank mixtures with other registered herbicides, then follow directions on the labels of those products which recommend tank mixing.

SPRAYING INSTRUCTIONS

Uniformly apply with properly calibrated ground equipment in sufficient water per acre to uniformly treat the area with a spray pressure of 25 to 50 psi. Maintain continuous agitation during spraying with good mechanical or bypass agitation.

STOMP 33 EC treatments in non-cropland areas and in Christmas tree plantations are most effective in controlling weeds when adequate rainfall is received within 30 days after application.

Avoid contact of spray solution with driveways, stone, wood, or other porous surfaces. Rinse immediately to avoid staining.

NONCROPLAND WEED CONTROL

STOMP 33 EC herbicide is recommended for pre-emergence control of most annual grasses and certain broadleaf weeds as they germinate on non-cropland areas such as railroad, utility, highway, and pipeline rights-of-way, highway guardrails, delineators, and sign posts, utility substations, and shelterbelts, and for post-emergence control of weeds in non-cropland areas. STOMP 33 EC herbicide are recommended for use in conjunction with other registered herbicides in tank mixtures.

* Registered Trademark of American Cyanamid Company

posts, utility substations, and shelterbelts, and for post-emergence control of weeds in non-cropland areas. STOMP 33 EC herbicide are recommended for use in conjunction with other registered herbicides in tank mixtures.

WEED CONTROL ORNAMENTS

STOMP 33 EC herbicide should not be applied to established ornamentals planted in non-cropland areas. If application is necessary, it must be taken that soil cracks that would result in best weed control. STOMP 33 EC herbicide should not be applied over the top of

EST ORN

Common Name
Azalea Coral Bell
Bamboo Heavenly
Birch River
Bottle Brush
Bowwood Common
Bush Honeysuckle
Common Oleander
Dogwood
European Fan Palm
Fir Balsam
Fritron
Holly Dwarf Burford
Holly Japanese Compact
Holly Savannah

Common Name	Scientific Name
Hosta	<i>Hosta latifolia</i>
Juniper Andromeda	<i>Juniperus andromeda</i>
Juniper Pfitzer	<i>Juniperus pfitzeri</i>
Juniper Sea Green	<i>Juniperus horizontalis ssp. murex</i>
Juniper Blue Chip	<i>Juniperus horizontalis</i>
Juniper Savins Broadleafed	<i>Juniperus horizontalis</i>
Linum Green	<i>Linum catharticum</i>
Linum Monroe White	<i>Linum catharticum</i>
Linum Variegated	<i>Linum catharticum variegatum</i>
Maple Red	<i>Acer rubrum</i>
Mondo Grass	<i>Ophiopogon japonicum</i>
Mountain Laurel	<i>Kalmia latifolia</i>
Natal Plum	<i>Caryocarpus glaberrima</i>
Pittosporum	<i>Pittosporum tobira</i>
Privet Japanese	<i>Ligustrum japonicum</i>
Sprea Bridal Wreath	<i>Spiraea vanhouttei</i>

GROUND COVER ORNAMENTALS

Common Name	Scientific Name
Capweed	<i>Antitrochium glandulosum</i>
Daisy Training African	<i>Osteospermum fruticosum</i>
Gazania	<i>Gazania splendens</i>
Rose of Sharon	<i>Hibiscus syriacus</i>
Iceplant Large Leaf	<i>Lamprolaima</i>
Ivy English	<i>Hedera helix</i>
Ivy Geranium	<i>Geranium ivy-leafed</i>
Murderer	<i>Murderer</i>
Pachysandra	<i>Pachysandra terminalis</i>
Potentilla	<i>Potentilla germanica</i>
Potentilla	<i>Potentilla fruticosa</i>

PRECAUTION!

Apply only to established plantings

WEED CONTROL IN CHRISTMAS TREE PLANTATIONS

STOMP 33 EC herbicide is recommended for pre-emergence control of the weed species listed on the label as they germinate in areas planted with the following Christmas trees.

Common Name	Scientific Name
Australian Pine	<i>Pinus nigra</i>
Balsam Fir	<i>Abies balsamica</i>
Colorado Blue Spruce	<i>Picea pungens</i>
Douglas Fir	<i>Pseudotsuga menziesii</i>
Scotts Pine	<i>Pinus sylvestris</i>
Virginia Pine	<i>Pinus virginiana</i>
White Fir	<i>Abies concolor</i>
White Spruce	<i>Picea glauca</i>

DO NOT apply STOMP 33 EC herbicide to Christmas tree seed or nursery beds. Apply STOMP 33 EC herbicide only to established Christmas tree plantations. STOMP 33 EC herbicide can be applied directly over the top of established Christmas trees.

STOMP 33 EC herbicide will not control established weeds. Therefore, areas to be treated should be free of established weeds at the time of treatment or STOMP 33 EC herbicide may be used in conjunction with herbicides registered for post-emergence use in Christmas trees. Consult the labels of these herbicides for suggested treatments. Do not use and label directions or restrictions for use in Christmas trees.

WEED SPECIES CONTROLLED

For pre-emergence control of the weed species listed apply STOMP 33 EC herbicide at the following rates:

Length of Control	STOMP 33 EC Rate
Short Term Control (2-4 months)	2.4 Quarts/Acre (1.8 fl. oz./1000 sq. ft.)
Long Term Control (6 months)	4.8 Quarts/Acre (3.5 fl. oz./1000 sq. ft.)

STOMP 33 EC herbicide will not control established weeds.

The efficacy of STOMP 33 EC herbicide will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation.

The following grass and broadleaf weeds are controlled by pre-emergence treatment with STOMP 33 EC herbicide at the rates recommended.

GRASSES CONTROLLED

Common Name	Scientific Name
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bluegrass Annual	<i>Poa annua</i>
Crabgrass	<i>Digitaria sp.</i>
Crownbuggrass	<i>Dactyloctenium aegyptium</i>
Foxtail Green	<i>Setaria faberii</i>
Foxtail Yellow	<i>Setaria viridis</i>
Goosegrass	<i>Eleusine indica</i>
Itchgrass	<i>Rottboellia exaltata</i>
Johnsongrass (from seed)	<i>Sorghum halepense</i>
Junglerice	<i>Echinochloa colona</i>
Lovegrass	<i>Eragrostis sp.</i>
Panicum Browntop	<i>Panicum fasciculatum</i>
Panicum Fall	<i>Panicum dichotomiflorum</i>
Panicum Texas	<i>Panicum texanum</i>
Sandbar Field	<i>Cenchrus incertus</i>
Signalgrass	<i>Brachiaria platyphylla</i>
Sprangletop Mexican	<i>Leptochloa univerrna</i>
Sprangletop Red	<i>Leptochloa fallax</i>
Witchgrass	<i>Panicum capillare</i>
Woolly Cupgrass	<i>Eriochloa villosa</i>

BROADLEAF WEEDS CONTROLLED

Common Name	Scientific Name
Carpetweed	<i>Mollis verticillata</i>
Chickweed	<i>Stellaria media</i>
Chickweed Annual	<i>Centaurea virginiana</i>
Clover	<i>Trifolium pratense</i>
Cudweed	<i>Gnaphalium sp.</i>
Fiddleneck	<i>Amsinckia intermedia</i>
Flaree	<i>Erodium cicutarium</i>
Henbit	<i>Lamium amplexicaule</i>
Knotted (Prostrate)	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scopulorum</i>
Lambquarters	<i>Chenopodium album</i>
Pigweed	<i>Amaranthus spp.</i>
Puncturevine	<i>Tribulus terrestris</i>
Purslane	<i>Portulaca oleracea</i>
Pusley Florida	<i>Richardia scabra</i>
Rocket London	<i>Sisymbrium officinale</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Smartweed Pennsylvania	<i>Polygonum pennsylvanicum</i>
Spurge Annual	<i>Euphorbia spp.</i>
Spurge Prostrate	<i>Euphorbia maculata</i>
Woods net yellow	<i>Oxalis stricta</i>
Yellow (Prostrate)	<i>Achillea millefolium</i>

BEST AVAILABLE COPY

