

FEB 2 1994

Mrs. Desiree L. Little
American Cyanamid Co.
P.O. Box 400
Princeton, NJ 08540

Dear Mrs. Little:

Subject: Supplemental Label for Use on Succulent Peas,
Dry Edible Peas, Lentils, and Chickpeas in
Washington, Idaho, Oregon and Montana
Pursuit DG Herbicide
EPA Reg. No. 241-350
Your submission dated Jan. 20, 1994

The labeling referred to above, submitted in connection
with registration under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, is acceptable. A stamped
copy is enclosed for your records.

Sincerely yours,

Robert J. Taylor
Product Manager 25
Furgicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

RD:RTaylor:eja:89:305-6800

FEB 2 1991

EPA Reg. No. 241-350

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

FOR USE IN SUCCULENT PEAS, DRY EDIBLE PEAS, LENTILS, AND CHICKPEAS

241-350
PURSUIT DG can be used on succulent peas, dry edible peas (including chickpeas) and lentils in Washington (Adams, Asotin, Columbia, Garfield, Spokane, Walla Walla and Whitman counties), Idaho (Benewah, Clearwater, Idaho, Kootenai, Latah, Lewis, and Nez Perce counties), Oregon (Union, Umatilla and Wallowa counties) and Montana only.

THIS LABEL EXPIRES DECEMBER 31, 1994

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to peas and lentils. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans. Plant peas or lentils at least 1/2 inch deep to reduce risk of crop injury.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

*Registered Trademark of American Cyanamid Company

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG applications must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT DG per year.

Allow at least 30 days between application and harvest of succulent peas.

Allow at least 60 days between application and harvest of dry edible peas, chickpeas, and lentils.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

(see reverse side)

BEST AVAILABLE COPY

Apply PURSUIT DG herbicide at a broadcast rate of 1 soluble bag (7.2 ounces per 6 2/3 acres) for preplant incorporated and preemergence applications. At this broadcast rate, one soluble bag of PURSUIT DG will treat 6 2/3 acres of edible beans.

APPLICATION INSTRUCTIONS

Preplant Incorporated Applications. Apply PURSUIT DG within 1 week before planting. DO NOT incorporate deeper than 3 inches. Refer to the PURSUIT DG leaflet label for incorporation instructions.

Preemergence Applications. Apply PURSUIT DG after planting, but prior to crop emergence.

PURSUIT DG may be tank-mixed with Sencor DF¹ or Lexone DF² to assist in the control of lambsquarters or mayweed-chamomile (dogfennel). Refer to the Sencor or Lexone label for proper application rates and restrictions.

WEEDS CONTROLLED

PURSUIT DG applied preplant incorporated or preemergence will control:

Buckwheat, wild	Nightshade, Eastern black
Kochia	Nightshade, hairy
Lambsquarters, common*	Pigweed, redroot
Mustard, wild	Shepherdspurse
Nightshade, black	Thistle, Russian

*preplant incorporated applications only

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime

Chickpeas	Peanuts
Lentils	Soybeans
Peas	IMI-Corn
- Four months after PURSUIT DG application:

Barley	Rye
Edible beans (dry, lima, snap)	Wheat

- Eighteen months after PURSUIT DG application:

Alfalfa	Popcorn
Cotton	Safflower
Lettuce	Sorghum
Oats	Sweet Corn

- Twenty-six months after PURSUIT DG application:
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

- Forty months after PURSUIT DG application:
Sugar beets

In the event of a crop loss, peas, lentils or chickpeas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

**Several seed companies have tested a wide range of inbreds for sensitivity to imazethapyr soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with imazethapyr the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into field treated previously with imazethapyr shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

BEST AVAILABLE COPY

¹Sencor is a trademark of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen
²Lexone is a trademark of E. I. duPont de Nemours and Company, Inc.