



Supplemental Labeling

PM 25
241-350
4/28

EPA Reg. No. 241-350

FOR USE IN SOYBEANS

TANK-MIXTURE OR SEQUENTIAL APPLICATIONS OF PURSUIT DG HERBICIDE AND STATUS¹ HERBICIDE FOR ENHANCED WEED CONTROL IN SOYBEANS

OBSERVE ALL PRECAUTIONARY AND WORKER PROTECTION STATEMENTS IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

STATUS herbicide may be tankmixed with PURSUIT DG for enhanced control of several broadleaf weeds. Additionally, STATUS may be applied sequentially, following a PURSUIT DG application. STATUS herbicide provides fast burn-down of susceptible broadleaf weeds. Temporary bronzing of the soybean leaves soon after application may occur. Soybeans soon outgrow early symptoms; yield will not be effected by adding STATUS to PURSUIT DG. In some cases the activity of PURSUIT DG may be reduced when mixed with STATUS. The reduction in activity may be overcome by delaying the application of STATUS herbicide 7 days following the application of PURSUIT DG.

WEEDS CONTROLLED

The addition of STATUS to PURSUIT DG at the recommended rates will enhance the control of several broadleaf weeds, including common and giant ragweed, pigweed species and waterhemp. (Refer to the STATUS label for additional weeds controlled).

APPLICATION RATES

PURSUIT DG:

Apply PURSUIT DG at the broadcast rate of 1.44 ounces per acre (1 soluble bag treats 5 acres). Refer to the following tables for the recommended rate of STATUS herbicide.

ACCEPTED
with COMMENTS
In EPA Letter Dated
MAR 21 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241-350

STATUS TANK-MIXTURE APPLICATION RATES

When tank-mixing STATUS with PURSUIT DG, apply STATUS at the following rates, depending on weed size:

Weeds/Weed Size	STATUS Rate (ounces per acre):*		
	8 - 10 oz.	12 - 14 oz.	16 - 20 oz.
Common ragweed Pigweed species, Waterhemp, tall common	1 - 4"	4 - 6"	6 - 8"
Giant Ragweed		1 - 6"	6 - 8"***

*Use the higher rate if common ragweed is present or the weed population is high.
 **Use the 24 ounce/acre rate if giant ragweed is 6 - 8 inches tall.

STATUS SEQUENTIAL APPLICATION RATES

When applying STATUS following a PURSUIT DG application (sequential), apply STATUS at the following rates:

Weeds/Weed Size	STATUS Rate (ounces per acre):*		
	10 - 12 oz.	14 - 16 oz.	18 - 24 oz.
Common ragweed Pigweed species, Waterhemp, tall common	1 - 4"	4 - 6"	6 - 8"
Giant Ragweed		1 - 6"	6 - 8"***

*Use the higher rate if common ragweed is present or the weed population is high.
 **Use the 24 ounce/acre rate if giant ragweed is 6 - 8 inches tall.

3/128

MIXING INSTRUCTIONS

POSTEMERGENCE APPLICATIONS OF PURSUIT DG AND STATUS REQUIRE THE ADDITION OF AN ADJUVANT AND A LIQUID FERTILIZER SOLUTION.

ADJUVANTS

- I. **SURFACTANTS:** Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 quart per 100 gallons.

OR

CROP OIL CONCENTRATE: Instead of a surfactant, a petroleum or vegetable seed based oil concentrate (such as SUN-IT II²) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other methylated seed oils at the rate of 1.5 to 2 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre. A non-ionic surfactant is recommended when crop response is of concern (i.e., under conditions of high relative humidity or the presence of dew on the crop leaves).

AND

II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs. per acre.

This label must be in the possession of the user at the time of herbicide application.

Refer to the STATUS label for additional information regarding application rates, restrictions, precautions, weeds controlled, adjuvants recommended and other information. **When tank-mixing herbicides, always follow the most restrictive label.**


¹Trademark of BASF Corporation

²Trademark of Agsco, Inc.

©Registered Trademark of American Cyanamid Company

PE-xxxxx 1/96 SD

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996





Supplemental Labeling

4/121

EPA Reg. No. 241-350
FOR USE IN SOYBEANS

NOT REVIEWED

In Accordance with PR Notice 82-2

Based On Draft Labeling Dated 3-21-96

TANK-MIXTURE OR SEQUENTIAL APPLICATIONS OF PURSUIT DG HERBICIDE AND STATUS¹ HERBICIDE FOR ENHANCED WEED CONTROL IN SOYBEANS

OBSERVE ALL PRECAUTIONARY AND WORKER PROTECTION STATEMENTS IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

STATUS herbicide may be tankmixed with PURSUIT DG for enhanced control of several broadleaf weeds. Additionally, STATUS may be applied sequentially, following a PURSUIT DG application. STATUS herbicide provides fast burn-down of susceptible broadleaf weeds. Temporary bronzing of the soybean leaves soon after application may occur. Soybeans soon outgrow early symptoms; yield will not be effected by adding STATUS to PURSUIT DG. In some cases the activity of PURSUIT DG may be reduced when mixed with STATUS. The reduction in activity may be overcome by delaying the application of STATUS herbicide 7 days following the application of PURSUIT DG.

WEEDS CONTROLLED

The addition of STATUS to PURSUIT DG at the recommended rates will enhance the control of several broadleaf weeds, including common and giant ragweed, pigweed species and waterhemp. (Refer to the STATUS label for additional weeds controlled).

APPLICATION RATES

PURSUIT DG:

Apply PURSUIT DG at the broadcast rate of 1.44 ounces per acre (1 soluble bag treats 5 acres). Refer to the following tables for the recommended rate of STATUS herbicide.

STATUS TANK-MIXTURE APPLICATION RATES

When tank-mixing STATUS with PURSUIT DG, apply STATUS at the following rates, depending on weed size:

Weeds/Weed Size	STATUS Rate (ounces per acre):*		
	8 - 10 oz.	12 - 14 oz.	16 - 20 oz.
Common ragweed Pigweed species, Waterhemp, tall common	1 - 4"	4 - 6"	6 - 8"
Giant Ragweed		1 - 6"	6 - 8"***

*Use the higher rate if common ragweed is present or the weed population is high.

**Use the 24 ounce/acre rate if giant ragweed is 6 - 8 inches tall.

STATUS SEQUENTIAL APPLICATION RATES

When applying STATUS following a PURSUIT DG application (sequential), apply STATUS at the following rates:

Weeds/Weed Size	STATUS Rate (ounces per acre):*		
	10 - 12 oz.	14 - 16 oz.	18 - 24 oz.
Common ragweed Pigweed species, Waterhemp, tall common	1 - 4"	4 - 6"	6 - 8"
Giant Ragweed		1 - 6"	6 - 8"***

*Use the higher rate if common ragweed is present or the weed population is high.

**Use the 24ounce/acre rate if giant ragweed is 6 - 8 inches tall.



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996



Supplemental Labeling

8/128

EPA Reg. No. 241-350

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, AND SOUTHERN AND ENGLISH PEAS

REFER TO THE MAP ON PAGE 6 FOR GEOGRAPHICAL USE AREA

READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash¹.

ACCEPTED
with COMMENTS
In EPA Letter Dated
MAR 21 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241-350

ACCEPTED
In EPA Letter Dated

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

¹Dash is a trademark of BASF Corporation

*Registered Trademark of American Cyanamid Company

DO NOT make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

USE RATE

0.72, 1.08 OR 1.44 OUNCES PER ACRE

Depending on soil type and edible bean or pea type, apply PURSUIT DG at a broadcast rate according to the following table.

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
0.72	10 acres
1.08	6-2/3 acres
1.44	5 acres

LIMA BEANS

DO NOT apply PURSUIT DG after June 30.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

NAVY, GREAT NORTHERN, RED KIDNEY, CRANBERRY, BLACK TURTLE, AND SMALL WHITE TYPE DRY BEANS

DO NOT apply to Domino variety black turtle beans.

DO NOT apply PURSUIT DG to sand or loamy sand soils.

DO NOT apply PURSUIT DG after June 20.

Allow at least 60 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

Soil Type	Use Rate (oz/acre)
Sands, loamy sands	DO NOT USE
Sandy loams and all other soils with ≤2% O.M.	0.72
All other soils with >2% O.M.	1.08

Preemergence Applications. Apply PURSUIT DG at the rates shown in the preceding table immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ENGLISH PEAS

DO NOT apply PURSUIT DG to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT DG to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. **DO NOT** apply PURSUIT DG if English peas have been treated with trifluralin.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. **DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.** If English peas have been treated with trifluralin, PURSUIT DG may be applied **ONLY** at 0.72 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT DG application may increase the likelihood and severity of crop injury.

SOUTHERN PEAS

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.44 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.44 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

Early Postemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.44 ounces per acre when peas are at least 3 inches in height. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence, or early postemergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

WEEDS CONTROLLED

PURSUIT DG applied to low organic matter soils at the broadcast rate of 0.72 ounces per acre preplant incorporated or preemergence will control light to moderate infestations of:

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

Postemergence applications of 1.08 ounces per acre must be made to weeds less than 2 inches tall for best results.

When applied as directed at the broadcast rate of 1.44 ounces per acre, PURSUIT DG will control or reduce competition from the weeds listed below:

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence.)

BROADLEAF WEEDS

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Anoda, spurred	C	2	1-2
Artichoke, Jerusalem		8	6-10
Buffalobur	C**		
Bristly starbur		2	1-2
Carpetweed	C		
Cocklebur, common	C**	8	1-8
Galinsoga	C		
Jimsonweed	C*	4	1-3
Kochia	C	4	1-3
Lambsquarters	C*	R	1-2
Mallow, Venice	R		
Momingglory			
entireleaf	R	2	1-2
ivyleaf	R	2	1-2
pitted	R	2	1-2
smallflower	C	4	1-3
tall	R	2	1-2
Mustard sp.	C	4	1-3
Nightshade,			
black	C	4	1-3
Eastern black	C	4	1-3
hairy	C	4	1-3
Pigweed,			
redroot	C	8	1-8
smooth	C	8	1-8
spiny	C	8	1-8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C*		

11/128

BROADLEAF WEEDS (Continued)

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Ragweed,			
common	R	4	1-3
giant	R	4	1-3
Sage, barnyard		R	1-3
Smartweed,			
ladysthumb	C	4	1-3
Pennsylvania	C	4	1-3
Spurge,			
prostrate	C	4	1-3
spotted	C	4	1-3
Sunflower, common	C*	4	1-3
Velvetleaf	C*	4	1-3
Thistle, Canada		R	1-3

*When soil applied, common lambsquarters, jimsonweed, prickly sida, velvetleaf and common sunflower are more consistently controlled by preplant incorporated treatments.

**Use soil applications for light to moderate infestations only. Must be preplant incorporated for best results.

GRASS WEEDS*

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Barnyardgrass	R	3	1-3
Crabgrass,			
large	R	3	1-3
smooth	R	3	1-3
Cupgrass, woolly		3**	1-3
Foxtail,			
giant	C	6	1-6
green	C	3	1-3
robust purple	C	3	1-3
robust white	C	3	1-3
yellow	C	3	1-3
Goosegrass	R		
Johnsongrass,			
seedling	C	6	1-8
rhizome		R	1-8
Panicum,			
fall	R		
Texas	R		

*When soil applied to grasses, more consistent control can be obtained from preplant incorporated treatments.

**PURSUIT DG controls emerged woolly cupgrass only.

12/128

GRASS WEEDS (Continued)

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Red rice		3	1 - 3
Shattercane	R	6	1 - 8
Signalgrass, broadleaf	R	4	1 - 8

SEDGES

Nutsedge, purple	R	R	1 - 3
yellow	R	R	1 - 3

DO NOT count cotyledon leaves when determining weed stage of growth.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - Peanuts
 - IMI-Corn™
 - Edible bean and pea types listed on this label (Refer to heading on page 1)
2. Two months after PURSUIT DG application:
 - Snap beans (DE, MD, PA and VA only)
3. Four months after PURSUIT DG application:
 - Barley
 - Edible bean and pea types not listed on this label (Refer to heading on page 1)
 - Rye
 - Wheat
4. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
5. Nine and one-half months after PURSUIT DG application:
 - Tobacco
6. Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet Corn
Popcorn	

7. Twenty-six months after PURSUIT DG application: Potatoes

8. Forty months after PURSUIT DG application:
 All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

ROTATIONAL CROP GUIDELINE (Continued)

In the event of a crop loss, the bean or pea crop may be replanted.


DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

USE AREA MAP

PURSUIT DG can be applied to the edible legume vegetables listed only in the states or parts of states shaded in the following map:


Use area includes: South Dakota east of U.S. 81, Nebraska east of U.S. 81, Kansas east of U.S. 81, Maryland and Virginia EXCLUDING THE DELMARVA PENINSULA, and other states or portions of states shaded in the following map.



PE-17242 Rev. 2 1/13/96 SD



American Cyanamid Company
 Agricultural Products Division
 Crop Protection Chemicals Department
 Wayne, NJ 07470 ©1996





Supplemental Labeling

NOT REVIEWED

In Accordance with PR Notice 82-2
Based On Draft Labeling Dated 3-21-96 EPA Reg. No. 241-350

14/128

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, AND SOUTHERN AND ENGLISH PEAS

REFER TO THE MAP ON PAGE 6 FOR GEOGRAPHICAL USE AREA

READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash¹.

¹Dash is a trademark of BASF Corporation

®Registered Trademark of American Cyanamid Company

Early Postemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.44 ounces per acre when peas are at least 3 inches in height. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

WEEDS CONTROLLED

PURSUIT DG applied to low organic matter soils at the broadcast rate of 0.72 ounces per acre preplant incorporated or preemergence will control light to moderate infestations of:

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence, or early postemergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

Postemergence applications of 1.08 ounces per acre must be made to weeds less than 2 inches tall for best results.

When applied as directed at the broadcast rate of 1.44 ounces per acre, PURSUIT DG will control or reduce competition from the weeds listed below:

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence.)

BROADLEAF WEEDS

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Anoda, spurred	C	2	1-2
Artichoke, Jerusalem		8	6-10
Buffalobur	C**		
Bristly starbur		2	1-2
Carpetweed	C		
Cocklebur, common	C**	8	1-8
Galinsoga	C		
Jimsonweed	C*	4	1-3
Kochia	C	4	1-3
Lambsquarters	C*	R	1-2
Mallow, Venice	R		
Momingglory			
entireleaf	R	2	1-2
ivyleaf	R	2	1-2
pitted	R	2	1-2
smallflower	C	4	1-3
tall	R	2	1-2
Mustard sp.	C	4	1-3
Nightshade,			
black	C	4	1-3
Eastern black	C	4	1-3
hairy	C	4	1-3
Pigweed,			
redroot	C	8	1-8
smooth	C	8	1-8
spiny	C	8	1-8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C*		

18/128

GRASS WEEDS (Continued)

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Red rice		3	1-3
Shattercane	R	6	1-8
Signalgrass, broadleaf	R	4	1-8
SEDGES			
Nutsedge,			
purple	R	R	1-3
yellow	R	R	1-3

DO NOT count cotyledon leaves when determining weed stage of growth.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - Peanuts
 - IMI-Corn™
 - Edible bean and pea types listed on this label (Refer to heading on page 1)
2. Two months after PURSUIT DG application:
 - Snap beans (DE, MD, PA and VA only)
3. Four months after PURSUIT DG application:
 - Barley
 - Edible bean and pea types not listed on this label (Refer to heading on page 1)
 - Rye
 - Wheat
4. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
5. Nine and one-half months after PURSUIT DG application:
 - Tobacco
6. Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet Corn
Popcorn	


7. Twenty-six months after PURSUIT DG application:
 - Potatoes

8. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.



19/128

ROTATIONAL CROP GUIDELINE (Continued)

In the event of a crop loss, the bean or pea crop may be replanted.


DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

USE AREA MAP

PURSUIT DG can be applied to the edible legume vegetables listed only in the states or parts of states shaded in the following map:

Use area includes: South Dakota east of U.S. 81, Nebraska east of U.S. 81, Kansas east of U.S. 81, Maryland and Virginia EXCLUDING THE DELMARVA PENINSULA, and other states or portions of states shaded in the following map.



CYANAMID
 American Cyanamid Company
 Agricultural Products Division
 Crop Protection Chemicals Department
 Wayne, NJ 07470 ©1996

PE-17242 Rev. 2 1/13/96 SD



Supplemental Labeling

20/128

EPA Reg. No. 241-350

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, SMALL WHITE TYPE DRY BEANS, LIMA BEANS, SOUTHERN AND ENGLISH PEAS, AND WHITE LUPINS

FOR DISTRIBUTION AND USE IN WISCONSIN ONLY

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended:

- Two months after PURSUIT DG application:
Snap beans

ACCEPTED
WITH COMMENTS
In EPA Letter Dated

MAR 21 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

241-350

Registered Trademark of American Cyanamid Company
PE-17276 Rev. 1 1/23/96 SD



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996



Supplemental Labeling

21/128

EPA Reg. No. 241-350

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, SMALL WHITE TYPE DRY BEANS, LIMA BEANS, SOUTHERN AND ENGLISH PEAS, AND WHITE LUPINS

FOR DISTRIBUTION AND USE IN WISCONSIN ONLY
NOT REVIEWED
Based On Draft Labeling Dated 3-21-96
With PR Notice 82-2
THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended:

- Two months after PURSUIT DG application:
Snap beans



®Registered Trademark of American Cyanamid Company
PE-17276 Rev. 1 1/23/96 SD





Supplemental Labeling

22/128

EPA Reg. No. 241-350

FOR USE IN LIMA BEANS ONLY

FOR DISTRIBUTION AND USE ONLY ON THE DELMARVA PENINSULA OF MARYLAND, DELAWARE, AND VIRGINIA

READ LABEL CAREFULLY

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 21 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

241-350

*Registered Trademark of American Cyanamid Company

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply PURSUIT DG after June 30.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
0.72	10 acres
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied to low organic matter soils at the broadcast rate of 0.72 ounces per acre preplant incorporated or preemergence will control light to moderate infestations of:

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime
 - Soybeans
 - Peanuts
 - IMI-Corn™
 - Lima beans
2. Two months after PURSUIT DG application:
 - Snap beans
3. Four months after PURSUIT DG application:
 - Barley
 - Other types of edible beans and peas
 - Rye
 - Wheat
4. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
5. Nine and one-half months after PURSUIT DG application:
 - Tobacco

™Trademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

6. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet corn |
| Popcorn | |

7. Twenty-six months after PURSUIT DG application:

- Potatoes

8. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

In the event of a crop loss, the bean or pea crop may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.





Supplemental Labeling

25/128

EPA Reg. No. 241-350

NOT REVIEWED

In Accordance with PR Notice 82-2

FOR USE IN LIMA BEANS ONLY

Filed On Draft Labeling Dated 3-21-96

FOR DISTRIBUTION AND USE ONLY ON THE DELMARVA PENINSULA OF
MARYLAND, DELAWARE, AND VIRGINIA

READ LABEL CAREFULLY

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

26/128

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply PURSUIT DG after June 30.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
0.72	10 acres
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied to low organic matter soils at the broadcast rate of 0.72 ounces per acre preplant incorporated or preemergence will control light to moderate infestations of:

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime
 - Soybeans
 - Peanuts
 - IMI-Corn™
 - Lima beans
2. Two months after PURSUIT DG application:
 - Snap beans
3. Four months after PURSUIT DG application:
 - Barley
 - Other types of edible beans and peas
 - Rye
 - Wheat
4. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
5. Nine and one-half months after PURSUIT DG application:
 - Tobacco

™Trademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

6. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet corn |
| Popcorn | |

7. Twenty-six months after PURSUIT DG application:

- Potatoes

8. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

*** Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.**

In the event of a crop loss, the bean or pea crop may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.





Supplemental Labeling

29/128

EPA Reg. No. 241-350

FOR USE IN GREAT NORTHERN, RED KIDNEY AND PINTO TYPE DRY BEANS

FOR DISTRIBUTION AND USE ONLY IN NEW MEXICO

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

with COMMENTS:
In EPA Letter Dated

MAR 21 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

241-350

*Registered Trademark of American Cyanamid Company

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after June 30. Allow at least 60 days between application and harvest.

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. **DO NOT** apply PURSUIT DG on sand or loamy sand soils.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preemergence or early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Soybeans
 - Peanuts
 - Edible bean types listed on this label (Refer to heading on page 1).
 - IMI-Corn™
- Four months after PURSUIT DG application:
 - Barley
 - Edible bean types not listed on this label (Refer to heading on page 1).
 - Peas
 - Rye
 - Wheat
- Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
- Nine and one-half months after PURSUIT DG application:
 - Tobacco

¹Dash is a trademark of BASF Corporation
™Trademark American Cyanamid Company



52/128

Supplemental Labeling

EPA Reg. No. 241-350

FOR USE IN GREAT NORTHERN, RED KIDNEY AND PINTO TYPE DRY BEANS

FOR DISTRIBUTION AND USE ONLY IN NEW MEXICO

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

NOT REVIEWED

In Accordance with PR Notice 82-2
Based On Draft Labeling Dated 3-21-96

*Registered Trademark of American Cyanamid Company

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after June 30. Allow at least 60 days between application and harvest.

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. **DO NOT** apply PURSUIT DG on sand or loamy sand soils.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preemergence or early postemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Soybeans
 - Peanuts
 - Edible bean types listed on this label (Refer to heading on page 1).
 - IMI-Corn™
- Four months after PURSUIT DG application:
 - Barley
 - Edible bean types not listed on this label (Refer to heading on page 1).
 - Peas
 - Rye
 - Wheat
- Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
- Nine and one-half months after PURSUIT DG application:
 - Tobacco

¹Dash is a trademark of BASF Corporation
™Trademark American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet corn |
| Popcorn | |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss Great Northern, red kidney or pinto beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

35/128





Supplemental Labeling

EPA Reg. No. 241-350

36/128

**FOR USE IN RED KIDNEY TYPE DRY BEANS, WHITE LUPINS,
LIMA BEANS, AND ENGLISH PEAS ONLY**

FOR DISTRIBUTION AND USE ONLY IN ILLINOIS, IOWA, AND WISCONSIN

READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 21 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241350

*Registered Trademark of American Cyanamid Company

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

RED KIDNEY TYPE DRY BEANS, WHITE LUPINS AND LIMA BEANS

DO NOT apply PURSUIT DG after June 30.

DO NOT apply PURSUIT DG to white lupins grown on sand or loamy sand soils.

Allow at least 30 days between application and harvest of lima beans.

Allow at least 60 days between application and harvest of red kidney beans or white lupins.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ENGLISH PEAS

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT DG to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. **DO NOT** apply PURSUIT DG if English peas have been treated with trifluralin.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT. For optimum weed control and crop safety, apply PURSUIT DG prior to 5 nodes before flowering. If English peas have been treated with trifluralin, PURSUIT DG may be applied ONLY at 0.72 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT DG application may increase the likelihood and severity of crop injury.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence, or early postemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Soybeans
 - Peanuts
 - Edible bean and pea types listed on this label (Refer to heading on page 1)
 - IMI-Corn™

¹Dash is a trademark of BASF Corporation
™Trademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

- 2. Four months after PURSUIT DG application:
 - Barley
 - Edible bean and pea types not listed on this label (Refer to heading on page 1).
 - Rye
 - Wheat

- 3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*

- 4. Nine and one-half months after PURSUIT DG application:
 - Tobacco

- 5. Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	

- 6. Twenty-six months after PURSUIT DG application:
 - Potatoes

- 7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, red kidney beans, white lupins, lima beans, or English peas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.





Supplemental Labeling

EPA Reg. No. 241-350

**FOR USE IN RED KIDNEY TYPE DRY BEANS, WHITE LUPINS,
LIMA BEANS, AND ENGLISH PEAS ONLY**

FOR DISTRIBUTION AND USE ONLY IN ILLINOIS, IOWA, AND WISCONSIN

READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

NOT REVIEWED

In Accordance with PR Notice 82-2
Based On Draft Labeling Dated 3-21-96

*Registered Trademark of American Cyanamid Company

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

RED KIDNEY TYPE DRY BEANS, WHITE LUPINS AND LIMA BEANS

DO NOT apply PURSUIT DG after June 30.

DO NOT apply PURSUIT DG to white lupins grown on sand or loamy sand soils.

Allow at least 30 days between application and harvest of lima beans.

Allow at least 60 days between application and harvest of red kidney beans or white lupins.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ENGLISH PEAS

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT DG to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. **DO NOT** apply PURSUIT DG if English peas have been treated with trifluralin.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT. For optimum weed control and crop safety, apply PURSUIT DG prior to 5 nodes before flowering. If English peas have been treated with trifluralin, PURSUIT DG may be applied **ONLY** at 0.72 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT DG application may increase the likelihood and severity of crop injury.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence, or early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Soybeans
 - Peanuts
 - Edible bean and pea types listed on this label (Refer to heading on page 1).
 - IMI-Corn™

¹Dash is a trademark of BASF Corporation

™Trademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

2. Four months after PURSUIT DG application:

- Barley
- Edible bean and pea types not listed on this label (Refer to heading on page 1).
- Rye
- Wheat

3. Eight and one-half months after PURSUIT DG application:

- Field corn
- Field corn grown for seed*

4. Nine and one-half months after PURSUIT DG application:

- Tobacco

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet corn |
| Popcorn | |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, red kidney beans, white lupins, lima beans, or English peas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

42/128





Supplemental Labeling

ACCEPTED
with COMMENTS
in EPA Letter Dated

MAR 21 1996

EPA Reg. No. 241-350

43/128

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
241-350

FOR USE IN ENGLISH PEAS ONLY

**FOR DISTRIBUTION AND USE ONLY ON THE DELMARVA PENINSULA OF
MARYLAND, DELAWARE, AND VIRGINIA**

READ LABEL CAREFULLY

THIS LABEL EXPIRES APRIL 30, 1997

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT DG per year.

*Registered Trademark of American Cyanamid Company

(see reverse side)

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply PURSUIT DG to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT DG to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. **DO NOT** apply PURSUIT DG if English peas have been treated with trifluralin.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Soybeans
 - Peanuts
 - IMI-Corn™
 - English peas and lima beans
- Two months after PURSUIT DG application:
 - Snap beans
- Four months after PURSUIT DG application:
 - Barley
 - Other types of edible beans and peas
 - Rye
 - Wheat

- Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
- Nine and one-half months after PURSUIT DG application:
 - Tobacco
- Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	
- Twenty-six months after PURSUIT DG application:
 - Potatoes
- Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

In the event of a crop loss, the bean or pea crop may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



™Trademark of American Cyanamid Company
PE-17282 Rev. 1 1/22/96 SD



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996



Supplemental Labeling

45/128

EPA Reg. No. 241-350

NOT REVIEWED

In Accordance with PR Notice 82-2

FOR USE IN ENGLISH PEAS ONLY Draft Labeling Dated 3-21-96

FOR DISTRIBUTION AND USE ONLY ON THE DELMARVA PENINSULA OF MARYLAND, DELAWARE, AND VIRGINIA

READ LABEL CAREFULLY

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT DG per year.

®Registered Trademark of American Cyanamid Company

(see reverse side)

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply PURSUIT DG to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. DO NOT apply PURSUIT DG to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. DO NOT apply PURSUIT DG if English peas have been treated with trifluralin.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Soybeans
 - Peanuts
 - IMI-Corn™
 - English peas and lima beans
- Two months after PURSUIT DG application:
 - Snap beans
- Four months after PURSUIT DG application:
 - Barley
 - Other types of edible beans and peas
 - Rye
 - Wheat

- Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
- Nine and one-half months after PURSUIT DG application:
 - Tobacco
- Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	
- Twenty-six months after PURSUIT DG application:
 - Potatoes

4. Forty months after PURSUIT DG application: All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

In the event of a crop loss, the bean or pea crop may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

™Trademark of American Cyanamid Company
PE-17282 Rev. 1 1/22/96 SD





Supplemental Labeling

47/128

EPA Reg. No. 241-350

FOR PREEMERGENCE USE IN SNAP BEANS GROWN IN NEW JERSEY ONLY

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to snap beans. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

*Registered Trademark of American Cyanamid Company

(see reverse side)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG applications must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT apply PURSUIT DG after June 20.

Allow at least 30 days between application and harvest.

DO NOT make more than one application of PURSUIT DG per year.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

ACCEPTED
with conditions
In EPA Letter Dated

MAR 21 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

241-350

APPLICATION INSTRUCTIONS

FOR USE IN THE STATE OF NEW JERSEY ONLY

Preemergence Applications. For suppression of pigweed, nightshade and common purslane, apply PURSUIT DG at the broadcast rate of 0.54 ounces per acre (one ECO-PAK soluble bag treats 13-1/3 acres) immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - Peanuts
 - Edible beans and peas
 - IMI-Corn™
2. Four months after PURSUIT DG application:
 - Barley
 - Rye
 - Wheat
3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
4. Nine and one-half months after PURSUIT DG application:
 - Tobacco
5. Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	
6. Twenty-six months after PURSUIT DG application:
 - Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, snap beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

™Trademark of American Cyanamid Company
PE-17275 Rev.1 1/23/96 SD



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996



Supplemental Labeling

49/128

EPA Reg. No. 241-350

FOR PREEMERGENCE USE IN SNAP BEANS GROWN IN NEW JERSEY ONLY

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

NOT REVIEWED

In Accordance with PR Notice 82-2

Based On Draft Labeling Dated 3.21.96

GENERAL INFORMATION

DIRECTIONS FOR USE

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to snap beans. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG applications must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT apply PURSUIT DG after June 20.

Allow at least 30 days between application and harvest.

DO NOT make more than one application of PURSUIT DG per year.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

*Registered Trademark of American Cyanamid Company

(see reverse side)

APPLICATION INSTRUCTIONS

FOR USE IN THE STATE OF NEW JERSEY ONLY

Preemergence Applications. For suppression of pigweed, nightshade and common purslane, apply PURSUIT DG at the broadcast rate of 0.54 ounces per acre (one ECO-PAK soluble bag treats 13-1/3 acres) immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - Peanuts
 - Edible beans and peas
 - IMI-Corn™
2. Four months after PURSUIT DG application:
 - Barley
 - Rye
 - Wheat
3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
4. Nine and one-half months after PURSUIT DG application:
 - Tobacco
5. Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	

6. Twenty-six months after PURSUIT DG application:

Potatoes

.....

..

.....

.....

.....

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, snap beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

™Trademark of American Cyanamid Company
PE-17275 Rev. 1/23/96 SD





Supplemental Labeling

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 21 1996

EPA Reg. No. 241-350

51/128

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 241-350

FOR USE IN SUCCULENT PEAS, DRY EDIBLE PEAS, LENTILS, CHICKPEAS, AND LIMA BEANS

PURSUIT DG can be used on succulent peas, dry edible peas (including chickpeas) and lentils in Washington (Adams, Asotin, Columbia, Garfield, Spokane, Walla Walla and Whitman counties), Idaho (Benewah, Clearwater, Idaho, Kootenai, Latah, Lewis, and Nez Perce counties), Oregon (Union, Umatilla and Wallowa counties) and Montana only.

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans. Plant peas, lentils, or lima beans at least 1/2 inch deep to reduce risk of crop injury.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT DG per year.

Allow at least 30 days between application and harvest of succulent peas and lima beans.

Allow at least 60 days between application and harvest of dry edible peas, chickpeas, and lentils.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

*Registered Trademark of American Cyanamid Company

(see reverse side)

USE RATE

(ECO-PAK soluble bag treats 6 2/3 acres which is equivalent to 1.1 ounces per acre)

Apply PURSUIT DG herbicide at a broadcast rate of 1 soluble bag (7.2 ounces per 6 2/3 acres) for preplant incorporated and preemergence applications. At this broadcast rate, one soluble bag of PURSUIT DG will treat 6 2/3 acres of edible beans.

APPLICATION INSTRUCTIONS

Preplant Incorporated Applications. Apply PURSUIT DG within 1 week before planting. DO NOT incorporate deeper than 3 inches. Refer to the PURSUIT DG leaflet label for incorporation instructions.

Preemergence Applications. Apply PURSUIT DG after planting, but prior to crop emergence.

PURSUIT DG may be tank-mixed with Sencor DF¹ of Lexone DF² to assist in the control of lambsquarters or mayweed-chamomile (dogfennel). Refer to the Sencor or Lexone label for proper application rates and restrictions.

WEEDS CONTROLLED

PURSUIT DG applied preplant incorporated or preemergence will control:

- | | |
|------------------------|---------------------------|
| Buckwheat, wild | Nightshade, Eastern black |
| Kochia | Nightshade, hairy |
| Lambsquarters, common* | Pigweed, redroot |
| Mustard, wild | Shepherdspurse |
| Nightshade, black | Thistle, Russian |

*preplant incorporated applications only

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:

Chickpeas	Peanuts
Lentils	Soybeans
Peas	IMI-Corn™
Lima beans	
- Four months after PURSUIT DG application:
 - Barley
 - Dry edible beans
 - Rye
 - Snap beans (3 months)
 - Wheat
- Nine and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed**
 - Tobacco

¹Sencor is a trademark of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen
²Lexone is a trademark of E. I. duPont de Nemours and Company, Inc.
 ™Trademark of American Cyanamid Company
 PE-17228 Rev. 2 1/11/96 SD

- Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet Corn
Popcorn	

- Twenty-six months after PURSUIT DG application:
 - Potatoes

- Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, peas, lentils, chickpeas, or lima beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

** Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996



Supplemental Labeling

53/12

EPA Reg. No. 241-350

NOT REVIEWED

In Accordance with PR Notice 82-2

Based On Draft Labeling Dated 3-21-96

FOR USE IN SUCCULENT PEAS, DRY EDIBLE PEAS, LENTILS, CHICKPEAS, AND LIMA BEANS

PURSUIT DG can be used on succulent peas, dry edible peas (including chickpeas) and lentils in Washington (Adams, Asotin, Columbia, Garfield, Spokane, Walla Walla and Whitman counties), Idaho (Benewah, Clearwater, Idaho, Kootenai, Latah, Lewis, and Nez Perce counties), Oregon (Union, Umatilla and Wallowa counties) and Montana only.

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans. Plant peas, lentils, or lima beans at least 1/2 inch deep to reduce risk of crop injury.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT DG per year.

Allow at least 30 days between application and harvest of succulent peas and lima beans.

Allow at least 60 days between application and harvest of dry edible peas, chickpeas, and lentils.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

USE RATE

(ECO-PAK soluble bag treats 6 2/3 acres which is equivalent to 1.1 ounces per acre)

Apply PURSUIT DG herbicide at a broadcast rate of 1 soluble bag (7.2 ounces per 6 2/3 acres) for preplant incorporated and preemergence applications. At this broadcast rate, one soluble bag of PURSUIT DG will treat 6 2/3 acres of edible beans.

APPLICATION INSTRUCTIONS

Preplant Incorporated Applications. Apply PURSUIT DG within 1 week before planting. DO NOT incorporate deeper than 3 inches. Refer to the PURSUIT DG leaflet label for incorporation instructions.

Preemergence Applications. Apply PURSUIT DG after planting, but prior to crop emergence.

PURSUIT DG may be tank-mixed with Sencor DF¹ or Lexone DF² to assist in the control of lambsquarters or mayweed-chamomile (dogfennel). Refer to the Sencor or Lexone label for proper application rates and restrictions.

WEEDS CONTROLLED

PURSUIT DG applied preplant incorporated or preemergence will control:

- | | |
|------------------------|---------------------------|
| Buckwheat, wild | Nightshade, Eastern black |
| Kochia | Nightshade, hairy |
| Lambsquarters, common* | Pigweed, redroot |
| Mustard, wild | Shepherdspurse |
| Nightshade, black | Thistle, Russian |

*preplant incorporated applications only

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:

Chickpeas	Peanuts
Lentils	Soybeans
Peas	IMI-Corn™
Lima beans	
2. Four months after PURSUIT DG application:
 - Barley
 - Dry edible beans
 - Rye
 - Snap beans (3 months)
 - Wheat
3. Nine and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed**
 - Tobacco

¹Sencor is a trademark of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen
²Lexone is a trademark of E. I. duPont de Nemours and Company, Inc.
™Trademark of American Cyanamid Company
PE-17228 Rev. 2 1/11/96 SD

4. Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet Corn
Popcorn	
5. Twenty-six months after PURSUIT DG application:
 - Potatoes

6. Forty months after PURSUIT DG application:
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, peas, lentils, chickpeas, or lima beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

** Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.





Supplemental Labeling

55/128

EPA Reg. No. 241-350

FOR USE IN GREAT NORTHERN AND RED KIDNEY TYPE DRY BEANS

FOR DISTRIBUTION AND USE ONLY IN COLORADO, KANSAS, NEBRASKA, WYOMING, TEXAS AND WISCONSIN

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

GENERAL INFORMATION

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG, in any manner not specifically described in this label.

ACCEPTED
with COMMENTS
In EPA Letter Dated
MAR 21 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

241-350

*Registered Trademark of American Cyanamid Company

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after June 30. Allow at least 60 days between application and harvest.

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. **DO NOT** apply PURSUIT DG on sand or loamy sand soils.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Basagran¹ may be tank-mixed with PURSUIT DG to control weeds not listed on the PURSUIT DG label. Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use only nonionic surfactants as a spray additive.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preemergence or early postemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Soybeans
 - Peanuts
 - Edible bean types listed on this label (Refer to heading on page 1).
 - IMI-Corn™
- Four months after PURSUIT DG application:
 - Barley
 - Edible bean types not listed on this label (Refer to heading on page 1).
 - Peas
 - Rye
 - Wheat
- Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
- Nine and one-half months after PURSUIT DG application:
 - Tobacco

¹Dash and Basagran are trademarks of BASF Corporation

™Trademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet Corn |
| Popcorn | |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.





Supplemental Labeling

EPA Reg. No. 241-350

FOR USE IN GREAT NORTHERN AND RED KIDNEY TYPE DRY BEANS

FOR DISTRIBUTION AND USE ONLY IN COLORADO, KANSAS, NEBRASKA, WYOMING, TEXAS AND WISCONSIN

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

GENERAL INFORMATION

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

NOT REVIEWED

In Accordance with PR Notice 82-2
Based On Draft Labeling Dated 3-21-96

*Registered Trademark of American Cyanamid Company

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after June 30. Allow at least 60 days between application and harvest.

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. **DO NOT** apply PURSUIT DG on sand or loamy sand soils.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Basagran¹ may be tank-mixed with PURSUIT DG to control weeds not listed on the PURSUIT DG label. Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use only nonionic surfactants as a spray additive.



DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preemergence or early postemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Soybeans
 - Peanuts
 - Edible bean types listed on this label (Refer to heading on page 1).
 - IMI-CornTM
- Four months after PURSUIT DG application:
 - Barley
 - Edible bean types not listed on this label (Refer to heading on page 1).
 - Peas
 - Rye
 - Wheat
- Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
- Nine and one-half months after PURSUIT DG application:
 - Tobacco

¹Dash and Basagran are trademarks of BASF Corporation
TMTrademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet Corn |
| Popcorn | |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:


All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.





Supplemental Labeling

62/128

EPA Reg. No. 241-350

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, AND SMALL WHITE TYPE DRY BEANS ONLY

**FOR DISTRIBUTION AND USE IN MICHIGAN ONLY
(EXCLUDING UPPER PENINSULA)**

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management, and tillage practices that eliminate competition and hardpan.

ACCEPTED
with comments
in EPA Letter Dated
MAR 21 1996
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
241-350

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray, drift or other indirect contact with PURSUIT DG.

*Registered Trademark of American Cyanamid Company

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply at any timing to Domino variety black turtle beans.

DO NOT apply PURSUIT DG to sand or loamy sand soils.

DO NOT apply PURSUIT DG after July 10.

Allow at least 60 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

Soil Type	Use Rate (oz/acre)
Sands, loamy sands	DO NOT USE
Sandy loams and all other soils with ≤ 2% O.M.	0.72
All other soils with > 2% O.M.	1.08

Preemergence Applications. Apply PURSUIT DG at the rates shown in the preceding table immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

Postemergence Applications. Apply PURSUIT DG at 0.72 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Basagran¹ may be tank-mixed with PURSUIT DG to control weeds not listed on the PURSUIT DG label. Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use only nonionic surfactants as a spray additive.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY. (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
0.72	10 acres
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied to low organic matter soils at the broadcast rate of 0.72 ounces per acre preplant incorporated, preemergence, or postemergence will control light to moderate infestations of:

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - Peanuts
 - IMI-Corn™
 - Edible bean and pea types listed on this label (Refer to heading on page 1)

¹Dash and Basagran are trademarks of BASF Corporation
 ™Trademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

2. Four months after PURSUIT DG application:

- Barley
- Edible bean and pea types not listed on this label (Refer to heading on page 1)
- Rye
- Wheat

3. Eight and one-half months after PURSUIT DG application:

- Field corn
- Field corn grown for seed*

4. Nine and one-half months after PURSUIT DG application:

- Tobacco

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet corn |
| Popcorn | |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.





Supplemental Labeling

65/128

EPA Reg. No. 241-350

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, AND SMALL WHITE TYPE DRY BEANS ONLY

**FOR DISTRIBUTION AND USE IN MICHIGAN ONLY
(EXCLUDING UPPER PENINSULA)**

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

NOT REVIEWED

In Accordance with PR Notice 82-2
Based On Draft Labeling Dated 3-21-96

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply at any timing to Domino variety black turtle beans.

DO NOT apply PURSUIT DG to sand or loamy sand soils.

DO NOT apply PURSUIT DG after July 10.

Allow at least 60 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

Soil Type	Use Rate (oz/acre)
Sands, loamy sands	DO NOT USE
Sandy loams and all other soils with ≤ 2% O.M.	0.72
All other soils with > 2% O.M.	1.08

Preemergence Applications. Apply PURSUIT DG at the rates shown in the preceding table immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.



Postemergence Applications. Apply PURSUIT DG at 0.72 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Basagran¹ may be tank-mixed with PURSUIT DG to control weeds not listed on the PURSUIT DG label. Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use only nonionic surfactants as a spray additive.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
0.72	10 acres
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied to low organic matter soils at the broadcast rate of 0.72 ounces per acre preplant incorporated, preemergence, or postemergence will control light to moderate infestations of:

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - Peanuts
 - IMI-Corn™
 - Edible bean and pea types listed on this label (Refer to heading on page 1)

¹Dash and Basagran are trademarks of BASF Corporation
™Trademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

- 2. Four months after PURSUIT DG application:
 - Barley
 - Edible bean and pea types not listed on this label (Refer to heading on page 1)
 - Rye
 - Wheat

- 3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*

- 4. Nine and one-half months after PURSUIT DG application:
 - Tobacco

- 5. Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	

- 6. Twenty-six months after PURSUIT DG application:
 - Potatoes

- 7. Forty months after PURSUIT DG application:


All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

68/128





Supplemental Labeling

69/128

EPA Reg. No. 241-350

FOR POSTEMERGENCE USE IN PINTO TYPE DRY BEANS

FOR DISTRIBUTION AND USE ONLY IN COLORADO, KANSAS, WYOMING, NEBRASKA, AND TEXAS

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

ACCEPTED
~~with COMMENTS~~
In EPA Letter Dated
MAR 21 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 241-350

*Registered Trademark of American Cyanamid Company

GENERAL INFORMATION

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

Some varieties of pinto beans may be inherently less tolerant to PURSUIT DG applications and crop injury could result. Contact the seed company for information and recommendations about postemergence applications of PURSUIT DG.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

70/128

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after July 15. Allow at least 60 days between application and harvest.

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash.¹

Basagran¹ may be tank-mixed with PURSUIT DG to control weeds not listed on the PURSUIT DG label. Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use only nonionic surfactants as a spray additive.

¹Dash and Basagran are trademarks of BASF Corporation

TMTrademark of American Cyanamid Company

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre early postemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Soybeans
 - Peanuts
 - Great Northern, red kidney, and pinto type dry beans
 - IMI-CornTM
- Four months after PURSUIT DG application:
 - Barley
 - Edible beans types other than Great Northern, red kidney, and pinto type dry beans
 - Peas
 - Rye
 - Wheat
- Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
- Nine and one-half months after PURSUIT DG application:
 - Tobacco

ROTATIONAL CROP GUIDELINE (continued)

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet corn |
| Popcorn | |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.


If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss Great Northern or red kidney beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.





Supplemental Labeling

EPA Reg. No. 241-350

FOR POSTEMERGENCE USE IN PINTO TYPE DRY BEANS

FOR DISTRIBUTION AND USE ONLY IN COLORADO, KANSAS, WYOMING, NEBRASKA, AND TEXAS

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

NOT REVIEWED

In Accordance with PR Notice 82-2
Based On Draft Labeling Dated 3-21-96

GENERAL INFORMATION

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

Some varieties of pinto beans may be inherently less tolerant to PURSUIT DG applications and crop injury could result. Contact the seed company for information and recommendations about postemergence applications of PURSUIT DG.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpan.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

*Registered Trademark of American Cyanamid Company

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after July 15. Allow at least 60 days between application and harvest.

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash.¹

Basagran[™] may be tank-mixed with PURSUIT DG to control weeds not listed on the PURSUIT DG label. Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use only nonionic surfactants as a spray additive.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre early postemergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - Peanuts
 - Great Northern, red kidney, and pinto type dry beans
 - IMI-Corn[™]
2. Four months after PURSUIT DG application:
 - Barley
 - Edible beans types other than Great Northern, red kidney, and pinto type dry beans
 - Peas
 - Rye
 - Wheat
3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
4. Nine and one-half months after PURSUIT DG application:
 - Tobacco

¹Dash and Basagran are trademarks of BASF Corporation
[™]Trademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet corn |
| Popcorn | |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.


If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss Great Northern or red kidney beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.







Supplemental Labeling

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 21 1996

76/128

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

EPA Reg. No. 241-350

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, PINTO, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, DRY EDIBLE PEAS, ENGLISH PEAS, LENTILS, AND CHICKPEAS

241-240

FOR DISTRIBUTION AND USE IN MINNESOTA (south of highway #210) AND SOUTH DAKOTA

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

Some crop varieties may be inherently less tolerant to PURSUIT DG applications and crop injury could result. Contact the seed company for information and recommendations about the use of PURSUIT DG postemergence.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

GENERAL INFORMATION

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

*Registered Trademark of American Cyanamid Company

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT apply to Domino variety black turtle beans.

DO NOT make more than one application of PURSUIT DG per year.

Allow at least 30 days between application and harvest of lima beans and English peas. Allow at least 60 days between application and harvest of dry beans, dry edible peas, lentils and chickpeas.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION INSTRUCTIONS

DO NOT apply PURSUIT DG to sand or loamy sand soils.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre within 1 week before planting. **DO NOT** incorporate deeper than 3 inches. Refer to the PURSUIT DG leaflet label for incorporation instructions.

DO NOT apply preplant incorporated to pinto beans.

Preemergence Applications: Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre after planting, but prior to crop emergence. If weeds are present at the time of application, a burndown herbicide such as Roundup² or Gramoxone Extra³ should be tank-mixed with PURSUIT DG to control emerged weeds.

DO NOT apply preemergence to pinto beans.

Postemergence Applications. FOR USE IN ENGLISH PEAS, NAVY, PINTO, GREAT NORTHERN AND RED KIDNEY TYPE DRY BEANS ONLY. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. **DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.**

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence or postemergence will control:

- Mustard, wild
- Nightshade, hairy
- Nightshade, black
- Pigweed, redroot
- Nightshade, Eastern black

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Chickpeas
 - Lentils
 - Dry edible peas
 - Peanuts
 - Edible beans (dry and lima listed on this label)
 - Soybeans
 - IMI-Corn™
2. Four months after PURSUIT DG application:
 - Barley
 - Edible beans (dry, snap) and peas not listed on this label
 - Rye
 - Wheat
3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*

¹Dash is a trademark of BASF Corporation
²Roundup is a trademark of Monsanto Agricultural Products Company
³Gramoxone Extra is a trademark of Zeneca, Inc.
[™]Trademark American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

4. Nine and one-half months after PURSUIT DG application:
Tobacco

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet corn |
| Popcorn | |

6. Twenty-six months after PURSUIT DG application:
Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, edible beans, dry edible peas, lentils or chickpeas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



79/128



Supplemental Labeling

NOT REVIEWED

In Accordance with PR Notice 82-2
Based On Draft Labeling Dated 3-21-96 EPA Reg. No. 241-350

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, PINTO, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, DRY EDIBLE PEAS, ENGLISH PEAS, LENTILS, AND CHICKPEAS

FOR DISTRIBUTION AND USE IN MINNESOTA (south of highway #210) AND SOUTH DAKOTA

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

GENERAL INFORMATION

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

Some crop varieties may be inherently less tolerant to PURSUIT DG applications and crop injury could result. Contact the seed company for information and recommendations about the use of PURSUIT DG postemergence.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

*Registered Trademark of American Cyanamid Company

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT apply to Domino variety black turtle beans.

DO NOT make more than one application of PURSUIT DG per year.

Allow at least 30 days between application and harvest of lima beans and English peas. Allow at least 60 days between application and harvest of dry beans, dry edible peas, lentils and chickpeas.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION INSTRUCTIONS

DO NOT apply PURSUIT DG to sand or loamy sand soils.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre within 1 week before planting. **DO NOT** incorporate deeper than 3 inches. Refer to the PURSUIT DG leaflet label for incorporation instructions.

DO NOT apply preplant incorporated to pinto beans.

Preemergence Applications: Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre after planting, but prior to crop emergence. If weeds are present at the time of application, a burndown herbicide such as Roundup² or Gramoxone Extra³ should be tank-mixed with PURSUIT DG to control emerged weeds.

DO NOT apply preemergence to pinto beans.

Postemergence Applications. FOR USE IN ENGLISH PEAS, NAVY, PINTO, GREAT NORTHERN AND RED KIDNEY TYPE DRY BEANS ONLY. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. **DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.**

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence or postemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Chickpeas
 - Lentils
 - Dry edible peas
 - Peanuts
 - Edible beans (dry and lima listed on this label)
 - Soybeans
 - IMI-Corn™
- Four months after PURSUIT DG application:
 - Barley
 - Edible beans (dry, snap) and peas not listed on this label
 - Rye
 - Wheat
- Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*

¹Dash is a trademark of BASF Corporation

²Roundup is a trademark of Monsanto Agricultural Products Company

³Gramoxone Extra is a trademark of Zeneca, Inc.

™Trademark American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

4. Nine and one-half months after PURSUIT DG application:
Tobacco

5. Eighteen months after PURSUIT DG application:
Alfalfa Safflower
Cotton Sorghum
Lettuce Sunflower
Oats Sweet corn
Popcorn

6. Twenty-six months after PURSUIT DG application:
Potatoes

7. Forty months after PURSUIT DG application:
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, edible beans, dry edible peas, lentils or chickpeas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.





Supplemental Labeling

83/128

EPA Reg. No. 241-350

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, AND PINTO TYPE DRY BEANS, AND DRY EDIBLE PEAS ONLY

FOR DISTRIBUTION AND USE IN NORTH DAKOTA AND NORTHERN MINNESOTA (see USE AREA below)

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 21 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

*Registered Trademark of American Cyanamid Company

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall) PURSUIT DG may cause injury to subsequent planted crops. Sugar beets are particularly sensitive to PURSUIT DG residues in soil. PURSUIT DG is not recommended in crop rotations that include sugar beets. See the **ROTATIONAL CROP** section of this supplemental label for rotations to other sensitive crops.

ROTATIONAL CROP GUIDELINE (continued)

5. Twenty-six months after PURSUIT DG application:

- Potatoes
- Flax

6. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.


If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996





Supplemental Labeling

EPA Reg. No. 241-350

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, AND PINTO TYPE DRY BEANS, AND DRY EDIBLE PEAS ONLY

NOT REVIEWED FOR DISTRIBUTION AND USE IN NORTH DAKOTA AND NORTHERN MINNESOTA (see USE AREA below)
En Accordance with PR Notice 82-2
Based On Draft Labeling Dated 3.21.96

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall) PURSUIT DG may cause injury to subsequent planted crops. Sugar beets are particularly sensitive to PURSUIT DG residues in soil. PURSUIT DG is not recommended in crop rotations that include sugar beets. See the ROTATIONAL CROP section of this supplemental label for rotations to other sensitive crops.

*Registered Trademark of American Cyanamid Company

USE AREA

PURSUIT DG may be used in North Dakota and in Minnesota north of highway #210.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply at any timing to Domino variety black turtle beans.

Pinto varieties UI-111 and Olathe are more sensitive to PURSUIT DG than other varieties.

DO NOT apply PURSUIT DG after July 10.

DO NOT feed treated forage, hay, or straw to livestock.

Allow at least 60 days between application and harvest.

Postemergence Applications. Apply PURSUIT DG at 0.72 ounces per acre (one ECO-PAK soluble bag treats 10 acres). A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 0.72 ounces per acre will control:

wild mustard
black nightshade*
eastern black nightshade*

*suppression

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - IMI-Corn™
 - Edible bean and pea types listed on this label (Refer to heading on page 1)
2. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
3. Nine and one-half months after PURSUIT DG application:
 - Wheat
 - Edible bean and pea types not listed on this label (Refer to heading on page 1)
4. Eighteen months after PURSUIT DG application:

Barley	Rye
Cotton	Safflower
Lettuce	Sorghum
Oats	Sunflower
Popcorn	Sweet Corn

¹Dash is a trademark of BASF Corporation

™Trademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

5. Twenty-six months after PURSUIT DG application:

- Potatoes
- Flax

6. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

89/128





Supplemental Labeling

90/128

EPA Reg. No. 241-350

FOR USE IN SOYBEANS

PURSUIT DG PLUS SCEPTER® OR SCEPTER® DG HERBICIDES FOR SUPPRESSION OF VOLUNTEER CORN AND ENHANCED CONTROL OF SUNFLOWERS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR WORKER PROTECTION STANDARD REQUIREMENTS, USE AREA, MIXING AND APPLICATION INFORMATION, AND WEEDS CONTROLLED.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

GENERAL INFORMATION

PURSUIT DG may be tank mixed with SCEPTER or SCEPTER DG for suppression of volunteer corn or for enhanced control of common sunflower in soybeans. Apply postemergence to soybeans only. In this label, SCEPTER refers to both formulations SCEPTER and SCEPTER DG. Be sure to refer to the correct label for additional information.

USE AREA

The application of PURSUIT DG plus SCEPTER or SCEPTER DG may be applied to states or portions of states described as Region 2 or Region 3 on the SCEPTER or SCEPTER DG labels, and the following counties in South Dakota: Yankton, Bon Homme, Hutchinson, McCook, Hanson, Davison, Miner, Lake and Kingsbury. Refer to the respective labels for the recommended use area. Do not use this tank mixture in North Dakota or in Minnesota north of state highway #210.

APPLICATION RATES

Apply the products at the following rate:

PURSUIT DG	1.44 oz/A*
AND	
SCEPTER	2 oz/A (liquid)
OR	
SCEPTER DG	0.53 oz/A**

ACCEPTED
with COMMENTS
In EPA Letter Dated
MAR 21 1996
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under Reg. No.
241-350

*At the rate of 1.44 ounces per acre, one 7.2 ounce soluble bag of PURSUIT DG will treat 5 acres.

**At the rate of 0.53 ounces per acre, one 14 ounce soluble bag of SCEPTER DG will treat 26.4 acres.

WEEDS SUPPRESSED/CONTROLLED

The tank mixture of PURSUIT DG plus SCEPTER or SCEPTER DG will suppress volunteer corn. Apply to volunteer corn up to 10 inches in height. The tank-mixture of PURSUIT DG and SCEPTER or SCEPTER DG will enhance the control of common sunflowers. Apply to sunflowers up to 3 inches in size. Refer to the SCEPTER or SCEPTER DG label for additional weeds controlled.

A postemergence application of PURSUIT DG plus SCEPTER products will not suppress volunteer IMI-Corn™ hybrids (field corn hybrids which possess tolerance or resistance to imidazolinone herbicides i.e., PURSUIT DG and SCEPTER).

ROTATIONAL CROP GUIDELINES

The following rotational crops may be planted after applying PURSUIT DG when tank-mixed with SCEPTER or SCEPTER DG at the recommended rates.

1. Anytime
Soybeans
2. Four months after PURSUIT DG + SCEPTER application:
Wheat
3. Nine and one half months after a PURSUIT DG + SCEPTER application:
Field Corn*
4. Eleven months after a PURSUIT DG + SCEPTER application.
Edible beans and peas
5. Eighteen months after PURSUIT DG + SCEPTER application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	
6. Twenty six months after PURSUIT DG + SCEPTER application.
Potatoes
7. Forty months after PURSUIT DG + SCEPTER application
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

NOTE: Following forty months after a PURSUIT DG + SCEPTER application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.



Supplemental Labeling

NOT REVIEWED
in Accordance with PR Notice 82-2
based On Draft Labeling Dated 3.21.76

EPA Reg. No. 241-350

FOR USE IN SOYBEANS

PURSUIT DG PLUS SCEPTER® OR SCEPTER® DG HERBICIDES FOR SUPPRESSION OF VOLUNTEER CORN AND ENHANCED CONTROL OF SUNFLOWERS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR WORKER PROTECTION STANDARD REQUIREMENTS, USE AREA, MIXING AND APPLICATION INFORMATION, AND WEEDS CONTROLLED.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

GENERAL INFORMATION

PURSUIT DG may be tank mixed with SCEPTER or SCEPTER DG for suppression of volunteer corn or for enhanced control of common sunflower in soybeans. Apply postemergence to soybeans only. In this label, SCEPTER refers to both formulations SCEPTER and SCEPTER DG. Be sure to refer to the correct label for additional information.

USE AREA

The application of PURSUIT DG plus SCEPTER or SCEPTER DG may be applied to states or portions of states described as Region 2 or Region 3 on the SCEPTER or SCEPTER DG labels, and the following counties in South Dakota: Yankton, Bon Homme, Hutchinson, McCook, Hanson, Davison, Miner, Lake and Kingsbury. Refer to the respective labels for the recommended use area. Do not use this tank mixture in North Dakota or in Minnesota north of state highway #210.

APPLICATION RATES

Apply the products at the following rate:	
PURSUIT DG	1.44 oz/A*
AND	
SCEPTER	2 oz/A (liquid)
OR	
SCEPTER DG	0.53 oz/A**

*At the rate of 1.44 ounces per acre, one 7.2 ounce soluble bag of PURSUIT DG will treat 5 acres.

**At the rate of 0.53 ounces per acre, one 14 ounce soluble bag of SCEPTER DG will treat 26.4 acres.

94/128

WEEDS SUPPRESSED/CONTROLLED

The tank mixture of PURSUIT DG plus SCEPTER or SCEPTER DG will suppress volunteer corn. Apply to volunteer corn up to 10 inches in height. The tank-mixture of PURSUIT DG and SCEPTER or SCEPTER DG will enhance the control of common sunflowers. Apply to sunflowers up to 3 inches in size. Refer to the SCEPTER or SCEPTER DG label for additional weeds controlled.

A postemergence application of PURSUIT DG plus SCEPTER products will not suppress volunteer IMI-Corn™ hybrids (field corn hybrids which possess tolerance or resistance to imidazolinone herbicides i.e., PURSUIT DG and SCEPTER).

ROTATIONAL CROP GUIDELINES

The following rotational crops may be planted after applying PURSUIT DG when tank-mixed with SCEPTER or SCEPTER DG at the recommended rates.

1. Anytime
Soybeans
2. Four months after PURSUIT DG + SCEPTER application:
Wheat
3. Nine and one half months after a PURSUIT DG + SCEPTER application:
Field Corn*
4. Eleven months after a PURSUIT DG + SCEPTER application.
Edible beans and peas
5. Eighteen months after PURSUIT DG + SCEPTER application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	
6. Twenty six months after PURSUIT DG + SCEPTER application.
Potatoes
7. Forty months after PURSUIT DG + SCEPTER application
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

NOTE: Following forty months after a PURSUIT DG + SCEPTER application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

95/
128

* Field corn may be planted 9 1/2 months following a postemergence application of PURSUIT DG when applied in tank-mixture with SCEPTER or SCEPTER DG at the recommended rates unless extreme drought conditions develop (less than 12 inches of rainfall or irrigation is received from two weeks prior to the date of PURSUIT DG application through November 15 of the same year). If the minimum rainfall requirement is not met, only field corn hybrids (IMI-Corn™) which possess tolerance or resistance to PURSUIT DG may be planted the spring of the year following a PURSUIT DG + SCEPTER application.

Use of PURSUIT DG and SCEPTER herbicides in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of these products and, therefore, rotational crop injury is always possible.

Except as stated above, DO NOT use this tank mixture in the same year as an application of SCEPTER, SCEPTER DG, SQUADRON®, SQUADRON® DG TRI-SCEPT®, SCEPTER® O.T.®, STEEL™ CP, DETAIL®, PURSUIT®, PURSUIT® DG, PURSUIT® PLUS EC, CONTOUR® or RESOLVE®. Also, DO NOT use the above tank mixture in the same year as an application of a product containing chlorimuron ethyl (e.g. Classic¹, Canopy¹, Concert¹, Gemini¹, Lorox Plus¹, Preview¹, Reliance¹, Synchrony¹ etc.).

96/128





Supplemental Labeling



EPA Reg. No. 241-350

97/128

FOR USE IN FIELD CORN (APPLY ONLY ON IMI-CORN™ INBREDS)

USE DIRECTIONS FOR THE USE OF PURSUIT DG FOR WEED CONTROL IN IMI-CORN INBREDS

OBSERVE ALL PRECAUTIONARY STATEMENTS ON THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

Apply PURSUIT DG herbicide only on selected field corn inbreds (IMI-Corn) warranted by the seed company to possess resistance/tolerance to direct application of PURSUIT DG. DO NOT apply PURSUIT DG to corn inbreds which lack genetic resistance/tolerance to PURSUIT DG herbicide. Contact your chemical dealer, seed supplier or American Cyanamid to obtain information regarding IMI-Corn inbreds.

PURSUIT DG can be applied as an early preplant, preplant incorporated, preemergence or postemergence treatment on IMI-Corn inbreds.

Crops growing under stressful environmental conditions can exhibit various injury symptoms which may be more pronounced if herbicides are used. Corn plants treated with PURSUIT DG may exhibit yellowing on new growth. Such effects occur infrequently and are temporary. Normal growth and appearance should resume within 1 to 2 weeks.

Naturally occurring biotypes* of some of the weeds listed on this label may not be effectively controlled by this and/or other products with either the ALS/AHAS enzyme inhibiting mode of action. Other herbicides with the ALS/AHAS enzyme inhibiting mode of action include

the sulfonylureas (e.g., Accent™, Basis™, Classic™, Concert™, Exceed™, Permit™, Pinnacle™, etc.), the sulfonamides (e.g., Broadstrike™, etc.) and the pyrimidyl benzoates (e.g. Staple™, etc.). If naturally occurring ALS/AHAS resistant biotypes are present in a field, Pursuit DG and/or any other ALS/AHAS enzyme inhibiting mode of action herbicide should be tank-mixed or applied sequentially with an appropriate registered herbicide having a different mode of action to ensure control.

* A weed biotype is a naturally occurring plant within a given species that has a slightly different, but distinct, genetic makeup from other plants.

Replanting: If replanting is necessary in a field previously treated with PURSUIT DG, the field may be replanted to IMI-Corn, soybeans, peanuts, lima beans or southern peas. Rework the soil no deeper than the treated zone. DO NOT apply a second treatment of PURSUIT DG herbicide.

MIXING INSTRUCTIONS

Fill the spray tank one-fourth to one-half full with clean water. Add the required number of PURSUIT DG soluble packets to the spray tank while agitating. Add adjuvants and fill the remainder of the tank with water.

TANK MIX COMBINATIONS WITH OTHER HERBICIDES

If other herbicides are tank-mixed with PURSUIT DG, while agitating, add components in the following order:

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 21 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241-350

98/128

1. Fill spray tank 1/4 to 1/2 full with clean water.
2. Add soluble packet products and thoroughly mix.
3. Add WP (wetttable powder), DG (dispersible granule), DF (dry flowable) or liquid flowable formulations not in soluble packets.
4. Add PURSUIT DG and thoroughly mix.
5. Add other aqueous solution products.
6. Add EC (emulsifiable concentrate) products.
7. Add surfactant or crop oil to the spray tank.
8. Add liquid fertilizer.
9. While agitating, fill the remainder of the tank with water.

If PURSUIT DG is applied in liquid fertilizer as a carrier, PURSUIT DG must first be dissolved in a water slurry. Premix PURSUIT DG in a minimum of one gallon of water per ECO-PAK prior to adding to the spray tank.

DO NOT add PURSUIT DG ECO-PAK packets directly into liquid fertilizer.

USE RATE
(1.44 OUNCES PER ACRE)
 (1 ECO-PAK soluble bag treats 5 acres)

Apply PURSUIT DG at a broadcast rate of 1 soluble bag per 5 acres (7.2 ounces) for all methods of application: early preplant, preplant incorporated, preemergence, and postemergence (including minimum and no-till). At this broadcast rate, one soluble bag PURSUIT DG will treat 5 acres of IMI-Corn.

NOTE: Only one application of PURSUIT DG may be made during the season.

PURSUIT DG HERBICIDE USE AREA
 (Not for use in California)

DO NOT apply PURSUIT DG herbicide to IMI-Corn in North Dakota or in Minnesota north of state highway 210.

APPLICATION INSTRUCTIONS
POSTEMERGENCE

Apply PURSUIT DG herbicide as a postemergence treatment to IMI-Corn when crop and weeds are actively growing. Apply PURSUIT DG before weeds exceed a height of 3 inches, unless otherwise indicated (see PURSUIT DG leaflet label). More restrictive crop growth stage limitations of tank-mix partners must be followed.

ADDITIVES:

POSTEMERGENCE APPLICATIONS OF PURSUIT DG REQUIRE THE ADDITION OF AN ADJUVANT AND A LIQUID FERTILIZER¹ SOLUTION.

I. ADJUVANTS

SURFACTANTS: Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 quart per 100 gallons of spray solution.

OR

CROP OIL CONCENTRATE: Instead of a surfactant, a petroleum or vegetable seed oil based crop oil concentrate (such as SUN-IT IITM) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other methylated seed oils at the rate of 1.5 - 2 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre.

AND

II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs per acre.

DO NOT apply PURSUIT DG herbicide with liquid fertilizer as a carrier for postemergence applications.

SOIL APPLICATIONS

PURSUIT DG can be applied as an early preplant, preplant incorporated, or preemergence treatment in IMI-Corn.

PURSUIT DG can be applied to the soil in liquid fertilizers, alone or in combination with PROWL[®] 3.3EC, Lasso EC[™], Lasso MT[™] or Dual[™]. Follow all PURSUIT DG label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. Always test the compatibility of PURSUIT DG herbicide with the liquid fertilizer before mixing in the spray tank.

¹ PURSUIT DG applications may be made with a non-ionic surfactant only (liquid fertilizer not required) in the state of Arkansas, Bootheel of Missouri, Tennessee, South Carolina, Louisiana, Mississippi, Alabama, and Georgia.

**SEQUENTIAL HERBICIDE
COMBINATIONS WITH PURSUIT DG**

**PURSUIT DG FOLLOWING EARLY PREPLANT
APPLICATION OF:**

2,4-D	Harness™
Bicep™, Bicep Lite™	Harness Extra
Bullet™	Lasso EC
Bronco™	Lasso MT
Dual	Roundup™
Extrazine II™	Surpass™
Frontier™	Surpass 100
Gramoxone Extra™	

**PURSUIT DG FOLLOWING PREPLANT
INCORPORATED APPLICATION OF:**

Atrazine ²	Harness
Bicep, Bicep Lite	Harness Extra
Bullet	Lariat™
Dual	Lasso EC, Lasso MT
Eradicane Extra™	Sutan+™
Extrazine II	Surpass
Frontier	Surpass 100

**PURSUIT DG FOLLOWING PREEMERGENCE
APPLICATION OF:**

Atrazine	Frontier
Banvel™, ²	Harness
Bicep, Bicep Lite	Harness Extra
Bladex™ 4L	Lariat
Bladex 90DF	Lasso EC, Lasso MT
Bronco	Marksman™, ²
Bullet	PROWL 3.3 EC
Clarity™, ²	Surpass
Dual	Surpass 100
Extrazine	

**TANK-MIX HERBICIDE
COMBINATIONS WITH PURSUIT DG**

**EARLY PREPLANT APPLICATION OF PURSUIT
DG TANK-MIXED WITH:**

2,4-D	Harness
Bronco	Lasso EC
Dual	Lasso MT
Frontier	Roundup
Gramoxone Extra	Surpass

**PREPLANT INCORPORATED APPLICATION OF
PURSUIT DG TANK-MIXED WITH:**

Atrazine	Frontier
Bicep, Bicep Lite	Harness
Bullet	Lariat
Dual	Lasso EC, Lasso MT
Eradicane Extra	Surpass
Extrazine	Sutan+

**PREEMERGENCE APPLICATION OF PURSUIT
DG TANK-MIXED WITH:**

Atrazine	Frontier
Banvel	Harness
Bicep, Bicep Lite	Harness Extra
Bladex 4L	Lariat
Bladex 90DF	Lasso EC, Lasso MT
Bronco	Marksman
Bullet	PROWL 3.3 EC
Clarity	Surpass
Dual	Surpass 100
Extrazine	

**POSTEMERGENCE APPLICATIONS OF PURSUIT
DG TANK-MIXED WITH:**

Atrazine ³	Frontier
Banvel ⁴	Lasso EC, Lasso MT
Basagran™, ²	Marksman
Buctril™, ^{2,3}	PROWL 3.3 EC
Clarity ⁴	Sencor DF™
Dual	Sencor 4L™

²In some cases the grass activity of PURSUIT DG will be reduced when used in combination with atrazine, Buctril, Banvel, Basagran, Clarity, or Marksman.

³Some corn leaf burn may result with Buctril or atrazine postemergence combinations with PURSUIT DG.

⁴Applications of Banvel or Clarity herbicide to corn during periods of rapid growth may result in temporary leaning.

DO NOT use crop oil concentrates as adjuvants in PURSUIT DG herbicide combinations with Banvel, Buctril, Clarity, Marksman, Sencor DF or Sencor 4L.

PURSUIT DG is active against many broadleaf and grass species. However, for long term weed management, alternate mode of action herbicides are recommended with PURSUIT DG. The application of a soil applied grass herbicide underlay will control grass weeds not on the PURSUIT DG label and enhance the control of certain broadleaf weeds such as common lambsquarters.

When PURSUIT DG herbicide is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions.

AERIAL APPLICATIONS

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre. When applied POSTEMERGENCE, the addition of a non-ionic surfactant AND fertilizer solution are required for optimum weed control. Apply a non-ionic surfactant at the rate of 1 quart per 100 gallons of spray mixture OR a crop oil concentrate at the rate of 1.5 - 2 pints per acre AND a liquid fertilizer at the rate of 1 quart per acre. (See instructions under APPLICATION INSTRUCTIONS - POSTEMERGENCE).

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS.

- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph. Coarse sprays (larger droplets) are less likely to drift.

Applicator is responsible for any loss or damage which results from spraying PURSUIT DG in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

When applied as directed at the broadcast rate of 1.44 ounces per acre, PURSUIT DG will control or reduce competition from the weeds listed below.

NOTE: C = Control R = Reduced Competition

BROADLEAF WEEDS

Weed Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Alligator weed		4	1-3
Anoda, spurred	C	2	1-2
Artichoke, Jerusalem		8	6-10
Buffalobur	C*	R	1-3
Bristly starbur		2	1-2
Carpetweed	C		
Cocklebur, common	R	8	1-8
Galinsoga	C		
Jimsonweed	C*	4	1-3
Kochia	C	4	1-3
Lambsquarters, common	C*	R	1-2
Mallow, Venice	R		
Marshelder	C	4	1-3
Morningglory, entireleaf	R	2	1-2
ivyleaf	R	2	1-2
pitted	R	2	1-2
smallflower	C	4	1-3
tall	R	2	1-2
Mustard sp.	C	4	1-3

Nightshade, black	C	4	1-3
Eastern black hairy	C	4	1-3
Pigweed, Palmer	C	8	1-8
redroot	C	8	1-8
smooth	C	8	1-8
spiny	C	8	1-8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C*		
Ragweed, common	R	4	1-3
giant	R	4	1-3
Sage, barnyard		R	1-3
Smartweed, ladythumb	C	4	1-3
Pennsylvania	C	4	1-3
Spurge, prostrate	C	4	1-3
spotted	C	4	1-3
Sunflower	C*	4	1-3
Velvetleaf	C*	4	1-3
Thistle, Canada		R	1-3

*When PURSUIT DG is soil applied, these weeds are more consistently controlled by preplant incorporated treatments.

DO NOT count cotyledon leaves when determining weed stage of growth.

GRASS WEEDS

Weed Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Barnyardgrass	R	3	1-3
Crabgrass, large	R	3	1-3
smooth	R	3	1-3
Cupgrass, woolly		3	1-3
Foxtail, giant	C	6	1-6
green	C	3	1-3
giant green	C	3	1-3
Goosegrass	R		
Johnsongrass, seedling	C	6	1-8
rhizome		R	1-8
Millet, wild proso	R	R	1-3
Panicum, fall	R		
Texas	R		

Red rice		3	1-3
Sandbur, field	R	R	≤1
Shattercane	R	6	1-8
Signalgrass,			
broadleaf	R	4	1-8
Sorghum alnum	R	6	1-3

SEDGES

Nutsedge,			
purple	R	R	1-3
yellow	R	R	1-3

*When PURSUIT DG is applied to grasses, preplant incorporated treatments of PURSUIT DG are more consistent.

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence.)

ROTATIONAL CROPS

The following rotational crops may be planted after applying PURSUIT DG at the recommended rate: (Planting earlier than the recommended interval may result in crop injury).

1. Anytime
 - Lima beans
 - Southern peas
 - Soybeans
 - Peanuts
 - IMI-Corn™ (resistant/tolerant to PURSUIT DG)
2. Four months after PURSUIT DG application:
 - Alfalfa
 - Rye
 - Wheat
 - Edible beans and peas (other than lima beans and Southern peas)
3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
4. Nine and one-half months after PURSUIT DG application:
 - Barley
 - Tobacco

5. Eighteen months after PURSUIT DG application:

Cotton	Safflower
Lettuce	Sorghum
Oats	Sunflower
Popcorn	Sweet corn

6. Twenty six months after PURSUIT DG application.

Potatoes

7. Forty months after PURSUIT DG application.

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH). If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

*Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

PRECAUTIONS

In the event of a crop loss due to weather, IMI-Corn inbreds and hybrids, soybeans, lima beans, southern peas or peanuts can be replanted if PURSUIT DG herbicide was applied alone at rates labelled for these crops. If PURSUIT DG herbicide was tank-mixed with other herbicides, such as atrazine, the label restrictions for these herbicides must also be followed.

There should be an interval of at least 45 days between an application of PURSUIT DG and corn harvest (silage, fodder, or grain). **DO NOT** graze or feed treated corn, forage, silage, fodder, or grain for at least 45 days after an application of PURSUIT DG.

All soil insecticides, including labelled banded or in-furrow applications, may be used in combination with Pioneer imidazolinone resistant (IR) corn inbreds.

Imidazolinone tolerant inbreds from other seed companies may occasionally exhibit injury symptoms when soil insecticides are used in combination with PURSUIT DG. USE banded applications of COUNTER® 15G, COUNTER® 20CR® or THIMET® in combination with PURSUIT DG herbicide application; **DO NOT USE COUNTER 15G** soil systemic insecticide products in-furrow with imidazolinone tolerant inbreds. American Cyanamid has not tested all inbreds in which the imidazolinone tolerance trait is claimed and cannot be responsible for factors which are beyond its control, such as growing conditions, environmental conditions, grower practices and the specific genetics of each inbred which may influence tolerance. Growers should contact the seed company for information regarding specific inbred tolerance to herbicide and insecticide applications.

This label must be in the possession of the user at the time of herbicide application.

For additional information regarding the use of PURSUIT DG herbicide, call telephone no. (800) 942-0500.

PE-xxxxx 3/96 SD

®Registered Trademark of American Cyanamid Company

™SUN-IT II is a Trademark of Agsco, Inc.

™Buctril is a Trademark of Rhone Poulenc Ag Company.

™Banvel, Clarity, Frontier, and Marksman are Trademarks of Sandoz Crop Protection Corporation.

™Basagran is a Trademark of BASF Corporation.

™Beacon, Bicep, Bicep Lite, Dual, and Exceed are Trademarks of Ciba Geigy Corporation.

™Bronco, Bullet, Harness, Harness Extra, Lariat, Lasso EC, Lasso MT, Permit and Roundup are Trademarks of Monsanto Agricultural Products Company.

™Sencor is a Trademark of Mobay Corporation.

™Gramoxone Extra, Eradicane Extra, Surpass, Surpass 100, and Sutan+ are Trademarks of Zeneca Agrochemicals.

™Accent, Basis, Bladex 4L, Bladex 90DF, Classic, Concert, Extrazine, Pinnacle, and Staple are Trademarks of E. I. DuPont de Nemours and Company, Inc.

™Imi-Corn is a Trademark of American Cyanamid Company

™Broadstrike is a Trademark of Dow Elanco Corp.



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996



Supplemental Labeling



EPA Reg. No. 241-350

NOT REVIEWED

In Accordance with PR Notice 82-2
Based On Draft Labeling Dated 3-21-96

**FOR USE IN FIELD CORN
(APPLY ONLY ON IMI-CORN™ INBREDS)**

USE DIRECTIONS FOR THE USE OF PURSUIT DG FOR WEED CONTROL IN IMI-CORN INBREDS

OBSERVE ALL PRECAUTIONARY STATEMENTS ON THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

Apply PURSUIT DG herbicide only on selected field corn inbreds (IMI-Corn) warranted by the seed company to possess resistance/tolerance to direct application of PURSUIT DG. **DO NOT** apply PURSUIT DG to corn inbreds which lack genetic resistance/tolerance to PURSUIT DG herbicide. Contact your chemical dealer, seed supplier or American Cyanamid to obtain information regarding IMI-Corn inbreds.

PURSUIT DG can be applied as an early preplant, preplant incorporated, preemergence or postemergence treatment on IMI-Corn inbreds.

Crops growing under stressful environmental conditions can exhibit various injury symptoms which may be more pronounced if herbicides are used. Corn plants treated with PURSUIT DG may exhibit yellowing on new growth. Such effects occur infrequently and are temporary. Normal growth and appearance should resume within 1 to 2 weeks.

Naturally occurring biotypes* of some of the weeds listed on this label may not be effectively controlled by this and/or other products with either the ALS/AHAS enzyme inhibiting mode of action. Other herbicides with the

ALS/AHAS enzyme inhibiting mode of action include the sulfonylureas (e.g., Accent™, Basis™, Classic™, Concert™, Exceed™, Permit™, Pinnacle™, etc.), the sulfonamides (e.g., Broadstrike™, etc.) and the pyrimidyl benzoates (e.g. Staple™, etc.). If naturally occurring ALS/AHAS resistant biotypes are present in a field, Pursuit DG and/or any other ALS/AHAS enzyme inhibiting mode of action herbicide should be tank-mixed or applied sequentially with an appropriate registered herbicide having a different mode of action to ensure control.

* A weed biotype is a naturally occurring plant within a given species that has a slightly different, but distinct, genetic makeup from other plants.

Replanting: If replanting is necessary in a field previously treated with PURSUIT DG, the field may be replanted to IMI-Corn, soybeans, peanuts, lima beans or southern peas. Rework the soil no deeper than the treated zone. **DO NOT** apply a second treatment of PURSUIT DG herbicide.

MIXING INSTRUCTIONS

Fill the spray tank one-fourth to one-half full with clean water. Add the required number of PURSUIT DG soluble packets to the spray tank while agitating. Add adjuvants and fill the remainder of the tank with water.

TANK MIX COMBINATIONS WITH OTHER HERBICIDES

1. Fill spray tank 1/4 to 1/2 full with clean water.
2. Add soluble packet products and thoroughly mix.
3. Add WP (wetttable powder), DG (dispersible granule), DF (dry flowable) or liquid flowable formulations not in soluble packets.
4. Add PURSUIT DG and thoroughly mix.
5. Add other aqueous solution products.
6. Add EC (emulsifiable concentrate) products.
7. Add surfactant or crop oil to the spray tank.
8. Add liquid fertilizer.
9. While agitating, fill the remainder of the tank with water.

If PURSUIT DG is applied in liquid fertilizer as a carrier, PURSUIT DG must first be dissolved in a water slurry. Premix PURSUIT DG in a minimum of one gallon of water per ECO-PAK prior to adding to the spray tank.

DO NOT add PURSUIT DG ECO-PAK packets directly into liquid fertilizer.

USE RATE
(1.44 OUNCES PER ACRE)
(1 ECO-PAK soluble bag treats 5 acres)

Apply PURSUIT DG at a broadcast rate of 1 soluble bag per 5 acres (7.2 ounces) for all methods of application: early preplant, preplant incorporated, preemergence, and postemergence (including minimum and no-till). At this broadcast rate, one soluble bag PURSUIT DG will treat 5 acres of IMI-Corn.

NOTE: Only one application of PURSUIT DG may be made during the season.

PURSUIT DG HERBICIDE USE AREA
(Not for use in California)

DO NOT apply PURSUIT DG herbicide to IMI-Corn in North Dakota or in Minnesota north of state highway 210.

APPLICATION INSTRUCTIONS
POSTEMERGENCE

Apply PURSUIT DG herbicide as a postemergence treatment to IMI-Corn when crop and weeds are actively growing. Apply PURSUIT DG before weeds exceed a height of 3 inches, unless otherwise indicated (see PURSUIT DG label). More restrictive crop growth stage limitations of tank-mix partners must be followed.

ADDITIVES:

POSTEMERGENCE APPLICATIONS OF PURSUIT DG REQUIRE THE ADDITION OF AN ADJUVANT AND A LIQUID FERTILIZER¹ SOLUTION.

I. ADJUVANTS

SURFACTANTS: Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 quart per 100 gallons of spray solution.

OR

CROP OIL CONCENTRATE: Instead of a surfactant, a petroleum or vegetable seed oil based crop oil concentrate (such as SUN-IT II™) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other methylated seed oils at the rate of 1.5 - 2 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre.

AND

II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs. per acre.

DO NOT apply PURSUIT DG herbicide with liquid fertilizer as a carrier for postemergence applications.

SOIL APPLICATIONS

PURSUIT DG can be applied as an early preplant, preplant incorporated, or preemergence treatment in IMI-Corn.

PURSUIT DG can be applied to the soil in liquid fertilizers, alone or in combination with PROWL® 3.3EC, Lasso EC™, Lasso MT™ or Dual™. Follow all PURSUIT DG label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. Always test the compatibility of PURSUIT DG herbicide with the liquid fertilizer before mixing in the spray tank.

¹ PURSUIT DG applications may be made with a non-ionic surfactant only (liquid fertilizer not required) in the state of Arkansas, Bootheel of Missouri, Tennessee, South Carolina, Louisiana, Mississippi, Alabama, and Georgia.

**SEQUENTIAL HERBICIDE
COMBINATIONS WITH PURSUIT DG**

**PURSUIT DG FOLLOWING EARLY PREPLANT
APPLICATION OF:**

2,4-D	Harness™
Bicep™, Bicep Lite™	Harness Extra
Bullet™	Lasso EC
Bronco™	Lasso MT
Dual	Roundup™
Extrazine II™	Surpass™
Frontier™	Surpass 100
Gramoxone Extra™	

**PURSUIT DG FOLLOWING PREPLANT
INCORPORATED APPLICATION OF:**

Atrazine ²	Harness
Bicep, Bicep Lite	Harness Extra
Bullet	Lariat™
Dual	Lasso EC, Lasso MT
Eradicane Extra™	Sutan+™
Extrazine II	Surpass
Frontier	Surpass 100

**PURSUIT DG FOLLOWING PREEMERGENCE
APPLICATION OF:**

Atrazine	Frontier
Banvel™,2	Harness
Bicep, Bicep Lite	Harness Extra
Bladex™ 4L	Lariat
Bladex 90DF	Lasso EC, Lasso MT
Bronco	Marksman™,2
Bullet	PROWL 3.3 EC
Clarity™,2	Surpass
Dual	Surpass 100
Extrazine	

**TANK-MIX HERBICIDE
COMBINATIONS WITH PURSUIT DG**

**EARLY PREPLANT APPLICATION OF PURSUIT
DG TANK-MIXED WITH:**

2,4-D	Harness
Bronco	Lasso EC
Dual	Lasso MT
Frontier	Roundup
Gramoxone Extra	Surpass

**PREPLANT INCORPORATED APPLICATION OF
PURSUIT DG TANK-MIXED WITH:**

Atrazine	Frontier
Bicep, Bicep Lite	Harness
Bullet	Lariat
Dual	Lasso EC, Lasso MT
Eradicane Extra	Surpass
Extrazine	Sutan+

**PREEMERGENCE APPLICATION OF PURSUIT
DG TANK-MIXED WITH:**

Atrazine	Frontier
Banvel	Harness
Bicep, Bicep Lite	Harness Extra
Bladex 4L	Lariat
Bladex 90DF	Lasso EC, Lasso MT
Bronco	Marksman
Bullet	PROWL 3.3 EC
Clarity	Surpass
Dual	Surpass 100
Extrazine	

**POSTEMERGENCE APPLICATIONS OF PURSUIT
DG TANK-MIXED WITH:**

Atrazine ³	Frontier
Banvel ⁴	Lasso EC, Lasso MT
Basagran™,2	Marksman
Buctril™,2,3	PROWL 3.3 EC
Clarity ⁴	Sencor DF™
Dual	Sencor 4L™

²In some cases the grass activity of PURSUIT DG will be reduced when used in combination with atrazine, Buctril, Banvel, Basagran, Clarity, or Marksman.

³Some corn leaf burn may result with Buctril or atrazine postemergence combinations with PURSUIT DG.

⁴Applications of Banvel or Clarity herbicide to corn during periods of rapid growth may result in temporary leaning.

DO NOT use crop oil concentrates as adjuvants in PURSUIT DG herbicide combinations with Banvel, Buctril, Clarity, Marksman, Sencor DF or Sencor 4L.

PURSUIT DG is active against many broadleaf and grass species. However, for long term weed management, alternate mode of action herbicides are recommended with PURSUIT DG. The application of a soil applied grass herbicide underlay will control grass weeds not on the PURSUIT DG label and enhance the control of certain broadleaf weeds such as common lambsquarters.

When PURSUIT DG herbicide is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions.

AERIAL APPLICATIONS

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre. When applied POSTEMERGENCE, the addition of a non-ionic surfactant AND fertilizer solution are required for optimum weed control. Apply a non-ionic surfactant at the rate of 1 quart per 100 gallons of spray mixture OR a crop oil concentrate at the rate of 1.5 - 2 pints per acre AND a liquid fertilizer at the rate of 1 quart per acre. (See instructions under APPLICATION INSTRUCTIONS - POSTEMERGENCE).

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS.

- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph. Coarse sprays (larger droplets) are less likely to drift.

Applicator is responsible for any loss or damage which results from spraying PURSUIT DG in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

When applied as directed at the broadcast rate of 1.44 ounces per acre, PURSUIT DG will control or reduce competition from the weeds listed below.

NOTE: C = Control R = Reduced Competition

BROADLEAF WEEDS

Weed Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Alligator weed		4	1-3
Anoda, spurred	C	2	1-2
Artichoke, Jerusalem		8	6-10
Buffalobur	C*	R	1-3
Bristly starbur		2	1-2
Carpetweed	C		
Cocklebur, common	R	8	1-8
Galinsoga	C		
Jimsonweed	C*	4	1-3
Kochia	C	4	1-3
Lambsquarters, common	C*	R	1-2
Mallow, Venice	R		
Marshelder	C	4	1-3
Morningglory, entireleaf	R	2	1-2
ivyleaf	R	2	1-2
pitted	R	2	1-2
smallflower	C	4	1-3
tall	R	2	1-2
Mustard sp.	C	4	1-3

Nightshade, black	C	4	1-3
Eastern black hairy	C	4	1-3
Pigweed, Palmer	C	8	1-8
redroot	C	8	1-8
smooth	C	8	1-8
spiny	C	8	1-8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C*		
Ragweed, common	R	4	1-3
giant	R	4	1-3
Sage, barnyard		R	1-3
Smartweed, ladythumb	C	4	1-3
Pennsylvania	C	4	1-3
Spurge, prostrate	C	4	1-3
spotted	C	4	1-3
Sunflower	C*	4	1-3
Velvetleaf	C*	4	1-3
Thistle, Canada		R	1-3

*When PURSUIT DG is soil applied, these weeds are more consistently controlled by preplant incorporated treatments.

DO NOT count cotyledon leaves when determining weed stage of growth.

GRASS WEEDS

Weed Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Barnyardgrass	R	3	1-3
Crabgrass, large	R	3	1-3
smooth	R	3	1-3
Cupgrass, woolly		3	1-3
Foxtail, giant	C	6	1-6
green	C	3	1-3
giant green	C	3	1-3
Goosegrass	R		
Johnsongrass, seedling	C	6	1-8
rhizome		R	1-8
Millet, wild proso	R	R	1-3
Panicum, fall	R		
Texas	R		

Red rice		3	1-3
Sandbur, field	R	R	≤1
Shattercane	R	6	1-8
Signalgrass, broadleaf	R	4	1-8
Sorghum almum	R	6	1-3

SEDGES

Nutsedge, purple	R	R	1-3
yellow	R	R	1-3

*When PURSUIT DG is applied to grasses, preplant incorporated treatments of PURSUIT DG are more consistent.

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence.)

ROTATIONAL CROPS

The following rotational crops may be planted after applying PURSUIT DG at the recommended rate: (Planting earlier than the recommended interval may result in crop injury).

1. Anytime
 - Lima beans
 - Southern peas
 - Soybeans
 - Peanuts
 - IMI-Corn™ (resistant/tolerant to PURSUIT DG)
2. Four months after PURSUIT DG application:
 - Alfalfa
 - Rye
 - Wheat
 - Edible beans and peas (other than lima beans and Southern peas)
3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
4. Nine and one-half months after PURSUIT DG application:
 - Barley
 - Tobacco

5. Eighteen months after PURSUIT DG application:

Cotton	Safflower
Lettuce	Sorghum
Oats	Sunflower
Popcorn	Sweet corn

6. Twenty six months after PURSUIT DG application.
Potatoes


7. Forty months after PURSUIT DG application.

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH). If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

*Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.





Supplemental Labeling

109/128

EPA Reg. No. 241-350

FOR USE IN SOYBEANS

TANK-MIXTURE OR SEQUENTIAL APPLICATIONS OF PURSUIT DG HERBICIDE AND PRESTIGE¹ HERBICIDE FOR ENHANCED GRASS CONTROL IN SOYBEANS

OBSERVE ALL PRECAUTIONARY AND WORKER PROTECTION STATEMENTS IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

In addition to broadleaf control, PURSUIT DG provides good to excellent control of several important grasses in soybeans. The addition of PRESTIGE herbicide to PURSUIT DG enhances the grass control, especially when heavy infestations of grass exist. It also provides control of grasses not controlled by PURSUIT DG. In some cases the activity of PRESTIGE may be reduced when mixed with PURSUIT DG. The reduction in activity may be overcome by delaying the application of PRESTIGE herbicide 7 days following the application of PURSUIT DG. If PRESTIGE is applied first, wait 3 days before applying PURSUIT DG.

For optimum control, apply the tank mixture to actively growing weeds at the sizes indicated in this Supplemental Labeling (or for sequential applications refer to application rates and weeds sizes indicated in the PURSUIT DG and PRESTIGE labels).

Refer to the PRESTIGE label for additional information regarding application rates, restrictions, precautions, weeds controlled, adjuvants recommended and other information. When tank-mixing herbicides, always follow the most restrictive label.

APPLICATION RATES

Apply PURSUIT DG at the rate of 1.44 ounces per acre. Refer to the table below for the appropriate rate of PRESTIGE herbicide for enhanced grass control.

ACCEPTED
~~with COMMENTS~~
In EPA Letter Dated

MAR 21 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

241-350

110/128

WEEDS CONTROLLED

The addition of PRESTIGE to PURSUIT DG at the recommended rates will control the grasses listed below. (Refer to the PRESTIGE label for other weeds controlled).

PRESTIGE Rate * (ounces per acre)	Annual Grasses Controlled	Size (inches)
12 oz.	Wild Proso Millet	4-10"
	Shattercane	3-12"
16 oz.	Foxtail, Giant	3-8"
	Junglerice	3-8"
	Panicum, Fall	3-8"
	Texas	3-8"
	Signalgrass, Broadleaf	3-8"
20 oz.	Volunteer Corn	4-10"
24 oz.	Barnyardgrass	3-8"
	Crabgrass, Large	3-6"
	Smooth	3-6"
	Cupgrass, woolly	3-8"
	Foxtail, Green	3-8"
	Yellow	3-8"
	Goosegrass	3-6"
	Johnsongrass, Seedling	3-8"
	Sprangletop, Red	3-8"
	Witchgrass	3-8"

*If a mixture of grasses are present, use the highest rate indicated for the grasses present.

MIXING INSTRUCTIONS

POSTEMERGENCE APPLICATIONS OF PURSUIT DG AND PRESTIGE REQUIRE THE ADDITION OF A CROP OIL CONCENTRATE AND A LIQUID FERTILIZER SOLUTION.

I. CROP OIL CONCENTRATE: A petroleum or vegetable seed based oil concentrate (such as SUN-IT II²) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other methylated seed oils at the rate of 1 to 1.5 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre.

AND

II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs. per acre.

This label must be in the possession of the user at the time of herbicide application.

¹Trademark of BASF Corporation.

²Trademark of Agsco, Inc.

Registered Trademark of American Cyanamid Company

PE-xxxxx 1/96 SD

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996

112/128

WEEDS CONTROLLED

The addition of PRESTIGE to PURSUIT DG at the recommended rates will control the grasses listed below. (Refer to the PRESTIGE label for other weeds controlled).

PRESTIGE Rate* (ounces per acre)	Annual Grasses Controlled	Size (inches)
12 oz.	Wild Proso Millet Shattercane	4-10" 3-12"
16 oz.	Foxtail, Giant Junglerice Panicum, Fall Texas Signalgrass, Broadleaf	3-8" 3-8" 3-8" 3-8" 3-8"
20 oz.	Volunteer Corn	4-10"
24 oz.	Barnyardgrass Crabgrass, Large Smooth Cupgrass, woolly Foxtail, Green Yellow Goosegrass Johnsongrass, Seedling Sprangletop, Red Witchgrass	3-8" 3-6" 3-6" 3-8" 3-8" 3-8" 3-6" 3-8" 3-8" 3-8"

*If a mixture of grasses are present, use the highest rate indicated for the grasses present.

MIXING INSTRUCTIONS

POSTEMERGENCE APPLICATIONS OF PURSUIT DG AND PRESTIGE REQUIRE THE ADDITION OF A CROP OIL CONCENTRATE AND A LIQUID FERTILIZER SOLUTION.

- I. **CROP OIL CONCENTRATE:** A petroleum or vegetable seed based oil concentrate (such as SUN-IT II²) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other methylated seed oils at the rate of 1 to 1.5 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre.

AND

II. **LIQUID FERTILIZER SOLUTION**

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs. per acre.

This label must be in the possession of the user at the time of herbicide application.



CYANAMID

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996

¹Trademark of BASF Corporation.

²Trademark of Agsco, Inc.

©Registered Trademark of American Cyanamid Company



Supplemental Labeling

113/128

NOT REVIEWED
EPA Reg. No. 241-350 In Accordance with PR Notice 82-2
Based On Draft Labeling Dated 3-21-96

~~FOR USE IN PEANUTS ONLY~~

FOR USE ONLY IN NORTH CAROLINA, SOUTH CAROLINA AND VIRGINIA

**COTTON PLANTING FOLLOWING PURSUIT DG APPLICATIONS TO PEANUTS -
ROTATIONAL CROP GUIDELINE MODIFICATION**

THIS LABEL EXPIRES MARCH 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

®Registered Trademark of American Cyanamid Company

(See reverse side)



174/128

ROTATIONAL CROPPING GUIDELINE

Cotton may be planted at the following intervals after a PURSUIT DG application if the criteria listed are met:

1. Nine and one-half months after an application of PURSUIT DG
 - PURSUIT DG applied to peanuts only, and
 - soil texture is sandy loam or loamy sand only, and
 - greater than 16 inches of rainfall and/or irrigation is received following application of PURSUIT DG through October of the application year.
 - Postemergence applications only. (Refer to the PURSUIT DG herbicide label for application instructions).

2. Eighteen months after an application of PURSUIT DG
 - If the above criteria are not met, rotation to cotton must be at least 18 months following an application of PURSUIT DG herbicide.

This label must be in the possession of the user at the time of herbicide application.



PE-17296 Rev. 1 1/96 SD



Supplemental Labeling

115/129

EPA Reg. No. 241-350 **NOT REVIEWED**
In Accordance with PR Notice 82-2
Based On Draft Labeling Dated 3.21.76
FOR USE IN SOYBEANS

TANK-MIXTURE OF PURSUIT DG + PINNACLE¹ HERBICIDES FOR POSTEMERGENCE CONTROL OF COMMON LAMBSQUARTERS IN SOYBEANS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.


GENERAL INFORMATION

For optimal weed control management, apply a soil applied grass herbicide such as PROWL[®], TRI-4[®], or Treflan² followed by PURSUIT DG postemergence. If common lambsquarters are not adequately controlled by the soil applied treatment, Pinnacle herbicide may be tank-mixed with PURSUIT DG for additional activity.

The addition of Pinnacle to PURSUIT DG may cause severe injury and/or stunting to soybeans, especially when applied under hot, humid conditions. The USER ASSUMES ALL RISKS AND CONSEQUENCES associated with applications of this tank-mixture to soybeans.

[®]Registered Trademarks of American Cyanamid Company
¹Trademark of E. I. duPont de Nemours and Company
²Trademark of Dow Elanco Company

(See reverse side)



APPLICATION RATES

116/128

When tank-mixing Pinnacle with PURSUIT DG, use the following rates:

PURSUIT DG - 1.44 ounces per acre

Pinnacle - 0.125 ounce per acre

Add to the spray mixture:

Non-ionic surfactant - 1 quart per 100 gallons (0.25% v/v)

AND

Liquid nitrogen based fertilizer (such as 28%N, 32%N, or 10-34-0) at the rate of 1-2 quarts per acre. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs./acres.

WEEDS CONTROLLED

This tank-mixture will improve the activity of PURSUIT DG on common lambsquarters. Refer to the PURSUIT DG label for additional weeds controlled.

RESTRICTIONS

Apply to 1-3 trifoliolate stage soybeans only.

Refer to both labels for information on methods of application, additional restrictions, precautions and rotational crop guidelines. Follow the more restrictive label.

This label must be in the possession of the user at the time of herbicide application.



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996



Supplemental Labeling

117/128

EPA Reg. No. 241-350

PURSUIT DG AND ROUNDUP¹ TANK-MIXTURES FOR WEED CONTROL IN NO-TILL SOYBEANS AND IN ROUNDUP READY¹ SOYBEANS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This labeling must be in the possession of the user at the time of herbicide application.

GENERAL INFORMATION

PURSUIT DG provides control of many broadleaf and grassy weeds when applied to the soil or postemergence to weeds. It also provides season-long control of many weeds. It can be applied no-till to soybeans either as a burndown prior to planting or postemergence to soybeans for weed control. Glyphosate containing products may be tank-mixed with PURSUIT DG for control of difficult weeds common to no-till production, such as marestail.

Roundup (or other glyphosate containing products) may be tank-mixed with PURSUIT DG for enhanced postemergence weed control in Roundup Ready soybeans.

BURNDOWN WEED CONTROL (No-till soybeans and stale seedbed)


Apply PURSUIT DG plus Roundup (or other glyphosate containing products) for burndown of weeds in no-till soybeans. This tank mixture may be applied prior to planting any soybean variety (including Roundup Ready soybeans). Roundup will enhance the control of weeds including marestail, common lambsquarters, common ragweed and waterhemp species. Refer to the glyphosate product label for recommended rates of application and weeds controlled.

ROUNDUP READY SOYBEANS (glyphosate resistant)

Roundup (or other glyphosate containing products) may be tank mixed with PURSUIT DG for postemergence weed control in Roundup Ready soybeans. Apply the combination of PURSUIT DG and glyphosate for general weed burndown and residual control of broadleaf and grassy weeds. Refer to the glyphosate product label for application rates and weeds controlled.

NOTE: Glyphosate containing products should be applied postemergence only to Roundup Ready soybean varieties (these varieties contain the glyphosate resistant gene). Use only glyphosate products that are registered for use on Roundup Ready soybeans.

(see other side)



118/128

PURSUIT DG may be tank-mixed with other herbicides (other than glyphosate) on Roundup Ready soybeans. Refer to the PURSUIT DG label for recommended herbicides. Refer to the other herbicide labels for weeds controlled and recommended applications rates.

RATES

Apply PURSUIT DG at the rate of 1.44 ounces per acre.

(In North Dakota and in Minnesota north of highway #210 apply 1.08 ounces per acre).

Refer to the Roundup (or other glyphosate product) label for recommended use rates.

MIXING INSTRUCTIONS

BURNDOWN OR POSTEMERGENCE APPLICATIONS REQUIRE THE ADDITION OF AN ADJUVANT AND FERTILIZER.

SURFACTANTS: Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 quart per 100 gallons.

OR

CROP OIL CONCENTRATE: If herbicides other than Roundup (glyphosate) are mixed with PURSUIT DG, a petroleum or vegetable based oil concentrate (such as SUN-IT II²) may be used instead of a surfactant. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or methylated seed oils at the rate of 1.5 to 2 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre. Do not use a crop oil concentrate when Roundup (or other glyphosate product) is tank-mixed with PURSUIT DG.

AND

FERTILIZER: Include a fertilizer in the spray solution. When Roundup (glyphosate) is used with PURSUIT DG, add spray grade ammonium sulfate at the rate of 2.5 lb. per acre. Instead of ammonium sulfate, a nitrogen-based liquid fertilizer (such as 28%N, 32%N or 10-34-0) may be used at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress.

Refer to the PURSUIT DG label and glyphosate product label for additional information regarding **WEEDS CONTROLLED, MIXING AND SPRAYING INSTRUCTIONS**. Refer to the Roundup label or other herbicide label for additional information regarding application rates, restrictions, precautions, weeds controlled, adjuvants recommended and other information.

When tank-mixing herbicides, always follow the more restrictive label.



¹Trademarks of Monsanto Agricultural Company
²Trademark of Agsco, Inc.
©Registered Trademark of American Cyanamid Company



Supplemental Labeling

NOT REVIEWED

EPA Reg. No. 241-350 In Accordance with PR Notice 82-2
Based On Draft Labeling Dated 3-21-76 119/128

~~PURSUIT DG AND ROUNDUP¹ TANK-MIXTURES FOR WEED CONTROL IN NO-TILL SOYBEANS~~ AND IN ROUNDUP READY¹ SOYBEANS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This labeling must be in the possession of the user at the time of herbicide application.

GENERAL INFORMATION

PURSUIT DG provides control of many broadleaf and grassy weeds when applied to the soil or postemergence to weeds. It also provides season-long control of many weeds. It can be applied no-till to soybeans either as a burndown prior to planting or postemergence to soybeans for weed control. Glyphosate containing products may be tank-mixed with PURSUIT DG for control of difficult weeds common to no-till production, such as marestalk.

Roundup (or other glyphosate containing products) may be tank-mixed with PURSUIT DG for enhanced postemergence weed control in Roundup Ready soybeans.

BURNDOWN WEED CONTROL (No-till soybeans and stale seedbed)


Apply PURSUIT DG plus Roundup (or other glyphosate containing products) for burndown of weeds in no-till soybeans. This tank mixture may be applied prior to planting any soybean variety (including Roundup Ready soybeans). Roundup will enhance the control of weeds including marestalk, common lambsquarters, common ragweed and waterhemp species. Refer to the glyphosate product label for recommended rates of application and weeds controlled.

ROUNDUP READY SOYBEANS (glyphosate resistant)

Roundup (or other glyphosate containing products) may be tank mixed with PURSUIT DG for postemergence weed control in Roundup Ready soybeans. Apply the combination of PURSUIT DG and glyphosate for general weed burndown and residual control of broadleaf and grassy weeds. Refer to the glyphosate product label for application rates and weeds controlled.

NOTE: Glyphosate containing products should be applied postemergence only to Roundup Ready soybean varieties (these varieties contain the glyphosate resistant gene). Use only glyphosate products that are registered for use on Roundup Ready soybeans.

(see other side)



PURSUIT DG may be tank-mixed with other herbicides (other than glyphosate) on Roundup Ready soybeans. Refer to the PURSUIT DG label for recommended herbicides. Refer to the other herbicide labels for weeds controlled and recommended applications rates.

RATES

Apply PURSUIT DG at the rate of 1.44 ounces per acre.

(In North Dakota and in Minnesota north of highway #210 apply 1.08 ounces per acre).

Refer to the Roundup (or other glyphosate product) label for recommended use rates.

MIXING INSTRUCTIONS

BURNDOWN OR POSTEMERGENCE APPLICATIONS REQUIRE THE ADDITION OF AN ADJUVANT AND FERTILIZER.

SURFACTANTS: Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 quart per 100 gallons.

OR


CROP OIL CONCENTRATE: If herbicides other than Roundup (glyphosate) are mixed with PURSUIT DG, a petroleum or vegetable based oil concentrate (such as SUN-IT II²) may be used instead of a surfactant. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or methylated seed oils at the rate of 1.5 to 2 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre. Do not use a crop oil concentrate when Roundup (or other glyphosate product) is tank-mixed with PURSUIT DG.

AND

FERTILIZER: Include a fertilizer in the spray solution. When Roundup (glyphosate) is used with PURSUIT DG, add spray grade ammonium sulfate at the rate of 2.5 lb. per acre. Instead of ammonium sulfate, a nitrogen-based liquid fertilizer (such as 28%N, 32%N or 10-34-0) may be used at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress.

Refer to the PURSUIT DG label and glyphosate product label for additional information regarding WEEDS CONTROLLED, MIXING AND SPRAYING INSTRUCTIONS. Refer to the Roundup label or other herbicide label for additional information regarding application rates, restrictions, precautions, weeds controlled, adjuvants recommended and other information.

When tank-mixing herbicides, always follow the more restrictive label.



¹Trademarks of Monsanto Agricultural Company

²Trademark of Agsco, Inc.

®Registered Trademark of American Cyanamid Company



121/128

Supplemental Labeling

NOT REVIEWED

EPA Reg. No. 241-350

In Accordance with PR Notice 82-2 **FOR USE IN ALFALFA ONLY**
Based On Draft Labeling Dated 3.21.96

For Use In California Only

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR MIXING AND APPLICATION INFORMATION, WORKER PROTECTION INFORMATION AND PRECAUTIONARY STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of herbicide application. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

SEEDLING ALFALFA

PURSUIT DG must be applied postemergence to seedling alfalfa. Apply PURSUIT DG when the seedling alfalfa is in the second (2nd) trifoliate stage or larger and when the majority of the weeds are 1-3 inches (see weed control tables on pages 4-6). For low growing weeds (such as mustards) apply PURSUIT DG before the rosette exceeds 3 inches. When PURSUIT DG is applied to seedling alfalfa, there may be a temporary reduction in growth. Alfalfa soon outgrows any effects of the herbicide.

ESTABLISHED ALFALFA

PURSUIT DG can be applied to established alfalfa in the fall, in the spring to dormant, or semi-dormant alfalfa (less than 3 inches of re-growth), or between cuttings. Any application should be made before significant alfalfa growth or re-growth (3 inches) to allow PURSUIT DG to reach the target weeds.

Replanting: If replanting is necessary in a field previously treated with PURSUIT DG, do not plant the field to alfalfa for 4 months following the PURSUIT DG application. Refer to the ROTATIONAL CROP GUIDELINE section on this label for plant-back interval of various crops.

MIXING INSTRUCTIONS

POSTEMERGENCE APPLICATIONS OF PURSUIT DG REQUIRE THE ADDITION OF AN ADJUVANT.

ADJUVANTS

SURFACTANTS: Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 quart per 100 gallons (0.25% V/V). An organo-silicone surfactant may be used in place of a non-ionic surfactant.

OR

CROP OIL CONCENTRATE: Instead of a surfactant, a petroleum or vegetable seed based oil concentrate (such as SUN-IT II¹) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other methylated seed oils at the rate of 1.5 to 2 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre.

122/128

Fill the spray tank one-fourth to one-half full with clean water. Use a calibrated measuring device to measure the required amount of PURSUIT DG. Add PURSUIT DG to the spray tank while agitating. Add adjuvants and fill the remainder of the tank with water.

TANK MIX COMBINATIONS WITH OTHER HERBICIDES

To control weeds not listed on the PURSUIT DG label, herbicides such as Buctril², 2,4-DB, POAST³ or POAST PLUS³ may be tankmixed with PURSUIT DG. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded.

SPRAYING INSTRUCTIONS

DO NOT apply when wind velocity is greater than 10 mph, or when spray may be carried to sensitive crops. Sensitive crops include leafy vegetables, cole crops, sugar beets, cotton, trees and vines.

GROUND APPLICATIONS

Uniformly apply with properly calibrated ground equipment in 10 or more gallons of water per acre. Use higher gallonage (i.e. 20 gallons of water per acre) for fields with dense vegetation or heavy weed pressure. A spray pressure of 20 to 40 psi is recommended.

Adjust the boom height to ensure proper coverage of weed foliage (according to the manufacturer's recommendation). Use only flat-fan nozzle tips only.

Avoid overlaps when spraying.

DO NOT apply by air in California.

APPLICATION INFORMATION

PURSUIT DG is effective in controlling a broad spectrum of broadleaf and grass weeds. Alfalfa has excellent tolerance to postemergence applications of PURSUIT DG after the second trifoliate leaf has expanded. Minor height reduction or slight leaf yellowing may occur soon after application. The effects are temporary and the alfalfa will soon recover.

Apply PURSUIT DG as an early postemergence treatment when the weeds are actively growing. Weeds are generally easier to control before they exceed 3 inches in height. Weeds under stress are less susceptible to control in cold or drought stress conditions.

PURSUIT DG should be applied a minimum of one hour before rainfall or overhead irrigation.

STAND ESTABLISHMENT

Apply PURSUIT DG after the alfalfa has 2 fully expanded trifoliate leaves. Weeds must not exceed the size listed in the WEEDS CONTROLLED tables. PURSUIT DG may be applied to summer, fall or spring seeded alfalfa.

Inter-seeded Oats

Oats inter-seeded with alfalfa will reduce soil erosion and allow the alfalfa to establish. Oats, however can compete with the alfalfa. An application of PURSUIT DG will kill or significantly reduce the growth of the oats and allow the alfalfa to establish with minimal erosion or competition from the oats. Apply PURSUIT to the oats when the oats have 3-4 leaves.

ESTABLISHED ALFALFA - DORMANT

PURSUIT DG may be applied to dormant alfalfa in the fall following the last cutting. PURSUIT DG may also be applied in the spring to dormant alfalfa, or as alfalfa breaks dormancy. Apply spring treatments prior to excessive alfalfa growth (less than 3 inches), to reduce spray interference.

ESTABLISHED ALFALFA - GROWING

For weed control during the season, apply PURSUIT DG following alfalfa cutting. Remove the hay from the field and apply PURSUIT DG prior to excessive alfalfa regrowth.

Perennial Grass Suppression

If perennial grasses (such as orchardgrass, fescues, bromes or timothy) are present in an alfalfa stand, PURSUIT DG will reduce the growth and competitive effect of the grass.

Do not apply to alfalfa during the last year of the stand.

USE RATE

(1.08 to 2.16 OUNCES PER ACRE)

Apply PURSUIT DG at a broadcast rate of 1.08 to 2.16 ounces per acre to seedling or established alfalfa. One soluble bag (7.2 ounces) will treat 5 acres at the rate of 1.44 ounces per acre.

NOTE: A maximum of 2.16 ounces of PURSUIT DG per year may be applied to alfalfa.

124/128

WEEDS CONTROLLED

When applied as directed, PURSUIT DG will control or reduce competition from the weeds listed below. Refer to the MIXING INSTRUCTIONS section on pages 1-2 for recommendations when weeds are at the maximum recommended growth stage, or are under stress.

NOTE: C = Control R = Reduced Competition

BROADLEAF WEEDS CONTROLLED

WEEDS 1-3 Inches*	PURSUIT DG APPLICATION RATE		
	1.08 Oz/A	1.44 Oz/A	2.16 Oz/A
Artichoke, Jerusalem	R	C	C
Beets, wild	C	C	C
Bedstraw, catchweed		C	C
Buckwheat, wild		C	C
Chickweed			
common	R	C	C
mouseear	R	C	C
Cocklebur, common	R	C	C
Dandelion		R	R
Dock,			
broadleaf (seedling)			R
curly (seedling)			R
Fiddleneck			R
Filaree			
redstem		R	C
whitestem		R	C
Flixweed	R	C	C
Goosefoot, nettleleaf	R	C	C
Grounsel, common			R
Henbit		R	C
Jimsonweed		C	C
Knotweed, prostrate		R	C
Kochia	R	C	C
Lambsquarter,			
common (1-2 leaves)		R	R
Lettuce, miners		C	C
Mallow,			
common		C	C
little		C	C
Marshelder		C	C
Morningglory,			
entireleaf			C
ivyleaf			C
pitted			C
smallflower		C	C
tall			C
Mustards			
tumble	C	C	C
wild	C	C	C
black	C	C	C
Nettle, burning		C	C

*Apply PURSUIT DG to weeds 1-3 inches tall or when rosettes are 1-3 inches wide. This will generally occur when weeds have 2-6 leaves.

125/128

BROADLEAF WEEDS CONTROLLED - Continued

WEEDS 1-3 Inches	PURSUIT DG APPLICATION RATE		
	1.08 Oz/A	1.44 Oz/A	2.16 Oz/A
Nightshade			
black	C	C	C
Eastern black	C	C	C
hairy	C	C	C
Oxtongue, bristly			R
Pennycress, field	C	C	C
Pepperweed,			
field	C	C	C
Virginia	R	C	C
Pigweed			
redroot	C	C	C
smooth	R	C	C
spiny		C	C
Radish, wild		R	C
Ragweed			
common		C	C
giant		C	C
Redmaids		C	C
Rocket			
London	C	C	C
yellow	R	C	C
Rockpurslane, desert			C
Shepherdspurse	C	C	C
Smartweed			
ladysthumb	R	C	C
Pennsylvania	R	C	C
swamp (seedling)		C	C
Spurge			
prostrate		R	C
spotted		R	C
petty		C	C
Spurry, corn		C	C
Sunflower, common	R	C	C
Swinecress		C	C
Tansymustard			
green	C	C	C
pinnate	C	C	C
Thistle, Russian	R	C	C
Velvetleaf	R	C	C
Wartcress, creeping		C	C
Watercress		C	C
Willowweed, panicle		C	C

126/128

GRASSES AND SEDGES CONTROLLED

WEEDS 1-3 Inches	PURSUIT DG APPLICATION RATE	
	1.44 Oz/A	2.16 Oz/A
Barnyardgrass	R	C
Bluegrass, annual		R
Canarygrass, littleseed	R	R
Cereals, volunteer		
barley	R	R
oats	R	R
wheat	R	R
Crabgrass,		
large	R	C
smooth	R	C
Cupgrass, woolly*	C	C
Foxtail,		
giant (1-6 leaves)	C	C
green	C	C
yellow	R	C
Johnsongrass,		
seedling (1-6 leaves)	C	C
rhizome	R	R
Millet, wild proso	R	C
Nutsedge,		
yellow	R	R
purple	R	R
Oats, wild	R	R
Rice, red	C	C
Shattercane (1-6 leaves)	C	C
Signalgrass, broadleaf	R	C
Quackgrass**		R

*PURSUIT DG controls emerged woolly cupgrass only.

**Quackgrass will be suppressed only when actively growing and before it exceeds 7 inches in height.

127/
128

ROTATIONAL CROP GUIDELINE

The following rotational crops may be planted after applying PURSUIT DG at the recommended rate: (Planting earlier than the recommended interval may result in crop injury).

1. Anytime.
 - Lima beans
 - Southern peas
 - Soybeans
 - Peanuts
 - IMI-Corn™ (resistant/tolerant to PURSUIT DG)
2. Four months after PURSUIT DG application:
 - Alfalfa
 - Rye
 - Wheat
 - Edible beans and peas (other than lima beans and Southern peas)
3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
4. Nine and one-half months after PURSUIT DG application:
 - Barley
 - Tobacco
5. Eighteen months after PURSUIT DG application:

Cotton	Lettuce
Oats	Popcorn
Safflower	Sorghum
Sunflower	Sweet corn
6. Twenty six months after PURSUIT DG application.
 - Potatoes
7. Forty months after PURSUIT DG application.*

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

*Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH). If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

EPA Reg. No. 241-350

FOR USE IN GREAT NORTHERN AND RED KIDNEY TYPE DRY BEANS

**FOR DISTRIBUTION AND USE ONLY IN COLORADO, KANSAS, NEBRASKA,
WYOMING, TEXAS AND WISCONSIN**

THIS LABEL EXPIRES APRIL 30, 1997

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

NOTICE

AMERICAN CYANAMID COMPANY
("CYANAMID") HAS OBTAINED A REGISTRATION
AS A SERVICE TO THOSE GROWERS WHO WISH
TO USE PURSUIT DG HERBICIDE AS AN AID IN
CONTROLLING WEEDS IN EDIBLE LEGUME
VEGETABLES. CYANAMID DOES NOT WARRANT
THE USE OF PURSUIT DG ON EDIBLE LEGUME
VEGETABLES. THE DECISION TO APPLY THIS
HERBICIDE MUST BE MADE BY THE GROWER
WITH THE UNDERSTANDING THAT REDUCED
CROP GROWTH, QUALITY, YIELD AND/OR
DELAYED MATURITY MAY RESULT EVEN WHEN
DIRECTIONS FOR USE ARE FOLLOWED
PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF
PURSUIT DG TO EDIBLE LEGUME VEGETABLE
CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL
FOR ADDITIONAL DISCLAIMER STATEMENTS.

GENERAL INFORMATION

Since crop maturity may be delayed, timing of harvest may
need to be adjusted accordingly. **DO NOT** apply PURSUIT
DG if planting is delayed and chance of frost prior to
maturity is likely

DO NOT apply PURSUIT DG if cold and/or wet conditions
are present or predicted to occur within one week of
application.

DO NOT apply PURSUIT DG postemergence after crop
has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices
have been utilized, including good soil fertility, proper crop
rotation, disease and insect management and tillage
practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label
directions is expected to result in normal growth of
rotational crops in most situations; however, various
environmental and agronomic factors make it impossible to
eliminate all risks associated with the use of this product
and, therefore, rotational crop injury is always possible.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a
manner inconsistent with its labeling. **DO NOT** apply this
product through any type of irrigation system. This label
must be in the possession of the user at the time of herbicide
application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not
specifically described in this label.

ACCEPTED
with COMMENTS
In EPA Letter Dated

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
amended, the pesticide
registered under EPA Reg. No.

241-350

*Registered Trademark of American Cyanamid Company

be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after June 30. Allow at least 60 days between application and harvest.

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. **DO NOT** apply PURSUIT DG on sand or loamy sand soils.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Basagran¹ may be tank-mixed with PURSUIT DG to control weeds not listed on the PURSUIT DG label. Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use only nonionic surfactants as a spray additive.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preemergence or early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Soybeans
 - Peanuts
 - Edible bean types listed on this label (Refer to heading on page 1).
 - IMI-Corn™
- Four months after PURSUIT DG application:
 - Barley
 - Edible bean types not listed on this label (Refer to heading on page 1).
 - Peas
 - Rye
 - Wheat
- Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
- Nine and one-half months after PURSUIT DG application:
 - Tobacco

¹Dash and Basagran are trademarks of BASF Corporation

™Trademark of American Cyanamid Company

157
1

ROTATIONAL CROP GUIDELINE (continued)

5. Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet Corn
Popcorn	

6. Twenty-six months after PURSUIT DG application:

Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



EPA Reg. No. 241-350

31751
r

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,
CRANBERRY, AND SMALL WHITE TYPE DRY BEANS ONLY**

**FOR DISTRIBUTION AND USE IN MICHIGAN ONLY
(EXCLUDING UPPER PENINSULA)**

THIS LABEL EXPIRES APRIL 30, 1997

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate competition and weeds.

ACCEPTED
**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
241-350**

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

*Registered Trademark of American Cyanamid Company

DO NOT make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply at any timing to Domino variety black turtle beans.

DO NOT apply PURSUIT DG to sand or loamy sand soils.

DO NOT apply PURSUIT DG after July 10.

Allow at least 60 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

Soil Type	Use Rate (oz/acre)
Sands, loamy sands	DO NOT USE
Sandy loams and all other soils with \leq 2% O.M.	0.72
All other soils with $>$ 2% O.M.	1.08

Preemergence Applications. Apply PURSUIT DG at the rates shown in the preceding table immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

weeds not listed on the PURSUIT DG label. Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use only nonionic surfactants as a spray additive.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
0.72	10 acres
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied to low organic matter soils at the broadcast rate of 0.72 ounces per acre preplant incorporated, preemergence, or postemergence will control light to moderate infestations of:

Nightshade, black
Nightshade, Eastern black
Pigweed, redroot

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild
Nightshade, black
Nightshade, Eastern black
Nightshade, hairy
Pigweed, redroot

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label.

- Anytime:
 - Soybeans
 - Peanuts
 - IMI-Corn™
 - Edible bean and pea types listed on this label (Refer to heading on page 1)

¹Dach and Basagran are trademarks of BASF Corporation.

²™ Trademark of American Cyanamid Company.

ROTATIONAL CROP GUIDELINE (continued)

2. Four months after PURSUIT DG application:

- Barley
- Edible bean and pea types not listed on this label (Refer to heading on page 1)
- Rye
- Wheat

3. Eight and one-half months after PURSUIT DG application:

- Field corn
- Field corn grown for seed*

4. Nine and one-half months after PURSUIT DG application:

- Tobacco

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet corn |
| Popcorn | |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



341 51

EPA Reg. No. 241-350

FOR POSTEMERGENCE USE IN PINTO TYPE DRY BEANS

**FOR DISTRIBUTION AND USE ONLY IN COLORADO, KANSAS, WYOMING, NEBRASKA,
AND TEXAS**

THIS LABEL EXPIRES APRIL 30, 1997

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

ACCEPTED
~~with COMMENTS~~
In EPA Letter Dated

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
241-350

*Registered Trademark of American Cyanamid Company

GENERAL INFORMATION

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

Some varieties of pinto beans may be inherently less tolerant to PURSUIT DG applications and crop injury could result. Contact the seed company for information and recommendations about postemergence applications of PURSUIT DG.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible

ROTATIONAL CROP GUIDELINE (continued)

5. Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	

6. Twenty-six months after PURSUIT DG application:

Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss Great Northern or red kidney beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



MAR 21 1995
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

EPA Reg. No. 241-350

377 51

**FOR USE IN NAVY, GREAT-NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY,
PINTO, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, DRY EDIBLE PEAS,
241-240 ENGLISH PEAS, LENTILS, AND CHICKPEAS**

**FOR DISTRIBUTION AND USE IN MINNESOTA (south of highway #210)
AND SOUTH DAKOTA**

THIS LABEL EXPIRES APRIL 30, 1997

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

NOTICE

AMERICAN CYANAMID COMPANY
("CYANAMID") HAS OBTAINED A REGISTRATION
AS A SERVICE TO THOSE GROWERS WHO WISH
TO USE PURSUIT DG HERBICIDE AS AN AID IN
CONTROLLING WEEDS IN EDIBLE LEGUME
VEGETABLES. CYANAMID DOES NOT WARRANT
THE USE OF PURSUIT DG ON EDIBLE LEGUME
VEGETABLES. THE DECISION TO APPLY THIS
HERBICIDE MUST BE MADE BY THE GROWER
WITH THE UNDERSTANDING THAT REDUCED
CROP GROWTH, QUALITY, YIELD AND/OR
DELAYED MATURITY MAY RESULT EVEN WHEN
DIRECTIONS FOR USE ARE FOLLOWED
PRECISELY.

DO NOT apply PURSUIT DG if cold and/or wet conditions
are present or predicted to occur within one week of
application.

DO NOT apply PURSUIT DG postemergence after crop
has begun to flower or crop injury may result.

Some crop varieties may be inherently less tolerant to
PURSUIT DG applications and crop injury could result.
Contact the seed company for information and
recommendations about the use of PURSUIT DG
postemergence.

USE PURSUIT DG ONLY if proper agronomic practices
have been utilized, including good soil fertility, proper crop
rotation, disease and insect management and tillage
practices that eliminate compaction and hardpans.

DISCLAIMER

**THE USER ASSUMES ALL RISKS AND
CONSEQUENCES ASSOCIATED WITH APPLICA-
TIONS OF PURSUIT DG TO EDIBLE LEGUME
VEGETABLE CROPS.**

**REFER TO THE PURSUIT DG LEAFLET LABEL
FOR ADDITIONAL DISCLAIMER STATEMENTS.**

GENERAL INFORMATION

Since crop maturity may be delayed, timing of harvest may
need to be adjusted accordingly. **DO NOT** apply PURSUIT
DG if planting is delayed and chance of frost prior to
maturity is likely.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a
manner inconsistent with its labeling. **DO NOT** apply this
product through any type of irrigation system. This label
must be in the possession of the user at the time of herbicide
application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not
specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and
leafy vegetables may be injured by spray drift or other
indirect contact with PURSUIT DG

*Registered Trademark of American Cyanamid Company

fertilizers or Dash.¹

DO NOT apply to Domino variety black turtle beans.

DO NOT make more than one application of PURSUIT DG per year.

Allow at least 30 days between application and harvest of lima beans and English peas. Allow at least 60 days between application and harvest of dry beans, dry edible peas, lentils and chickpeas.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION INSTRUCTIONS

DO NOT apply PURSUIT DG to sand or loamy sand soils.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre within 1 week before planting. **DO NOT** incorporate deeper than 3 inches. Refer to the PURSUIT DG leaflet label for incorporation instructions.

DO NOT apply preplant incorporated to pinto beans.

Preemergence Applications: Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre after planting, but prior to crop emergence. If weeds are present at the time of application, a burndown herbicide such as Roundup² or Gramoxone Extra³ should be tank-mixed with PURSUIT DG to control emerged weeds.

DO NOT apply preemergence to pinto beans.

The nonionic surfactant must contain at least 60% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. **DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.**

ECO-PAK CONVERSION TABLE

<u>Use Rate (oz/acre)</u>	<u>1 ECO-PAK Soluble Bag Treats</u>
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence or postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Chickpeas
 - Lentils
 - Dry edible peas
 - Peanuts
 - Edible beans (dry and lima listed on this label)
 - Soybeans
 - IMI-Corn™
- Four months after PURSUIT DG application:
 - Barley
 - Edible beans (dry, snap) and peas not listed on this label
 - Rye
 - Wheat
- Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*

¹Dash is a trademark of BASF Corporation

²Roundup is a trademark of Monsanto Agricultural Products Company

³Gramoxone Extra is a trademark of Zeneca, Inc

*Trademark American Cyanamid Company

23

ROTATIONAL CROP GUIDELINE (continued)

4. Nine and one-half months after PURSUIT DG application:

Tobacco

5. Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	

6. Twenty-six months after PURSUIT DG application:

Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, edible beans, dry edible peas, lentils or chickpeas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



107 51

EPA Reg. No. 241-350

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY,
BLACK TURTLE, CRANBERRY, AND PINTO TYPE DRY BEANS,
AND DRY EDIBLE PEAS ONLY**

**FOR DISTRIBUTION AND USE IN NORTH DAKOTA AND
NORTHERN MINNESOTA (see USE AREA below)**

THIS LABEL EXPIRES APRIL 30, 1997

**) OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

NOTICE

GENERAL INFORMATION

AMERICAN CYANAMID COMPANY
("CYANAMID") HAS OBTAINED A REGISTRATION
AS A SERVICE TO THOSE GROWERS WHO WISH
TO USE PURSUIT DG HERBICIDE AS AN AID IN
CONTROLLING WEEDS IN EDIBLE LEGUME
VEGETABLES. CYANAMID DOES NOT WARRANT
THE USE OF PURSUIT DG ON EDIBLE LEGUME
VEGETABLES. THE DECISION TO APPLY THIS
HERBICIDE MUST BE MADE BY THE GROWER
WITH THE UNDERSTANDING THAT REDUCED
CROP GROWTH, QUALITY, YIELD AND/OR
DELAYED MATURITY MAY RESULT EVEN WHEN
**) DIRECTIONS FOR USE ARE FOLLOWED
PRECISELY.**

PURSUIT DG application methods and use rates vary for
different types of edible legume vegetables. Additionally,
geographical restrictions apply. Apply PURSUIT DG only
to the legume vegetables specifically listed in this label.
DO NOT apply PURSUIT DG in geographical regions
outside those specifically defined in this label for each type
of legume vegetable.

Since crop maturity may be delayed, timing of harvest may
need to be adjusted accordingly. **DO NOT** apply PURSUIT
DG if planting is delayed and chance of frost prior to
maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions
are present or predicted to occur within one week of
application.

DO NOT apply PURSUIT DG postemergence after crop
has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices
have been utilized, including good soil fertility, proper crop
rotation, disease and insect management and tillage
practices that eliminate compaction and hardpans.

Under some conditions (such as heavy texture soil, high
organic matter, low pH or low rainfall) PURSUIT DG may
cause injury to subsequent planted crops. Sugar beets are
particularly sensitive to PURSUIT DG residues in soil.
PURSUIT DG is not recommended in crop rotations that
include sugar beets. See the ROTATIONAL CROP section
of this supplemental label for rotations to other sensitive
crops.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF
PURSUIT DG TO EDIBLE LEGUME VEGETABLE
CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL
FOR ADDITIONAL DISCLAIMER STATEMENTS.

ACCEPTED
~~with COMMENTS~~
In EPA Letter Dated

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

*Registered Trademark of American Cyanamid Company

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply at any timing to Domino variety black turtle beans.

Pinto varieties UI-111 and Olathe are more sensitive to PURSUIT DG than other varieties.

DO NOT apply PURSUIT DG after July 10

DO NOT feed treated forage, hay, or straw to livestock

Allow at least 60 days between application and harvest

gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 0.72 ounces per acre will control:

wild mustard
black nightshade*
eastern black nightshade*

*suppression

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
Soybeans
IMI-Corn™
Edible bean and pea types listed on this label
(Refer to heading on page 1)
2. Eight and one-half months after PURSUIT DG application:
Field corn
Field corn grown for seed*
3. Nine and one-half months after PURSUIT DG application:
Wheat
Edible bean and pea types not listed on this label
(Refer to heading on page 1)
4. Eighteen months after PURSUIT DG application
Barley Rye
Cotton Safflower
Lettuce Sorghum
Oats Sunflower
Popcorn Sweet Corn

¹Dash is a trademark of BASF Corporation.

™ Trademark of American Cyanamid Company.

ROTATIONAL CROP GUIDELINE (continued)

5. Twenty-six months after PURSUIT DG application:

- Potatoes
- Flax

6. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



1137 51

EPA Reg. No. 241-350

FOR USE IN SOYBEANS

**PURSUIT DG PLUS SCEPTER® OR SCEPTER® DG HERBICIDES
FOR SUPPRESSION OF VOLUNTEER CORN
AND ENHANCED CONTROL OF SUNFLOWERS**

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR WORKER PROTECTION STANDARD REQUIREMENTS, USE AREA, MIXING AND APPLICATION INFORMATION, AND WEEDS CONTROLLED.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

GENERAL INFORMATION

PURSUIT DG may be tank mixed with SCEPTER or SCEPTER DG for suppression of volunteer corn or for enhanced control of common sunflower in soybeans. Apply postemergence to soybeans only. In this label, SCEPTER refers to both formulations SCEPTER and SCEPTER DG. Be sure to refer to the correct label for additional information.

USE AREA

The application of PURSUIT DG plus SCEPTER or SCEPTER DG may be applied to states or portions of states described as Region 2 or Region 3 on the SCEPTER or SCEPTER DG labels, and the following counties in South Dakota: Yankton, Bon Homme, Hutchinson, McCook, Hanson, Davison, Miner, Lake and Kingsbury. Refer to the respective labels for the recommended use area. Do not use this tank mixture in North Dakota or in Minnesota north of state highway #210.

APPLICATION RATES

Apply the products at the following rate:

PURSUIT DG	1.44 oz/A*
AND	
SCEPTER	2 oz/A (liquid)
OR	
SCEPTER DG	0.53 oz/A**

ACCEPTED
with ~~COMMENTS~~
In EPA Letter Dated
MAY 11 1988

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under Reg. No.
241-350

*At the rate of 1.44 ounces per acre, one 7.2 ounce soluble bag of PURSUIT DG will treat 5 acres.

**At the rate of 0.53 ounces per acre, one 14 ounce soluble bag of SCEPTER DG will treat 26.4 acres.

®Registered Trademarks of American Cyanamid Company

* Field corn may be planted 9 1/2 months following a postemergence application of PURSUIT DG when applied in tank-mixture with SCEPTER or SCEPTER DG at the recommended rates unless extreme drought conditions develop (less than 12 inches of rainfall or irrigation is received from two weeks prior to the date of PURSUIT DG application through November 15 of the same year). If the minimum rainfall requirement is not met, only field corn hybrids (IMI-Corn™) which possess tolerance or resistance to PURSUIT DG may be planted the spring of the year following a PURSUIT DG + SCEPTER application.

Use of PURSUIT DG and SCEPTER herbicides in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of these products and, therefore, rotational crop injury is always possible.

Except as stated above, DO NOT use this tank mixture in the same year as an application of SCEPTER, SCEPTER DG, SQUADRON®, SQUADRON® DG TRI-SCEPT®, SCEPTER® O.T.®, STEEL™ CP, DETAIL®, PURSUIT®, PURSUIT® DG, PURSUIT® PLUS EC, CONTOUR® or RESOLVE®. Also, DO NOT use the above tank mixture in the same year as an application of a product containing chlorimuron ethyl (e.g. Classic¹, Canopy¹, Concert¹, Gemini¹, Lorcx Plus¹, Preview¹, Reliance¹, Synchrony¹ etc.)

®/™ Trademarks of American Cyanamid Company

¹ Trademarks of E.I. duPont de Nemours & Company, Inc.

PE-xxxxx 1/96 SD

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Division
Wayne, NJ 07470 ©1996

EPA Reg. No. 241-350

**FOR USE IN FIELD CORN
(APPLY ONLY ON IMI-CORN™ INBREDS)**

**USE DIRECTIONS FOR THE USE OF PURSUIT DG FOR WEED CONTROL IN
IMI-CORN INBREDS**

OBSERVE ALL PRECAUTIONARY STATEMENTS ON THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

Apply PURSUIT DG herbicide only on selected field corn inbreds (IMI-Corn) warranted by the seed company to possess resistance/tolerance to direct application of PURSUIT DG. DO NOT apply PURSUIT DG to corn inbreds which lack genetic resistance/tolerance to PURSUIT DG herbicide. Contact your chemical dealer, seed supplier or American Cyanamid to obtain information regarding IMI-Corn inbreds.

PURSUIT DG can be applied as an early preplant, preplant incorporated, preemergence or postemergence treatment on IMI-Corn inbreds.

Crops growing under stressful environmental conditions can exhibit various injury symptoms which may be more pronounced if herbicides are used. Corn plants treated with PURSUIT DG may exhibit yellowing on new growth. Such effects occur infrequently and are temporary. Normal growth and appearance should resume within 1 to 2 weeks.

Naturally occurring biotypes* of some of the weeds listed on this label may not be effectively controlled by this and/or other products with either the ALS/AHAS enzyme inhibiting mode of action. Other herbicides with the ALS/AHAS enzyme inhibiting mode of action include

the sulfonylureas (e.g., Accent™, Basis™, Classic™, Concert™, Exceed™, Permit™, Pinnacle™, etc.), the sulfonamides (e.g., Broadstrike™, etc.) and the pyrimidyl benzoates (e.g. Staple™, etc.). If naturally occurring ALS/AHAS resistant biotypes are present in a field, Pursuit DG and/or any other ALS/AHAS enzyme inhibiting mode of action herbicide should be tank-mixed or applied sequentially with an appropriate registered herbicide having a different mode of action to ensure control.

* A weed biotype is a naturally occurring plant within a given species that has a slightly different, but distinct, genetic makeup from other plants.

Replanting: If replanting is necessary in a field previously treated with PURSUIT DG, the field may be replanted to IMI-Corn, soybeans, peanuts, lima beans or southern peas. Rework the soil no deeper than the treated zone. DO NOT apply a second treatment of PURSUIT DG herbicide.

MIXING INSTRUCTIONS

Fill the spray tank one-fourth to one-half full with clean water. Add the required number of PURSUIT DG soluble packets to the spray tank while agitating. Add adjuvants and fill the remainder of the tank with water.

**TANK MIX COMBINATIONS WITH OTHER
HERBICIDES**

If other herbicides are tank-mixed with PURSUIT DG, while agitating, add components in the following order:

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

MAR 21 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241-350

- in soluble packets.
4. Add PURSUIT DG and thoroughly mix.
 5. Add other aqueous solution products.
 6. Add EC (emulsifiable concentrate) products.
 7. Add surfactant or crop oil to the spray tank.
 8. Add liquid fertilizer.
 9. While agitating, fill the remainder of the tank with water.

If PURSUIT DG is applied in liquid fertilizer as a carrier, PURSUIT DG must first be dissolved in a water slurry. Premix PURSUIT DG in a minimum of one gallon of water per ECO-PAK prior to adding to the spray tank.

DO NOT add PURSUIT DG ECO-PAK packets directly into liquid fertilizer.

USE RATE
(1.44 OUNCES PER ACRE)
(1 ECO-PAK soluble bag treats 5 acres)

Apply PURSUIT DG at a broadcast rate of 1 soluble bag per 5 acres (7.2 ounces) for all methods of application: early preplant, preplant incorporated, preemergence, and postemergence (including minimum and no-till). At this broadcast rate, one soluble bag PURSUIT DG will treat 5 acres of IMI-Corn.

NOTE: Only one application of PURSUIT DG may be made during the season.

PURSUIT DG HERBICIDE USE AREA
(Not for use in California)

DO NOT apply PURSUIT DG herbicide to IMI-Corn in North Dakota or in Minnesota north of state highway 210.

APPLICATION INSTRUCTIONS
POSTEMERGENCE

Apply PURSUIT DG herbicide as a postemergence treatment to IMI-Corn when crop and weeds are actively growing. Apply PURSUIT DG before weeds exceed a height of 3 inches, unless otherwise indicated (see PURSUIT DG leaflet label). More restrictive crop growth stage limitations of tank-mix partners must be followed.

ADDITIVES:

POSTEMERGENCE APPLICATIONS OF PURSUIT DG REQUIRE THE ADDITION OF AN ADJUVANT AND A LIQUID FERTILIZER¹ SOLUTION.

the surfactant at the rate of 1 quart per 100 gallons of spray solution.

OR

CROP OIL CONCENTRATE: Instead of a surfactant, a petroleum or vegetable seed oil based crop oil concentrate (such as SUN-IT II™) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other methylated seed oils at the rate of 1.5 - 2 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre.

AND

II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs per acre.

DO NOT apply PURSUIT DG herbicide with liquid fertilizer as a carrier for postemergence applications.

SOIL APPLICATIONS

PURSUIT DG can be applied as an early preplant, preplant incorporated, or preemergence treatment in IMI-Corn.

PURSUIT DG can be applied to the soil in liquid fertilizers, alone or in combination with PROWL® 3.3EC, Lasso EC™, Lasso MT™ or Dual™. Follow all PURSUIT DG label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. Always test the compatibility of PURSUIT DG herbicide with the liquid fertilizer before mixing in the spray tank.

¹ PURSUIT DG applications may be made with a non-ionic surfactant only (liquid fertilizer not required) in the state of Arkansas, Bootheel of Missouri, Tennessee, South Carolina, Louisiana, Mississippi, Alabama, and Georgia.

**SEQUENTIAL HERBICIDE
COMBINATIONS WITH PURSUIT DG**

**PURSUIT DG FOLLOWING EARLY PREPLANT
APPLICATION OF:**

2,4-D	Harness™
Bicep™, Bicep Lite™	Harness Extra
Bullet™	Lasso EC
Bronco™	Lasso MT
Dual	Roundup™
Extrazine II™	Surpass™
Frontier™	Surpass 100
Gramoxone Extra™	

**PURSUIT DG FOLLOWING PREPLANT
INCORPORATED APPLICATION OF:**

Atrazine ²	Harness
Bicep, Bicep Lite	Harness Extra
Bullet	Lariat™
Dual	Lasso EC, Lasso MT
Eradicane Extra™	Sutan+™
Extrazine II	Surpass
Frontier	Surpass 100

**PURSUIT DG FOLLOWING PREEMERGENCE
APPLICATION OF:**

Atrazine	Frontier
Banvel™, ²	Harness
Bicep, Bicep Lite	Harness Extra
Bladex™ 4L	Lariat
Bladex 90DF	Lasso EC, Lasso MT
Bronco	Marksman™, ²
Bullet	PROWL 3.3 EC
Clarity™, ²	Surpass
Dual	Surpass 100
Extrazine	

**TANK-MIX HERBICIDE
COMBINATIONS WITH PURSUIT DG**

**EARLY PREPLANT APPLICATION OF PURSUIT
DG TANK-MIXED WITH:**

2,4-D	Harness
Bronco	Lasso EC
Dual	Lasso MT
Frontier	Roundup
Gramoxone Extra	Surpass

**PREPLANT INCORPORATED APPLICATION OF
PURSUIT DG TANK-MIXED WITH:**

Atrazine	Frontier
Bicep, Bicep Lite	Harness
Bullet	Lariat
Dual	Lasso EC, Lasso MT
Eradicane Extra	Surpass
Extrazine	Sutan+

**PREEMERGENCE APPLICATION OF PURSUIT
DG TANK-MIXED WITH:**

Atrazine	Frontier
Banvel	Harness
Bicep, Bicep Lite	Harness Extra
Bladex 4L	Lariat
Bladex 90DF	Lasso EC, Lasso MT
Bronco	Marksman
Bullet	PROWL 3.3 EC
Clarity	Surpass
Dual	Surpass 100
Extrazine	

**POSTEMERGENCE APPLICATIONS OF PURSUIT
DG TANK-MIXED WITH:**

Atrazine ³	Frontier
Banvel ⁴	Lasso EC, Lasso MT
Basagran™, ²	Marksman
Buctril™, ^{2,3}	PROWL 3.3 EC
Clarity ⁴	Sencor DF™
Dual	Sencor 4L™

²In some cases the grass activity of PURSUIT DG will be reduced when used in combination with atrazine, Buctril, Banvel, Basagran, Clarity, or Marksman.

³Some corn leaf burn may result with Buctril or atrazine postemergence combinations with PURSUIT DG.

⁴Applications of Banvel or Clarity herbicide to corn during periods of rapid growth may result in temporary leaning.

DO NOT use crop oil concentrates as adjuvants in PURSUIT DG herbicide combinations with Banvel, Buctril, Clarity, Marksman, Sencor DF or Sencor 4L.

PURSUIT DG is active against many broadleaf and grass species. However, for long term weed management, alternate mode of action herbicides are recommended with PURSUIT DG. The application of a soil applied grass herbicide underlay will control grass weeds not on the PURSUIT DG label and enhance the control of certain broadleaf weeds such as common lamb-quarters.

When PURSUIT DG herbicide is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions.

AERIAL APPLICATIONS

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre. When applied POSTEMERGENCE, the addition of a non-ionic surfactant AND fertilizer solution are required for optimum weed control. Apply a non-ionic surfactant at the rate of 1 quart per 100 gallons of spray mixture OR a crop oil concentrate at the rate of 1.5 - 2 pints per acre AND a liquid fertilizer at the rate of 1 quart per acre. (See instructions under APPLICATION INSTRUCTIONS - POSTEMERGENCE).

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS.

- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph. Coarse sprays (larger droplets) are less likely to drift.

Applicator is responsible for any loss or damage which results from spraying PURSUIT DG in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

When applied as directed at the broadcast rate of 1.44 ounces per acre, PURSUIT DG will control or reduce competition from the weeds listed below.

NOTE: C = Control R = Reduced Competition

BROADLEAF WEEDS

Weed Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Alligator weed		4	1-3
Anoda, spurred	C	2	1-2
Artichoke, Jerusalem		8	6-10
Buffalobur	C*	R	1-3
Bristly starbur		2	1-2
Carpetweed	C		
Cocklebur, common	R	8	1-8
Galinsoga	C		
Jimsonweed	C*	4	1-3
Kochia	C	4	1-3
Lambsquarters, common	C*	R	1-2
Mallow, Venice	R		
Marshelder	C	4	1-3
Morningglory, entireleaf	R	2	1-2
ivyleaf	R	2	1-2
pitted	R	2	1-2
smallflower	C	4	1-3
tall	R	2	1-2
Mustard sp.	C	4	1-3

Nightshade, black	C	4	1-3
Eastern black hairy	C	4	1-3
Pigweed, Palmer	C	8	1-8
redroot	C	8	1-8
smooth	C	8	1-8
spiny	C	8	1-8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C*		
Ragweed, common	R	4	1-3
giant	R	4	1-3
Sage, barnyard		R	1-3
Smartweed, ladythumb	C	4	1-3
Pennsylvania	C	4	1-3
Spurge, prostrate	C	4	1-3
spotted	C	4	1-3
Sunflower	C*	4	1-3
Velvetleaf	C*	4	1-3
Thistle, Canada		R	1-3

*When PURSUIT DG is soil applied, these weeds are more consistently controlled by preplant incorporated treatments.

DO NOT count cotyledon leaves when determining weed stage of growth.

GRASS WEEDS

Weed Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Barnyardgrass	R	3	1-3
Crabgrass, large	R	3	1-3
smooth	R	3	1-3
Cupgrass, woolly		3	1-3
Foxtail, giant	C	6	1-6
green	C	3	1-3
giant green	C	3	1-3
Goosegrass	R		
Johnsongrass, seedling	C	6	1-8
rhizome		R	1-8
Millet, wild proso	R	R	1-3
Panicum, fall	R		
Texas	R		

Red rice		3	1-3
Sandbur, field	R	R	≤1
Shattercane	R	6	1-8
Signalgrass, broadleaf	R	4	1-8
Sorghum alnum	R	6	1-3

SEDGES

Nutsedge, purple	R	R	1-3
yellow	R	R	1-3

*When PURSUIT DG is applied to grasses, preplant incorporated treatments of PURSUIT DG are more consistent.

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence.)

ROTATIONAL CROPS

The following rotational crops may be planted after applying PURSUIT DG at the recommended rate: (Planting earlier than the recommended interval may result in crop injury).

1. Anytime
 - Lima beans
 - Southern peas
 - Soybeans
 - Peanuts
 - IMI-Corn™ (resistant/tolerant to PURSUIT DG)
2. Four months after PURSUIT DG application:
 - Alfalfa
 - Rye
 - Wheat
 - Edible beans and peas (other than lima beans and Southern peas)
3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
4. Nine and one-half months after PURSUIT DG application:
 - Barley
 - Tobacco

5. Eighteen months after PURSUIT DG application:

Cotton	Safflower
Lettuce	Sorghum
Oats	Sunflower
Popcorn	Sweet corn

6. Twenty six months after PURSUIT DG application.
Potatoes

7. Forty months after PURSUIT DG application.
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH). If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

*Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

PRECAUTIONS

In the event of a crop loss due to weather, IMI-Corn inbreds and hybrids, soybeans, lima beans, southern peas or peanuts can be replanted if PURSUIT DG herbicide was applied alone at rates labelled for these crops. If PURSUIT DG herbicide was tank-mixed with other herbicides, such as atrazine, the label restrictions for these herbicides must also be followed.

There should be an interval of at least 45 days between an application of PURSUIT DG and corn harvest (silage, fodder, or grain). DO NOT graze or feed treated corn, forage, silage, fodder, or grain for at least 45 days after an application of PURSUIT DG.

All soil insecticides, including labelled banded or in-furrow applications, may be used in combination with Pioneer imidazolinone resistant (IR) corn inbreds.

Imidazolinone tolerant inbreds from other seed companies may occasionally exhibit injury symptoms when soil insecticides are used in combination with PURSUIT DG. USE banded applications of COUNTER® 15G, COUNTER® 20CR® or THIMET® in combination with PURSUIT DG herbicide application; DO NOT USE COUNTER 15G soil systemic insecticide products in-furrow with imidazolinone tolerant inbreds. American Cyanamid has not tested all inbreds in which the imidazolinone tolerance trait is claimed and cannot be responsible for factors which are beyond its control, such as growing conditions, environmental conditions, grower practices and the specific genetics of each inbred which may influence tolerance. Growers should contact the seed company for information regarding specific inbred tolerance to herbicide and insecticide applications.

This label must be in the possession of the user at the time of herbicide application.

For additional information regarding the use of PURSUIT DG herbicide, call telephone no. (800) 942-0500.

PE-xxxx 3/96 SD

- ® Registered Trademark of American Cyanamid Company
- ™ SUN-IT II is a Trademark of Agsco, Inc.
- ™ Buctril is a Trademark of Rhone Poulenc Ag Company.
- ™ Banvel, Clarity, Frontier, and Marksman are Trademarks of Sandoz Crop Protection Corporation.
- ™ Basagran is a Trademark of BASF Corporation.
- ™ Beacon, Bicep, Bicep Lite, Dual, and Exceed are Trademarks of Ciba Geigy Corporation.
- ™ Bronco, Bullet, Harness, Harness Extra, Lariat, Lasso EC, Lasso MT, Permit and Roundup are Trademarks of Monsanto Agricultural Products Company.
- ™ Sencor is a Trademark of Mobay Corporation.
- ™ Gramoxone Extra, Eradicane Extra, Surpass, Surpass 100, and Sutan+ are Trademarks of Zeneca Agrochemicals.
- ™ Accent, Basis, Bladex 4L, Bladex 90DF, Classic, Concert, Extrazine, Pinnacle, and Staple are Trademarks of E. I. DuPont de Nemours and Company, Inc.
- ™ Imi-Corn is a Trademark of American Cyanamid Company
- ™ Broadstrike is a Trademark of Dow Elanco Corp.



141 25 241-350 12/1/81

EPA Reg. No. 241-350

FOR USE IN SOYBEANS

TANK-MIXTURE OR SEQUENTIAL APPLICATIONS OF PURSUIT DG
HERBICIDE AND PRESTIGE¹ HERBICIDE FOR ENHANCED
GRASS CONTROL IN SOYBEANS

OBSERVE ALL PRECAUTIONARY AND WORKER PROTECTION STATEMENTS IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

In addition to broadleaf control, PURSUIT DG provides good to excellent control of several important grasses in soybeans. The addition of PRESTIGE herbicide to PURSUIT DG enhances the grass control, especially when heavy infestations of grass exist. It also provides control of grasses not controlled by PURSUIT DG. In some cases the activity of PRESTIGE may be reduced when mixed with PURSUIT DG. The reduction in activity may be overcome by delaying the application of PRESTIGE herbicide 7 days following the application of PURSUIT DG. If PRESTIGE is applied first, wait 3 days before applying PURSUIT DG.

For optimum control, apply the tank mixture to actively growing weeds at the sizes indicated in this Supplemental Labeling (or for sequential applications refer to application rates and weeds sizes indicated in the PURSUIT DG and PRESTIGE labels).

Refer to the PRESTIGE label for additional information regarding application rates, restrictions, precautions, weeds controlled, adjuvants recommended and other information. **When tank-mixing herbicides, always follow the most restrictive label.**

APPLICATION RATES

Apply PURSUIT DG at the rate of 1.44 ounces per acre. Refer to the table below for the appropriate rate of PRESTIGE herbicide for enhanced grass control.

ACCEPTED
with COMMENTS
In EPA Letter Dated
5/1/81

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241-350

PRESTIGE Rate * (ounces per acre)	Annual Grasses Controlled	Size (inches)
12 oz.	Wild Proso Millet Shattercane	4-10" 3-12"
16 oz.	Foxtail, Giant Jung'erie Panicum, Fall Texas Signalgrass, Broadleaf	3-8" 3-8" 3-8" 3-8" 3-8"
20 oz.	Volunteer Corn	4-10"
24 oz.	Barnyardgrass Crabgrass, Large Smooth Cupgrass, woolly Foxtail, Green Yellow Goosegrass Johnsongrass, Seedling Sprangletop, Red Witchgrass	3-8" 3-6" 3-6" 3-8" 3-8" 3-8" 3-6" 3-8" 3-8" 3-8"

*If a mixture of grasses are present, use the highest rate indicated for the grasses present

MIXING INSTRUCTIONS

POSTEMERGENCE APPLICATIONS OF PURSUIT DG AND PRESTIGE REQUIRE THE ADDITION OF A CROPOIL CONCENTRATE AND A LIQUID FERTILIZER SOLUTION

- I. CROP OIL CONCENTRATE:** A petroleum or vegetable seed based oil concentrate (such as SUN-IT II²) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other methylated seed oils at the rate of 1 to 1.5 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre

AND

II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs. per acre

This label must be in the possession of the user at the time of herbicide application

© 1994 by Dow Agrosciences
 All rights reserved.
 Printed in the United States of America

Manufactured by Dow Agrosciences
 Agricultural Products Division
 Crop Protection Chemicals Department
 Wayne, NJ 07093-1199

EPA Reg. No. 241-350

FOR USE IN SOYBEANS

**TANK-MIXTURE OR SEQUENTIAL APPLICATIONS OF PURSUIT DG
HERBICIDE AND STATUS¹ HERBICIDE FOR ENHANCED
WEED CONTROL IN SOYBEANS**

OBSERVE ALL PRECAUTIONARY AND WORKER PROTECTION STATEMENTS IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. SEE THE PURSUIT DG LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

STATUS herbicide may be tankmixed with PURSUIT DG for enhanced control of several broadleaf weeds. Additionally, STATUS may be applied sequentially, following a PURSUIT DG application. STATUS herbicide provides fast burn-down of susceptible broadleaf weeds. Temporary bronzing of the soybean leaves soon after application may occur. Soybeans soon outgrow early symptoms; yield will not be effected by adding STATUS to PURSUIT DG. In some cases the activity of PURSUIT DG may be reduced when mixed with STATUS. The reduction in activity may be overcome by delaying the application of STATUS herbicide 7 days following the application of PURSUIT DG.

WEEDS CONTROLLED

The addition of STATUS to PURSUIT DG at the recommended rates will enhance the control of several broadleaf weeds, including common and giant ragweed, pigweed species and waterhemp. (Refer to the STATUS label for additional weeds controlled).

APPLICATION RATES

PURSUIT DG:

Apply PURSUIT DG at the broadcast rate of 1.44 ounces per acre (1 soluble bag treats 5 acres). Refer to the following tables for the recommended rate of STATUS herbicide.

ACCEPTED
~~FOR COMMENTS~~
In EPA Letter Dated

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241-350

When tank-mixing STATUS with PURSUIT DG, apply STATUS at the following rates, depending on weed size:

	STATUS Rate (ounces per acre):*		
	8 - 10 oz.	12 - 14 oz.	16 - 20 oz.
Weeds/Weed Size	1 - 4"	4 - 6"	6 - 8"
Common ragweed Pigweed species, Waterhemp, tall common			
Giant Ragweed		1 - 6"	6 - 8"***

*Use the higher rate if common ragweed is present or the weed population is high.

**Use the 24 ounce/acre rate if giant ragweed is 6 - 8 inches tall.

STATUS SEQUENTIAL APPLICATION RATES

When applying STATUS following a PURSUIT DG application (sequential), apply STATUS at the following rates:

	STATUS Rate (ounces per acre):*		
	10 - 12 oz.	14 - 16 oz.	18 - 24 oz.
Weeds/Weed Size	1 - 4"	4 - 6"	6 - 8"
Common ragweed Pigweed species, Waterhemp, tall common			
Giant Ragweed		1 - 6"	6 - 8"***

*Use the higher rate if common ragweed is present or the weed population is high.

**Use the 24 ounce/acre rate if giant ragweed is 6 - 8 inches tall.

MIXING INSTRUCTIONS

POSTEMERGENCE APPLICATIONS OF PURSUIT DG AND STATUS REQUIRE THE ADDITION OF AN ADJUVANT AND A LIQUID FERTILIZER SOLUTION.

ADJUVANTS

- I. **SURFACTANTS:** Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 quart per 100 gallons.

OR

CROP OIL CONCENTRATE: Instead of a surfactant, a petroleum or vegetable seed based oil concentrate (such as SUN-IT II²) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other methylated seed oils at the rate of 1.5 to 2 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre. A non-ionic surfactant is recommended when crop response is of concern (i.e., under conditions of high relative humidity or the presence of dew on the crop leaves).

AND

II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs. per acre.

This label must be in the possession of the user at the time of herbicide application.

Refer to the STATUS label for additional information regarding application rates, restrictions, precautions, weeds controlled, adjuvants recommended and other information. **When tank-mixing herbicides, always follow the most restrictive label.**

¹ Trademark of BASF Corporation

² Trademark of Agsco, Inc.

®Registered Trademark of American Cyanamid Company

© 1988 American Cyanamid Company

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470

EPA Reg. No. 241-350

6/01

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, AND SOUTHERN AND ENGLISH PEAS

REFER TO THE MAP ON PAGE 6 FOR GEOGRAPHICAL USE AREA

READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations, however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash[®].

ACCEPTED
with COMMENTS
In EPA Letter Dated

ACCEPTED
In EPA Letter Dated

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

† Data on file with EPA, Registration Division.
* Registered Under Federal Insecticide, Fungicide, and Rodenticide Act.

treatments when PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

USE RATE

0.72, 1.08 OR 1.44 OUNCES PER ACRE

Depending on soil type and edible bean or pea type, apply PURSUIT DG at a broadcast rate according to the following table.

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
0.72	10 acres
1.08	6-2/3 acres
1.44	5 acres

LIMA BEANS

DO NOT apply PURSUIT DG after June 30

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide.

Preemergence Applications Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

NAVY, GREAT NORTHERN, RED KIDNEY, CRANBERRY, BLACK TURTLE, AND SMALL WHITE TYPE DRY BEANS

DO NOT apply to Domino variety black turtle beans

DO NOT apply PURSUIT DG to sand or loamy sand soils

DO NOT apply PURSUIT DG after June 20

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications Apply PURSUIT DG within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide. Refer to the following table for use rates.

wid. \leq 2% O.M.	0.72
All other soils with $>$ 2% O.M.	1.08

Preemergence Applications Apply PURSUIT DG at the rates shown in the preceding table immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ENGLISH PEAS

DO NOT apply PURSUIT DG to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT DG to English peas that have been or will be treated with trifluralin.

Preemergence Applications Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. **DO NOT** apply PURSUIT DG if English peas have been treated with trifluralin.

Postemergence Applications Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. **DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.** If English peas have been treated with trifluralin, PURSUIT DG may be applied **ONLY** at 0.72 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT DG application may increase the likelihood and severity of crop injury.

SOUTHERN PEAS

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications Apply PURSUIT DG at the broadcast rate of 1.44 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide.

Preemergence Applications Apply PURSUIT DG at the broadcast rate of 1.44 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

Early Postemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.44 ounces per acre when peas are at least 3 inches in height. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

WEEDS CONTROLLED

PURSUIT DG applied to low organic matter soils at the broadcast rate of 0.72 ounces per acre preplant incorporated or preemergence will control light to moderate infestations of:

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence, or early postemergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

Postemergence applications of 1.08 ounces per acre must be made to weeds less than 2 inches tall for best results.

When applied as directed at the broadcast rate of 1.44 ounces per acre, PURSUIT DG will control or reduce competition from the weeds listed below:

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence.)

BROADLEAF WEEDS

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Anoda, spurred	C	2	1 - 2
Artichoke, Jerusalem		8	6 - 10
Buffalobur	C**		
Bristly starbur		2	1 - 2
Carpetweed	C		
Cocklebur, common	C**	8	1 - 8
Galinsoga	C		
Jimsonweed	C*	4	1 - 3
Kochia	C	4	1 - 3
Lambsquarters	C*	R	1 - 2
Mallow, Venice	R		
Morningglory			
entireleaf	R	2	1 - 2
ivyleaf	R	2	1 - 2
pitted	R	2	1 - 2
smallflower	C	4	1 - 3
tall	R	2	1 - 2
Mustard sp	C	4	1 - 3
Nightshade,			
black	C	4	1 - 3
Eastern black	C	4	1 - 3
hairy	C	4	1 - 3
Pigweed,			
redroot	C	8	1 - 8
smooth	C	8	1 - 8
spiny	C	8	1 - 8
Poinsettia, wild	C		
Puncturevine	C		
Purshane, common	C		
Pusley, Florida	C		
Sida, prickly	C*		

290

BROADLEAF WEEDS (Continued)

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Ragweed,			
common	R	4	1 - 3
giant	R	4	1 - 3
Sage, barnyard		R	1 - 3
Smartweed,			
ladysthumb	C	4	1 - 3
Pennsylvania	C	4	1 - 3
Spurge,			
prostrate	C	4	1 - 3
spotted	C	4	1 - 3
Sunflower, common	C*	4	1 - 3
Velvetleaf	C*	4	1 - 3
Thistle, Canada		R	1 - 3

*When soil applied, common lambsquarters, jimsonweed, prickly sida, velvetleaf and common sunflower are more consistently controlled by preplant incorporated treatments.

**Use soil applications for light to moderate infestations only. Must be preplant incorporated for best results.

GRASS WEEDS*

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Barnyardgrass	R	3	1 - 3
Crabgrass,			
large	R	3	1 - 3
smooth	R	3	1 - 3
Cupgrass, woolly		3**	1 - 3
Foxtail,			
giant	C	6	1 - 6
green	C	3	1 - 3
robust purple	C	3	1 - 3
robust white	C	3	1 - 3
yellow	C	3	1 - 3
Goosegrass	R		
Johnsongrass,			
seedling	C	6	1 - 8
rhizome		R	1 - 8
Panicum,			
fall	R		
Texas	R		

*When soil applied to grasses, more consistent control can be obtained from preplant incorporated treatments

**PURSUTE DG controls emerged woolly cupgrass only

GRASS WEEDS (Continued)

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Red rice		3	1 - 3
Shattercane	R	6	1 - 8
Signalgrass, broadleaf	R	4	1 - 8

SEDGES

Nutsedge,			
purple	R	R	1 - 3
yellow	R	R	1 - 3

DO NOT count cotyledon leaves when determining weed stage of growth.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - Peanuts
 - IMI-Corn™
 - Edible bean and pea types listed on this label (Refer to heading on page 1)
2. Two months after PURSUIT DG application:
 - Snap beans (DE, MD, PA and VA only)
3. Four months after PURSUIT DG application:
 - Barley
 - Edible bean and pea types not listed on this label (Refer to heading on page 1)
 - Rye
 - Wheat
4. Eight and one-half months after PURSUIT DG application
 - Field corn
 - Field corn grown for seed*
5. Nine and one-half months after PURSUIT DG application
 - Tobacco
6. Eighteen months after PURSUIT DG application

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet Corn
Popcorn	

7. Twenty-six months after PURSUIT DG application:
Potatoes

8. Forty months after PURSUIT DG application:
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

ROTATIONAL CROP GUIDELINE (Continued)

In the event of a crop loss, the bean or pea crop may be replanted.


DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

USE AREA MAP

PURSUIT DG can be applied to the edible legume vegetables listed only in the states or parts of states shaded in the following map:

Use area includes: South Dakota east of U.S. 81, Nebraska east of U.S. 81, Kansas east of U.S. 81, Maryland and Virginia EXCLUDING THE DELMARVA PENINSULA, and other states or portions of states shaded in the following map.



10951
EPA Reg. No. 241-350

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,
CRANBERRY, SMALL WHITE TYPE DRY BEANS, LIMA BEANS,
SOUTHERN AND ENGLISH PEAS, AND WHITE LUPINS**

FOR DISTRIBUTION AND USE IN WISCONSIN ONLY

THIS LABEL EXPIRES APRIL 30, 1997

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended:

1. Two months after PURSUIT DG application:
Snap beans

ACCEPTED
with COMMENTS
In EPA Letter Dated

*Registered Trademark of American Cyanamid Company
EPA 17276 Rev. 1-12/96 SD



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

FOR USE IN LIMA BEANS ONLY

**FOR DISTRIBUTION AND USE ONLY ON THE DELMARVA PENINSULA OF
MARYLAND, DELAWARE, AND VIRGINIA**

READ LABEL CAREFULLY

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application

DO NOT apply by air

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

*Registered Trademark of American Cyanamid Company

indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply PURSUIT DG after June 30.

Allow at least 30 days between application and harvest

Preplant Incorporated Applications Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide

Preemergence Applications Apply PURSUIT DG at the broadcast rate of 0.72 ounces per acre to sand and loamy sand soils, or 1.08 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
0.72	10 acres
1.08	6-2/3 acres

moderate infestations of:

Nightshade, black
Nightshade, Eastern black
Pigweed, redroot

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild
Nightshade, black
Nightshade, Eastern black
Nightshade, hairy
Pigweed, redroot

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime
Soybeans
Peanuts
IMI-Corn™
Lima beans
2. Two months after PURSUIT DG application:
Snap beans
3. Four months after PURSUIT DG application:
Barley
Other types of edible beans and peas
Rye
Wheat
4. Eight and one-half months after PURSUIT DG application
Field corn
Field corn grown for seed*
5. Nine and one-half months after PURSUIT DG application
Tobacco

ROTATIONAL CROP GUIDELINE (continued)

6. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet corn |
| Popcorn | |

7. Twenty-six months after PURSUIT DG application:

- Potatoes

8. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

In the event of a crop loss, the bean or pea crop may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



16/51
EPA Reg. No. 241-350

**FOR USE IN GREAT NORTHERN, RED KIDNEY AND PINTO
TYPE DRY BEANS**

FOR DISTRIBUTION AND USE ONLY IN NEW MEXICO

THIS LABEL EXPIRES APRIL 30, 1997

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

PRECAUTIONS
In EPA Letter Dated

*Registered Trademark of American Cyanamid Company

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
and amendments for the pesticide
registration under EPA Reg. No.
241-350

Use only nonionic surfactants as a spray additive for post applications

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after June 30. Allow at least 60 days between application and harvest

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. **DO NOT** apply PURSUIT DG on sand or loamy sand soils

Postemergence Applications Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

(oz/acre)

1.08

Treats

6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preemergence or early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label

- 1 Anytime
 - Soybeans
 - Peanuts
 - Edible bean types listed on this label (Refer to heading on page 1)
 - IMI-Corn™
- 2 Four months after PURSUIT DG application
 - Barley
 - Edible bean types not listed on this label (Refer to heading on page 1)
 - Peas
 - Rye
 - Wheat
- 3 Eight and one-half months after PURSUIT DG application
 - Field corn
 - Field corn grown for seed*
- 4 Nine and one-half months after PURSUIT DG application
 - Tobacco

¹Dash is a trademark of BASF Corporation

™Trademark American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

5. Eighteen months after PURSUIT DG application

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet corn |
| Popcorn | |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss Great Northern, red kidney or pinto beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



EPA Reg. No. 241-350

1975

**FOR USE IN RED KIDNEY TYPE DRY BEANS, WHITE LUPINS,
LIMA BEANS, AND ENGLISH PEAS ONLY**

FOR DISTRIBUTION AND USE ONLY IN ILLINOIS, IOWA, AND WISCONSIN

READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations, however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

ACCEPTED
~~with conditions~~
In EPA Letter Dated

*Registered Trademark of American Cyanamid Company

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

[Signature]

DO NOT make more than one application of PURSUIT DG per year

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures

RED KIDNEY TYPE DRY BEANS, WHITE LUPINS AND LIMA BEANS

DO NOT apply PURSUIT DG after June 30

DO NOT apply PURSUIT DG to white lupins grown on sand or loamy sand soils

Allow at least 30 days between application and harvest of lima beans

Allow at least 60 days between application and harvest of red kidney beans or white lupins

Preplant Incorporated Applications Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide

Preemergence Applications Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide

ENGLISH PEAS

Allow at least 30 days between application and harvest

Preplant Incorporated Applications Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT DG to English peas that have been or will be treated with trifluralin

ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT. For optimum weed control and crop safety, apply PURSUIT DG prior to 5 nodes before flowering. If English peas have been treated with trifluralin, PURSUIT DG may be applied ONLY at 0.72 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT DG application may increase the likelihood and severity of crop injury.

ECO-PAK CONVERSION TABLE

<u>Use Rate (oz/acre)</u>	<u>1 ECO-PAK Soluble Bag Treats</u>
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence, or early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label

- 1. Anytime
 - Soybeans
 - Peanuts
 - Edible bean and pea types listed on this label (Refer to heading on page 11)
 - IMI-Corn™

† Trademark of Dow Chemical Company

™ Trademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

2. Four months after PURSUIT DG application.

Barley
Edible bean and pea types not listed on this label (Refer to heading on page 1).
Rye
Wheat

3. Eight and one-half months after PURSUIT DG application.

Field corn
Field corn grown for seed*

4. Nine and one-half months after PURSUIT DG application.

Tobacco

5. Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	

6. Twenty-six months after PURSUIT DG application

Potatoes

7. Forty months after PURSUIT DG application.

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, red kidney beans, white lupins, lima beans, or English peas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



EPA Reg. No. 241-350

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
241-350

FOR USE IN ENGLISH PEAS ONLY

**FOR DISTRIBUTION AND USE ONLY ON THE DELMARVA PENINSULA OF
MARYLAND, DELAWARE, AND VIRGINIA**

READ LABEL CAREFULLY

THIS LABEL EXPIRES APRIL 30, 1997

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

DISCLAIMER

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. DO NOT apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application

DO NOT apply by air.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations, however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products

DO NOT make more than one application of PURSUIT DG per year.

*Registered Trademark of American Cyanamid Company

DG cannot be mixed with any product containing a label prohibiting such mixtures

DO NOT apply PURSUIT DG to sand or loamy sand soils

Allow at least 30 days between application and harvest

Preplant Incorporated Applications Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT DG to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. **DO NOT** apply PURSUIT DG if English peas have been treated with trifluralin.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label

- Anytime
 - Soybeans
 - Peanuts
 - IMI-CornTM
 - English peas and lima beans
- Two months after PURSUIT DG application
 - Snap beans
- Four months after PURSUIT DG application
 - Barley
 - Other types of edible beans and peas
 - Rye
 - Wheat

6. Eighteen months after PURSUIT DG application

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	

7. Twenty-six months after PURSUIT DG application

- Potatoes

8. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

- * Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

In the event of a crop loss, the bean or pea crop may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

TMTrademark of American Cyanamid Company
EPL 1282 Rev. 1-1-2279-500



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996

EPA Reg. No. 241-350

**FOR PREEMERGENCE USE IN SNAP BEANS GROWN
IN NEW JERSEY ONLY**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to snap beans. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG applications must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT apply PURSUIT DG after June 20.

Allow at least 30 days between application and harvest

DO NOT make more than one application of PURSUIT DG per year.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

ACCEPTED
with ~~the~~ label
In EPA Letter Dated

*Registered Trademark of American Cyanamid Company

(see reverse side)

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

Preemergence Applications For suppression of pigweed, nightshade and common purslane, apply PURSUIT DG at the broadcast rate of 0.54 ounces per acre (one ECO-PAK soluble bag treats 13-1/3 acres) immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- 1 Anytime:
 - Soybeans
 - Peanuts
 - Edible beans and peas
 - IMI-Corn™
- 2 Four months after PURSUIT DG application:
 - Barley
 - Rye
 - Wheat
- 3 Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
- 4 Nine and one-half months after PURSUIT DG application:
 - Tobacco
- 5 Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	
- 6 Twenty-six months after PURSUIT DG application:
 - Potatoes

field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, snap beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

™ Trademark of American Cyanamid Company
© 1996, Rev. 1-1-96, SD



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 © 1996

EPA Reg. No. 241-350

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, this pesticide
registered under EPA Reg. No.

**FOR USE IN SUCCULENT PEAS, DRY EDIBLE PEAS,
LENTILS, CHICKPEAS, AND LIMA BEANS**

PURSUIT DG can be used on succulent peas, dry edible peas (including chickpeas) and lentils in Washington (Adams, Asotin, Columbia, Garfield, Spokane, Walla Walla and Whitman counties), Idaho (Benewah, Clearwater, Idaho, Kootenai, Latah, Lewis, and Nez Perce counties), Oregon (Union, Umatilla and Wallowa counties) and Montana only.

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans. Plant peas, lentils, or lima beans at least 1/2 inch deep to reduce risk of crop injury.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT DG per year.

Allow at least 30 days between application and harvest of succulent peas and lima beans.

Allow at least 60 days between application and harvest of dry edible peas, chickpeas, and lentils.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION INSTRUCTIONS

Preplant Incorporated Applications. Apply PURSUIT DG within 1 week before planting. DO NOT incorporate deeper than 3 inches. Refer to the PURSUIT DG leaflet label for incorporation instructions.

Preemergence Applications. Apply PURSUIT DG after planting, but prior to crop emergence.

PURSUIT DG may be tank-mixed with Sencor DF¹ or Lexone DF² to assist in the control of lambsquarters or mayweed-chamomile (dogfennel). Refer to the Sencor or Lexone label for proper application rates and restrictions.

WEEDS CONTROLLED

PURSUIT DG applied preplant incorporated or preemergence will control:

Buckwheat, wild	Nightshade, Eastern black
Kochia	Nightshade, hairy
Lambsquarters, common*	Pigweed, redroot
Mustard, wild	Shepherdspurse
Nightshade, black	Thistle, Russian

*preplant incorporated applications only

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label

1. Anytime

Chickpeas	Peanuts
Lentils	Soybeans
Peas	IMI-Corn TM
Lima beans	
2. Four months after PURSUIT DG application
 - Barley
 - Dry edible beans
 - Rye
 - Snap beans (3 months)
 - Wheat
3. Nine and one-half months after PURSUIT DG application
 - Field corn
 - Field corn grown for seed**
 - Tobacco

¹Sencor is a trademark of the Parent Company of Eubank, Inc. or Bayer GmbH, Leverkusen.

²Lexone is a trademark of E. I. duPont de Nemours and Company, Inc.

TMTrademark of American Cyanamid Company.

FD-1228 Rev. 2-1-11-76 SD

6. Forty months after PURSUIT DG application.

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, peas, lentils, chickpeas, or lima beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

** Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470

© 1976