

PM 25 241-350 3-11-97 10/25

Please read Instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 11-30-93

(A)		United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number 207927
		Application for Pesticide:		

Section I

1. Company/Product Number 241-350	2. EPA Product Manager P. Errico	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) PURSUIT® DG herbicide	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) American Cyanamid Company Agricultural Research Center P.O. Box 400 Princeton, NJ 08543-0400 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain below <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<table style="width:100%;"> <tr> <td style="width:50%; border: 1px solid black;"> <input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - explain below. </td> <td style="width:50%; text-align: center; vertical-align: middle;"> NOTIFICATION MAR 11 1997 </td> </tr> </table>	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - explain below.	NOTIFICATION MAR 11 1997
<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - explain below.	NOTIFICATION MAR 11 1997		

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
 Notification of changes in packaging label statements and label expiration date, and incorporation of approved wording per PR Notice 95-2. This notification is consistent with the provisions of PR Notice 95-2 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 95-2 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No * Certification must be submitted.	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted.	If "Yes," Unit Package wgt. No. per container	If "Yes," Package wgt. No. per container	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) of Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner In Which Label Is Affixed To Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other (_____) <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Janet M. Overholt	Title Product Registrations Manager	Telephone No. (Include Area Code) 609-716-2410

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (S.amped)
2. Signature 	3. Title Product Registrations Manager	[Stamp Area]
4. Typed Name Janet M. Overholt	5. Date 2/26/97	



2 of 25

THIS VOLUME DOES NOT CONTAIN 40 CFR 158 DATA

American Cyanamid Company
Agricultural Products Research Division
P.O. Box 400
Princeton, NJ 08543-0400
Phone (609) 716-2000
General Fax (609) 275-3582

February 26, 1997

Dr. Philip V. Errico
Product Manager (25)
Registration Division (H-7505C)
Office of Pesticide Programs
U.S. Environmental Protection Agency
Crystal Mall 2, Room 266A
1921 Jefferson Davis Highway
Arlington, VA 22202

Attention: Edward J. Allen

RE: PURSUIT® DG Herbicide (EPA Reg. No. 241-350):
Notification of Label Changes for Four Supplemental Labels
for Use in Edible Legume Vegetables
OPP Id No. 207927

Dear Dr. Errico:

American Cyanamid Company hereby notifies EPA of minor changes to four supplemental labels for PURSUIT DG herbicide (EPA Reg. No. 241-350) use in edible legume vegetables. The labels include the following uses:

- ◆ Wisconsin in navy, great northern, red kidney, black turtle, cranberry, small white type dry beans, lima beans, southern and English Peas, and white lupins
- ◆ North Dakota and Northern Minnesota in navy, great northern, red kidney, black turtle, cranberry, and pinto type dry beans, and dry edible peas
- ◆ English peas on the Delmarva Peninsula
- ◆ Michigan in navy, great northern, red kidney, black turtle, cranberry, and small white type dry beans.

The changes to the labels include the extension or removal of the self-imposed expiration date, removal of reference to acres treated per package size, and revision of Rotational Crop Guidelines to agree with the stamp-approved full label.



Philip V. Errico
PURSUIT® DG Herbicide

February 26, 1997
Page 2

In support of this notification, we are submitting an application form, data certification form, and revised labels. Copies of the stamp-approved supplemental and full labels are also included. Five additional copies of each of the revised labels are also enclosed.

Thank you for processing this notification. If there are any questions concerning this submission, please contact me at 609-716-2410.

Respectfully submitted,

A handwritten signature in cursive script that reads 'Janet M. Overholt'.

Janet M. Overholt
Product Registrations Manager
U.S. Plant Regulatory Affairs

encl.

® Registered Trademark of American Cyanamid Company

**Revisions to Michigan Label for Use in
Navy, Great Northern, Red Kidney, Black Turtle, Cranberry, and
Small White Type Dry Beans**

- Changed the self-imposed expiration date to April 30, 1999.
- Removed ECO-PAK® CONVERSION TABLE section, because ECO-PAK water-soluble bags can now be marketed in more than one size.
- Revised Rotational Crop Guidelines for alfalfa and barley to agree with the stamp-approved full label.

® Registered Trademark of American Cyanamid Company.



Supplemental Labeling

EPA Reg. No. 241-350

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,
CRANBERRY, AND SMALL WHITE TYPE DRY BEANS ONLY**

**FOR DISTRIBUTION AND USE IN MICHIGAN ONLY
(EXCLUDING UPPER PENINSULA)**

THIS LABEL EXPIRES APRIL 30, 1997

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply at any timing to Domino variety black turtle beans.

DO NOT apply PURSUIT DG to sand or loamy sand soils.

DO NOT apply PURSUIT DG after July 10.

Allow at least 60 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

Soil Type	Use Rate (oz/acre)
Sands, loamy sands	DO NOT USE
Sandy loams and all other soils with ≤ 2% O.M.	0.72
All other soils with > 2% O.M.	1.08

Preemergence Applications. Apply PURSUIT DG at the rates shown in the preceding table immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

Postemergence Applications. Apply PURSUIT DG at 0.72 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Basagran¹ may be tank-mixed with PURSUIT DG to control weeds not listed on the PURSUIT DG label. Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use only nonionic surfactants as a spray additive.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
0.72	10 acres
1.08	6-2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied to low organic matter soils at the broadcast rate of 0.72 ounces per acre preplant incorporated, preemergence, or postemergence will control light to moderate infestations of:

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- I. Anytime:
 - Soybeans
 - Peanuts
 - IMI-CORN™
 - Edible bean and pea types listed on this label (Refer to heading on page 1)

¹Dash and Basagran are trademarks of BASF Corporation
™ Trademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

2. Four months after PURSUIT DG application:

Alfalfa Barley

Edible bean and pea types not listed on this label (Refer to heading on page 1)

Rye

Wheat

3. Eight and one-half months after PURSUIT DG application:

Field corn

Field corn grown for seed*

4. Nine and one-half months after PURSUIT DG application:

Barley

Tobacco

5. Eighteen months after PURSUIT DG application:

Alfalfa

Safflower

Cotton

Sorghum

Lettuce

Sunflower

Oats

Sweet corn

Popcorn

6. Twenty-six months after PURSUIT DG application:

Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



CYANAMID

American Cyanamid Company
North America Agricultural Products Division
Crop Protection Products Department
One Campus Drive, Parsippany, NJ 07054 ©1997

Revisions to Label for English Peas on the Delmarva Peninsula

- Changed the self-imposed expiration date to April 30, 1999.
- Removed ECO-PAK® CONVERSION TABLE section, because ECO-PAK water-soluble bags can now be marketed in more than one size.
- Revised Rotational Crop Guidelines for alfalfa and barley to agree with the stamp-approved full label.

® Registered Trademark of American Cyanamid Company.



Supplemental Labeling

EPA Reg. No. 241-350

FOR USE IN ENGLISH PEAS ONLY

FOR DISTRIBUTION AND USE ONLY ON THE DELMARVA PENINSULA OF
MARYLAND, DELAWARE, AND VIRGINIA

READ LABEL CAREFULLY

THIS LABEL EXPIRES APRIL 30, 1997

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply PURSUIT DG to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. DO NOT apply PURSUIT DG to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. DO NOT apply PURSUIT DG if English peas have been treated with trifluralin.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6 2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Soybeans
 - Peanuts
 - IMI-CORN™
 - English peas and lima beans
- Two months after PURSUIT DG application:
 - Snap beans
- Four months after PURSUIT DG application:
 - Alfalfa Barley
 - Other types of edible beans and peas
 - Rye
 - Wheat

- Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
- Nine and one-half months after PURSUIT DG application:
 - Barley
 - Tobacco
- Eighteen months after PURSUIT DG application:
 - Alfalfa
 - Cotton
 - Lettuce
 - Oats
 - Popcorn
 - Safflower
 - Sorghum
 - Sunflower
 - Sweet corn
- Twenty-six months after PURSUIT DG application:
 - Potatoes
- Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

In the event of a crop loss, the bean or pea crop may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



American Cyanamid Company
North America Agricultural Products Division
Crop Protection Products Department
One Campus Drive, Parsippany, NJ 07054 ©1997

11/01/25

**Revisions to North Dakota and Northern Minnesota Label for Use in
Navy, Great Northern, Red Kidney, Black Turtle, Cranberry, and
Pinto Type Dry Beans, and Dry Edible Peas**

- Changed the self-imposed expiration date to April 30, 1999.
- Removed reference to acres treated per ECO-PAK® water-soluble bag, because the bags can now be marketed in more than one size.
- Revised Rotational Crop Guidelines to agree with the stamp-approved full label.

® Registered Trademark of American Cyanamid Company.



Supplemental Labeling

EPA Reg. No. 241-350

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY,
BLACK TURTLE, CRANBERRY, AND PINTO TYPE DRY BEANS,
AND DRY EDIBLE PEAS ONLY**

**FOR DISTRIBUTION AND USE IN NORTH DAKOTA AND
NORTHERN MINNESOTA (see USE AREA below)**

THIS LABEL EXPIRES APRIL 30, 1999⁷

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall) PURSUIT DG may cause injury to subsequent planted crops. Sugar beets are particularly sensitive to PURSUIT DG residues in soil. PURSUIT DG is not recommended in crop rotations that include sugar beets. See the **ROTATIONAL CROP** section of this supplemental label for rotations to other sensitive crops.

USE AREA

PURSUIT DG may be used in North Dakota and in Minnesota north of highway #210.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply at any timing to Domino variety black turtle beans.

Pinto varieties UI-111 and Olathe are more sensitive to PURSUIT DG than other varieties.

DO NOT apply PURSUIT DG after July 10.

DO NOT feed treated forage, hay, or straw to livestock.

Allow at least 60 days between application and harvest.

Postemergence Applications. Apply PURSUIT DG at 0.72 ounces per acre (one ECO-PAK soluble bag treats 10 acres). A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 0.72 ounces per acre will control:

- wild mustard
- black nightshade*
- eastern black nightshade*
- *suppression

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - IMI-CORN™
 - Edible bean and pea types listed on this label (Refer to heading on page 1)
2. Four Months after PURSUIT DG application
 - Alfalfa
 - Edible bean and pea types not listed on this label
 - Wheat
- ~~3.2~~ Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
- ~~4.3~~ Nine and one-half months after PURSUIT DG application:
 - Barley
 - Wheat
 - Edible bean and pea types not listed on this label (Refer to heading on page 1)
- ~~5.4~~ Eighteen months after PURSUIT DG application:

Barley	Rye
Cotton	Safflower
Lettuce	Sorghum
Oats	Sunflower
Popcorn	Sweet Corn

¹Dash is a trademark of BASF Corporation
™Trademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

6.5. Twenty-six months after PURSUIT DG application:

- Potatoes
- Flax

7.6. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

15 of 25

**Revisions to Wisconsin Label for Use in
Navy, Great Northern, Red Kidney, Black Turtle, Cranberry, Small White Type
Dry Beans, Lima Beans, Southern and English Peas, and White Lupins**

- Removed the self-imposed expiration date.



Supplemental Labeling

EPA Reg. No. 241-350

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, SMALL WHITE TYPE DRY BEANS, LIMA BEANS, SOUTHERN AND ENGLISH PEAS, AND WHITE LUPINS

FOR DISTRIBUTION AND USE IN WISCONSIN ONLY

~~THIS LABEL EXPIRES APRIL 30, 1997~~

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended:

- Two months after PURSUIT DG application:
Snap beans

*Registered Trademark of American Cyanamid Company
PE-17276 Rev. 2-1 2/6/97 4/23/96 SD



American Cyanamid Company
North America Agricultural Products Division
Crop Protection Products Department
One Campus Drive, Parsippany, NJ 07054 ©1997



Supplemental Labeling

EPA Reg. No. 241-350

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,
CRANBERRY, AND SMALL WHITE TYPE DRY BEANS ONLY**

**FOR DISTRIBUTION AND USE IN MICHIGAN ONLY
(EXCLUDING UPPER PENINSULA)**

THIS LABEL EXPIRES APRIL 30, 1999

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply at any timing to Domino variety black turtle beans.

DO NOT apply PURSUIT DG to sand or loamy sand soils.

DO NOT apply PURSUIT DG after July 10.

Allow at least 60 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

Soil Type	Use Rate (oz/acre)
Sands, loamy sands	DO NOT USE
Sandy loams and all other soils with ≤ 2% O.M.	0.72
All other soils with > 2% O.M.	1.08

Preemergence Applications. Apply PURSUIT DG at the rates shown in the preceding table immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

Postemergence Applications. Apply PURSUIT DG at 0.72 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Basagran¹ may be tank-mixed with PURSUIT DG to control weeds not listed on the PURSUIT DG label. Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use only nonionic surfactants as a spray additive.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

WEEDS CONTROLLED

PURSUIT DG applied to low organic matter soils at the broadcast rate of 0.72 ounces per acre preplant incorporated, preemergence, or postemergence will control light to moderate infestations of:

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - Peanuts
 - IMI-CORN™
 - Edible bean and pea types listed on this label (Refer to heading on page 1)

¹Dash and Basagran are trademarks of BASF Corporation
™Trademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

- 2. Four months after PURSUIT DG application:
 - Alfalfa
 - Edible bean and pea types not listed on this label (Refer to heading on page 1)
 - Rye
 - Wheat

- 3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*

- 4. Nine and one-half months after PURSUIT DG application:
 - Barley
 - Tobacco

- 5. Eighteen months after PURSUIT DG application:

Cotton	Safflower
Lettuce	Sorghum
Oats	Sunflower
Popcorn	Sweet corn

- 6. Twenty-six months after PURSUIT DG application:
 - Potatoes


- 7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.





Supplemental Labeling

EPA Reg. No. 241-350

FOR USE IN ENGLISH PEAS ONLY

**FOR DISTRIBUTION AND USE ONLY ON THE DELMARVA PENINSULA OF
MARYLAND, DELAWARE, AND VIRGINIA**

READ LABEL CAREFULLY

THIS LABEL EXPIRES APRIL 30, 1999

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT DG per year.

2/af-25

Refer to the PURSUIT DG leaflet label for approved combination treatments. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply PURSUIT DG to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. DO NOT apply PURSUIT DG to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. DO NOT apply PURSUIT DG if English peas have been treated with trifluralin.

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated or preemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - Peanuts
 - IMI-CORN™
 - English peas and lima beans
2. Two months after PURSUIT DG application:
 - Snap beans
3. Four months after PURSUIT DG application:
 - Alfalfa
 - Other types of edible beans and peas
 - Rye
 - Wheat

4. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
5. Nine and one-half months after PURSUIT DG application:
 - Barley
 - Tobacco
6. Eighteen months after PURSUIT DG application:

Cotton	Safflower
Lettuce	Sorghum
Oats	Sunflower
Popcorn	Sweet corn
7. Twenty-six months after PURSUIT DG application:
 - Potatoes
8. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.


* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

In the event of a crop loss, the bean or pea crop may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

™Trademark of American Cyanamid Company
PE-17282 Rev. 2 2/6/97 SD





Supplemental Labeling

22/95

EPA Reg. No. 241-350

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY,
BLACK TURTLE, CRANBERRY, AND PINTO TYPE DRY BEANS,
AND DRY EDIBLE PEAS ONLY**

**FOR DISTRIBUTION AND USE IN NORTH DAKOTA AND
NORTHERN MINNESOTA (see USE AREA below)**

THIS LABEL EXPIRES APRIL 30, 1999

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. DO NOT apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall) PURSUIT DG may cause injury to subsequent planted crops. Sugar beets are particularly sensitive to PURSUIT DG residues in soil. PURSUIT DG is not recommended in crop rotations that include sugar beets. See the ROTATIONAL CROP section of this supplemental label for rotations to other sensitive crops.

USE AREA

PURSUIT DG may be used in North Dakota and in Minnesota north of highway #210.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply at any timing to Domino variety black turtle beans.

Pinto varieties UI-111 and Olathe are more sensitive to PURSUIT DG than other varieties.

DO NOT apply PURSUIT DG after July 10.

DO NOT feed treated forage, hay, or straw to livestock.

Allow at least 60 days between application and harvest.

Postemergence Applications. Apply PURSUIT DG at 0.72 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 0.72 ounces per acre will control:

- wild mustard
- black nightshade*
- eastern black nightshade*
- *suppression

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - IMI-CORN™
 - Edible bean and pea types listed on this label (Refer to heading on page 1)
2. Four Months after PURSUIT DG application
 - Alfalfa
 - Edible bean and pea types not listed on this label
 - Wheat
3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
4. Nine and one-half months after PURSUIT DG application:
 - Barley
5. Eighteen months after PURSUIT DG application:

Cotton	Rye
Lettuce	Safflower
Oats	Sorghum
Popcorn	Sunflower
	Sweet Corn

¹Dash is a trademark of BASF Corporation
™Trademark of American Cyanamid Company

24 of 25

ROTATIONAL CROP GUIDELINE (continued)

6.. Twenty-six months after PURSUIT DG application:

Potatoes
Flax

7.. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



Supplemental Labeling

EPA Reg. No. 241-350

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,
CRANBERRY, SMALL WHITE TYPE DRY BEANS, LIMA BEANS,
SOUTHERN AND ENGLISH PEAS, AND WHITE LUPINS**

FOR DISTRIBUTION AND USE IN WISCONSIN ONLY

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

GENERAL INFORMATION

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended:

1. Two months after PURSUIT DG application:
Snap beans



American Cyanamid Company
North America Agricultural Products Division
Crop Protection Products Department
One Campus Drive, Parsippany, NJ 07054

PM 25 241-350 3-11-97 10/24

Please read Instructions on reverse before completing form.

Form Approved OMB No. 2070-0060. Approval expires 11-30-93

(A) 	United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number 207929
	Application for Pesticide:		

Section I

1. Company/Product Number 241-350	2. EPA Product Manager P. Errico	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) PURSUIT® DG herbicide	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) American Cyanamid Company Agricultural Research Center P.O. Box 400 Princeton, NJ 08543-0400 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain below <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<div style="text-align: right; font-weight: bold; font-size: 1.2em;">NOTIFICATION</div> Final printed labels in response to Agency letter dated MAR 11 1997 <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - explain below.
---	---

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of change to expiration date, incorporation of approved wording, addition of other states and a specific legume name per PR Notice 95-2. This notification is consistent with the provisions of PR Notice 95-2 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 95-2 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted.		If "Yes," Unit Package wgt.	No. per container	If "Yes," Package wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label Is Affixed To Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other (_____)		

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Janet M. Overholt	Title Product Registrations Manager	Telephone No. (Include Area Code) 609-716-2410
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Product Registrations Manager	
4. Typed Name Janet M. Overholt	5. Date 2/26/97	

2/24



THIS VOLUME DOES NOT CONTAIN 40 CFR 158 DATA

American Cyanamid Company

Agricultural Products Research Division
P O Box 400
Princeton NJ 08543 0400
Phone (609) 716 2000
General Fax (609) 275 3582

February 26, 1997

Dr Philip V Errico
Product Manager (25)
Registration Division (H-7505C)
Office of Pesticide Programs
U S Environmental Protection Agency
Crystal Mall 2, Room 266A
1921 Jefferson Davis Highway
Arlington, VA 22202

Attention Edward J Allen

RE: PURSUIT® DG Herbicide (EPA Reg. No 241-350):
Notification of Label Changes for Three Supplemental Labels
for Use in Edible Legume Vegetables
OPP Id No 207929

Dear Dr Errico

American Cyanamid Company hereby notifies EPA of minor changes to three supplemental labels for PURSUIT DG herbicide (EPA Reg No 241-350) use in edible legume vegetables. The labels include the following uses

- ◆ New Mexico in great northern, red kidney and pinto type dry beans
- ◆ Minnesota and South Dakota in navy, great northern, red kidney, black turtle, cranberry, pinto, and small white type dry beans, lima beans, dry edible peas, English peas, lentils, and chickpeas
- ◆ Postemergence use in snap beans in West Texas and New Mexico

Revisions common to all three labels include removal or changing of the expiration date and changes in the Rotational Crop Guidelines to agree with the full label. Other changes to individual labels are the addition of another specific legume name, addition of states to the supplemental label use area and deletion and addition of advisory statements concerning liquid fertilizers

3 of 24




Philip V. Errico
PURSUIT® DG Herbicide

February 26, 1997
Page 2

In support of this notification, we are submitting an application form, data certification form, and revised labels. Copies of the stamp-approved supplemental and full labels are also enclosed. Five copies of each revised label are included.

Thank you for processing this notification. If you have any questions concerning this submission, please contact me at 609-716-2410.

Respectfully submitted,

Janet M. Overholt
Product Registrations Manager
U.S. Plant Regulatory Affairs

encl.
© Registered Trademark of American Cyanamid Company.

**Revisions to New Mexico Label for Use in
Great Northern, Red Kidney and Pinto Type Dry Beans**

- Deleted self-imposed expiration date.
- Deleted advisory statement concerning liquid fertilizers. Added advisory phrase that liquid fertilizers may increase crop injury.
- Removed reference to acres treated per ECO-PAK® water-soluble bag, because the bags can now be marketed in more than one size.
- Revised Rotational Crop Guidelines for alfalfa and barley to agree with the stamp-approved full label.

® Registered Trademark of American Cyanamid Company.



Supplemental Labeling

5/24

EPA Reg. No. 241-350

FOR USE IN GREAT NORTHERN, RED KIDNEY AND PINTO TYPE DRY BEANS

FOR DISTRIBUTION AND USE ONLY IN NEW MEXICO

~~THIS LABEL EXPIRES APRIL 30, 1997~~

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

*Registered Trademark of American Cyanamid Company

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use ~~only~~ a nonionic surfactants as a spray additive for post applications. Liquid fertilizers may increase crop injury.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT DG per year. DO NOT apply PURSUIT DG after June 30. Allow at least 60 days between application and harvest.

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. DO NOT apply PURSUIT DG on sand or loamy sand soils.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6 2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preemergence or early postemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Soybeans
 - Peanuts
 - Edible bean types listed on this label (Refer to heading on page 1).
 - IMI-CORN™
- Four months after PURSUIT DG application:
 - Barley Alfalfa
 - Edible bean types not listed on this label (Refer to heading on page 1).
 - Peas
 - Rye
 - Wheat
- Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
- Nine and one-half months after PURSUIT DG application:
 - Barley
 - Tobacco

¹Dash is a trademark of BASF Corporation
™Trademark American Cyanamid Company

7/24

ROTATIONAL CROP GUIDELINE (continued)

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet corn |
| Popcorn | |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss Great Northern, red kidney or pinto beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

**Revisions to Minnesota and South Dakota Label for Use in
Navy, Great Northern, Red Kidney, Black Turtle, Cranberry, Pinto, and Small
White Type Dry Beans, Lima Beans, Dry Edible Peas, English Peas, Lentils, and
Chickpeas**

- Added White Lupins to the list of specific legume vegetables (succulent or dried) for this use area.
- Deleted self-imposed expiration date.
- Removed reference to acres treated per ECO-PAK® water-soluble bag, because the bags can now be marketed in more than one size.
- Revised Rotational Crop Guidelines for alfalfa and barley to agree with the stamp-approved full label. Also, clarified which crops can be planted anytime after application.



Supplemental Labeling

7 of 24

EPA Reg. No. 241-350

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, PINTO, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, DRY EDIBLE PEAS, ENGLISH PEAS, LENTILS, WHITE LUPINS AND CHICKPEAS

FOR DISTRIBUTION AND USE IN MINNESOTA (south of highway #210) AND SOUTH DAKOTA

~~THIS LABEL EXPIRES APRIL 30, 1997~~

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

GENERAL INFORMATION

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

Some crop varieties may be inherently less tolerant to PURSUIT DG applications and crop injury could result. Contact the seed company for information and recommendations about the use of PURSUIT DG postemergence.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

*Registered Trademark of American Cyanamid Company

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT apply to Domino variety black turtle beans.

DO NOT make more than one application of PURSUIT DG per year.

Allow at least 30 days between application and harvest of lima beans and English peas. Allow at least 60 days between application and harvest of dry beans, dry edible peas, lentils and chickpeas.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION INSTRUCTIONS

DO NOT apply PURSUIT DG to sand or loamy sand soils.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre within 1 week before planting. **DO NOT** incorporate deeper than 3 inches. Refer to the PURSUIT DG leaflet label for incorporation instructions.

DO NOT apply preplant incorporated to pinto beans.

Preemergence Applications: Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre after planting, but prior to crop emergence. If weeds are present at the time of application, a burndown herbicide such as Roundup² or Gramoxone Extra³ should be tank-mixed with PURSUIT DG to control emerged weeds.

DO NOT apply preemergence to pinto beans.

Postemergence Applications. FOR USE IN ENGLISH PEAS, NAVY, PINTO, GREAT NORTHERN AND RED KIDNEY TYPE DRY BEANS ONLY. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. **DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.**

ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6 2/3 acres

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence or postemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
 - Chickpeas
 - Lentils
 - Dry edible peas
 - Peanuts
 - Edible beans and peas (dry and lima listed on this label)
 - Soybeans
 - IMI-CORN™
- Four months after PURSUIT DG application:
 - Barley Alfalfa
 - Edible beans (dry, snap) and peas not listed on this label
 - Rye
 - Wheat
- Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*

¹Dash is a trademark of BASF Corporation

²Roundup is a trademark of Monsanto Agricultural Products Company

³Gramoxone Extra is a trademark of Zeneca, Inc.

™Trademark American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

4. Nine, and one-half months after PURSUIT DG application:

- Barley
- Tobacco

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Alfalfa | Safflower |
| Cotton | Sorghum |
| Lettuce | Sunflower |
| Oats | Sweet corn |
| Popcorn | |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, edible beans, dry edible peas, lentils or chickpeas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

12/94

**Revisions to Label for Postemergence Use in Snap Beans
in West Texas and New Mexico**

- Added Arkansas, Missouri, and Oklahoma to the use area for this label.
- Changed the self-imposed expiration date to April 30, 1999.
- Removed reference to acres treated per ECO-PAK® water-soluble bag, because the bags can now be marketed in more than one size.
- Revised Rotational Crop Guidelines for alfalfa and barley to agree with the stamp-approved full label.

® Registered Trademark of American Cyanamid Company.



Supplemental Labeling

EPA Reg. No. 241-350

**FOR POSTEMERGENCE USE IN SNAP BEANS GROWN
IN ARKANSAS, MISSOURI, OKLAHOMA, WEST TEXAS AND NEW
MEXICO ONLY**

(see USE AREA below)

**FOR USE ONLY BY GROWERS CONTRACTED BY ALLEN CANNING
COMPANY. THIS LABEL EXPIRES APRIL 30, 1997**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION
INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING.
APPLY WITH GROUND EQUIPMENT ONLY.**

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS. SPECIFICALLY, CYANAMID DISCLAIMS ANY LIABILITY FOR CLAIMS, CAUSES OF ACTION, FINES, PENALTIES, DAMAGES, INCLUDING CONSEQUENTIAL DAMAGES, LOSSES, LIABILITIES, JUDGMENTS, AND EXPENSES ARISING OUT OF OR RELATING TO ANY CROP LOSS OR OTHER INJURY TO PERSONS OR PROPERTY RESULTING FROM THE USE OF PURSUIT DG HERBICIDE ON EDIBLE LEGUMES.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

USE AREA

For use in the Texas Counties of Bailey, Castro, Lamb, and Parmer, and the New Mexico Counties of Curry and Roosevelt and the states of Arkansas, Missouri and Oklahoma.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to snap beans. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

14 of 24

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG applications must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash¹.

Allow at least 30 days between application and harvest.

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after June 30.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION INSTRUCTIONS

Postemergence Applications. For suppression of pigweed and nightshade, apply PURSUIT DG at the broadcast rate of 0.54 ounces per acre (~~one ECO-PAK soluble bag treats 13 1/3 acres~~) in a tank-mix combination with Basagran¹. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use only nonionic surfactants as a spray additive.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

15 of 24

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:

Soybeans	Edible beans and peas
Peanuts	IMI™-CORN
2. Four months after PURSUIT DG application:

Alfalfa	Barley	Rye
Wheat		
3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
4. Nine and one-half months after PURSUIT DG application:
 - Barley
 - Tobacco
5. Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	
6. Twenty-six months after PURSUIT DG application:
 - Potatoes
7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, snap beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

©PURSUIT DG is a registered trademark of American Cyanamid Company

™IMI-CORN is a Trademark of American Cyanamid Company

‡Dash and Basagran are trademarks of BASF Corporation





Supplemental Labeling

EPA Reg. No. 241-350

**FOR POSTEMERGENCE USE IN SNAP BEANS GROWN
IN ARKANSAS, MISSOURI, OKLAHOMA, WEST TEXAS AND NEW
MEXICO ONLY**

(see USE AREA below)

**FOR USE ONLY BY GROWERS CONTRACTED BY ALLEN CANNING
COMPANY. THIS LABEL EXPIRES APRIL 30, 1999.**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION
INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING.
APPLY WITH GROUND EQUIPMENT ONLY.**

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS. SPECIFICALLY, CYANAMID DISCLAIMS ANY LIABILITY FOR CLAIMS, CAUSES OF ACTION, FINES, PENALTIES, DAMAGES, INCLUDING CONSEQUENTIAL DAMAGES, LOSSES, LIABILITIES, JUDGMENTS, AND EXPENSES ARISING OUT OF OR RELATING TO ANY CROP LOSS OR OTHER INJURY TO PERSONS OR PROPERTY RESULTING FROM THE USE OF PURSUIT DG HERBICIDE ON EDIBLE LEGUMES.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

USE AREA

For use in the Texas Counties of Bailey, Castro, Lamb, and Parmer, and the New Mexico Counties of Curry and Roosevelt and the states of Arkansas, Missouri and Oklahoma.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to snap beans. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG applications must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash¹.

Allow at least 30 days between application and harvest.

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after June 30.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION INSTRUCTIONS

Postemergence Applications. For suppression of pigweed and nightshade, apply PURSUIT DG at the broadcast rate of 0.54 ounces per acre in a tank-mix combination with Basagran¹. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use only nonionic surfactants as a spray additive.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

180424

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- 1. Anytime:
 - Soybeans Edible beans and peas
 - Peanuts IMI™-CORN
- 2. Four months after PURSUIT DG application:
 - Alfalfa Rye
 - Wheat
- 3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
- 4. Nine and one-half months after PURSUIT DG application:
 - Barley
 - Tobacco
- 5. Eighteen months after PURSUIT DG application:
 - Cotton Safflower
 - Lettuce Sorghum
 - Oats Sunflower
 - Popcorn Sweet corn

- 6. Twenty-six months after PURSUIT DG application:
 - Potatoes

- 7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, snap beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

®PURSUIT DG is a registered trademark of American Cyanamid Company

™IMI-CORN is a Trademark of American Cyanamid Company

¹Dash and Basagran are trademarks of BASF Corporation



Supplemental Labeling

EPA Reg. No. 241-350

FOR USE IN GREAT NORTHERN, RED KIDNEY AND PINTO TYPE DRY BEANS

FOR DISTRIBUTION AND USE ONLY IN NEW MEXICO

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

20 of 24

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use a nonionic surfactant as a spray additive for post applications. Liquid fertilizers may increase crop injury.

DO NOT use crop oils, petroleum oils, or Dash.¹

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after June 30. Allow at least 60 days between application and harvest.

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. **DO NOT** apply PURSUIT DG on sand or loamy sand soils.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preemergence or early postemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - Peanuts
 - Edible bean types listed on this label (Refer to heading on page 1).
 - IMI-CORN™
2. Four months after PURSUIT DG application:
 - Alfalfa
 - Edible bean types not listed on this label (Refer to heading on page 1).
 - Peas
 - Rye
 - Wheat
3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*
4. Nine and one-half months after PURSUIT DG application:
 - Barley
 - Tobacco

¹Dash is a trademark of BASF Corporation
™Trademark American Cyanamid Company

2/16/97

ROTATIONAL CROP GUIDELINE (continued)

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Cotton | Safflower |
| Lettuce | Sorghum |
| Oats | Sunflower |
| Popcorn | Sweet corn |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss Great Northern, red kidney or pinto beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.



Supplemental Labeling

EPA Reg. No. 241-350

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY,
PINTO, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, DRY EDIBLE PEAS,
ENGLISH PEAS, LENTILS, WHITE LUPINS AND CHICKPEAS

FOR DISTRIBUTION AND USE IN MINNESOTA (south of highway #210)
AND SOUTH DAKOTA

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

GENERAL INFORMATION

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

Some crop varieties may be inherently less tolerant to PURSUIT DG applications and crop injury could result. Contact the seed company for information and recommendations about the use of PURSUIT DG postemergence.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

23 of 24

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT apply to Domino variety black turtle beans.

DO NOT make more than one application of PURSUIT DG per year.

Allow at least 30 days between application and harvest of lima beans and English peas. Allow at least 60 days between application and harvest of dry beans, dry edible peas, lentils and chickpeas.

When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION INSTRUCTIONS

DO NOT apply PURSUIT DG to sand or loamy sand soils.

Preplant Incorporated Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre within 1 week before planting. **DO NOT** incorporate deeper than 3 inches. Refer to the PURSUIT DG leaflet label for incorporation instructions.

DO NOT apply preplant incorporated to pinto beans.

Preemergence Applications: Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre after planting, but prior to crop emergence. If weeds are present at the time of application, a burndown herbicide such as Roundup² or Gramoxone Extra³ should be tank-mixed with PURSUIT DG to control emerged weeds.

DO NOT apply preemergence to pinto beans.

Postemergence Applications. FOR USE IN ENGLISH PEAS, NAVY, PINTO, GREAT NORTHERN AND RED KIDNEY TYPE DRY BEANS ONLY. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. **DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.**

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence or postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Peanuts
 - Edible beans and peas listed on this label
 - Soybeans
 - IMI-CORN™
2. Four months after PURSUIT DG application:
 - Alfalfa
 - Edible beans and peas not listed on this label
 - Rye
 - Wheat
3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*

¹Dash is a trademark of BASF Corporation

²Roundup is a trademark of Monsanto Agricultural Products Company

³Gramoxone Extra is a trademark of Zeneca, Inc.

™Trademark American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

4. Nine and one-half months after PURSUIT DG application:

- Barley
- Tobacco

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Cotton | Safflower |
| Lettuce | Sorghum |
| Oats | Sunflower |
| Popcorn | Sweet corn |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, edible beans, dry edible peas, lentils or chickpeas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.