



Page 1 of 3

Supplemental Labeling

EPA Reg. No. 241-350

FOR USE IN GREAT NORTHERN, PINTO AND RED KIDNEY TYPE DRY BEANS

FOR DISTRIBUTION AND USE ONLY IN COLORADO, KANSAS, NEBRASKA, WYOMING, TEXAS AND WISCONSIN

THIS LABEL EXPIRES APRIL 30, 1999

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT DG HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT DG ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT DG TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT DG LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

GENERAL INFORMATION

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT DG in any manner not specifically described in this label.

ACCEPTED with COMMENTS
In EPA Letter Dated

APR 14 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

*Registered Trademark of American Cyanamid Company

241-350

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Uniformly apply PURSUIT DG using 10 or more gallons of water per acre with flat fan nozzle tips only, at a spray pressure of 20 to 40 psi.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications. **DO NOT** use crop oils, petroleum oils, or Dash.¹

Liquid Fertilizer can only be added when PURSUIT and Basagran¹ are tank-mixed. See postemergence tankmix instructions.

DO NOT make more than one application of PURSUIT DG per year. **DO NOT** apply PURSUIT DG after July 15. Allow at least 60 days between application and harvest.

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

PURSUIT DG should be used as part of a total weed management program involving the use of a registered soil-applied herbicide as a tank-mix or sequential partner.

Preemergence Applications. Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 1 day after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. **DO NOT** apply PURSUIT DG on sand or loamy sand soils.

Postemergence Applications. Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Postemergence Tankmix Combination. The addition of at least 1 pint of Basagran¹ may be tank-mixed with PURSUIT DG to control weeds not listed on the PURSUIT DG label. Refer to the Basagran label for proper application rates and

restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use a nonionic surfactant as a spray additive. Liquid fertilizer can be included with the nonionic surfactant with this tank mix only.

DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preemergence or early postemergence will control:

- | | |
|---------------------------|-------------------|
| Mustard, wild | Nightshade, hairy |
| Nightshade, black | Pigweed, redroot |
| Nightshade, Eastern black | |

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

1. Anytime:
 - Soybeans
 - Peanuts
 - Edible bean types listed on this label (Refer to heading on page 1).
 - IMI-CORNTM
2. Four months after PURSUIT DG application:
 - Alfalfa
 - Edible bean types not listed on this label (Refer to heading on page 1).
 - Peas
 - Rye
 - Wheat
3. Eight and one-half months after PURSUIT DG application:
 - Field corn
 - Field corn grown for seed*

¹Dash and Basagran are trademarks of BASF Corporation
TMTrademark of American Cyanamid Company

ROTATIONAL CROP GUIDELINE (continued)

4. Nine and one-half months after PURSUIT DG application:

- Barley
- Tobacco

5. Eighteen months after PURSUIT DG application:

- | | |
|---------|------------|
| Cotton | Safflower |
| Lettuce | Sorghum |
| Oats | Sunflower |
| Popcorn | Sweet Corn |

6. Twenty-six months after PURSUIT DG application:

- Potatoes

7. Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.