


PM 25

241-350

9-25-78

1074



SEP 25 1998

EPA Reg. No. 241-350

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, PURSUIT DG herbicide is registered under EPA Reg. No. 241-350

**PURSUIT DG HERBICIDE FOR ROSEAN RED KIDNEY TYPE DRY BEANS, WHITE LUPINS, LIMA BEANS, AND ENGLISH PEAS ONLY**

**FOR DISTRIBUTION AND USE ONLY IN ILLINOIS, IOWA, AND WISCONSIN**

**READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT DG LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

**DISCLAIMER**

PURSUIT DG application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT DG only to the legume vegetables specifically listed in this labeling. **DO NOT** apply PURSUIT DG in geographical regions outside those specifically defined in this labeling for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT DG application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT DG if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT DG if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT DG postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT DG ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

THIS PRODUCT WHEN USED ON EDIBLE LEGUME VEGETABLE CROPS MAY LEAD TO CROP INJURY, LOSS, OR DAMAGE. CYANAMID RECOMMENDS THAT THE USER AND/OR GROWER TEST THIS PRODUCT IN ORDER TO DETERMINE ITS SUITABILITY FOR SUCH INTENDED USE. CYANAMID MAKES THIS PRODUCT AVAILABLE TO THE USER AND/OR GROWER SOLELY TO THE EXTENT THE BENEFIT AND UTILITY, IN THE SOLE OPINION OF THE USER AND/OR GROWER, OUTWEIGH THE EXTENT OF POTENTIAL INJURY ASSOCIATED WITH THE USE OF THIS PRODUCT. THE DECISION TO USE OR NOT TO USE THIS HERBICIDE MUST BE MADE BY EACH INDIVIDUAL PURSUIT DG USER AND/OR GROWER ON THE BASIS OF POSSIBLE CROP INJURY FROM PURSUIT DG, THE SEVERITY OF WEED INFESTATION, THE COST OF ALTERNATIVE WEED CONTROLS, AND OTHER FACTORS. CYANAMID INTENDS THAT BECAUSE OF THE RISK OF FAILURE TO PERFORM OR CROP DAMAGE THAT ALL SUCH USE IS AT THE USER'S AND/OR GROWER'S RISK.

AMERICAN CYANAMID DISCLAIMS ANY LIABILITY FOR CLAIMS, CAUSES OF ACTION, FINES, PENALTIES, DAMAGES, INCLUDING CONSEQUENTIAL INCIDENTS AND DAMAGES, LOSSES, LIABILITIES, JUDGMENTS, AND EXPENSES ARISING OUT OF OR RELATING TO INJURY TO PERSONS, CROPS, OR PROPERTY RESULTING FROM THE USE OF PURSUIT DG HERBICIDE ON EDIBLE LEGUMES CONTRARY TO THE LABEL INSTRUCTIONS. AMERICAN CYANAMID COMPANY WARRANTS ONLY THAT

**DISCLAIMER (continued)**

**THE MATERIAL CONTAINED HEREIN CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL AND IS REASONABLY FIT FOR THE USE THEREIN DESCRIBED WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE, SUBJECT TO THE RISKS REFERRED TO ABOVE.**

**CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.**

**BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF PURSUIT DG. In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This labeling must be in the possession of the user at the time of herbicide application.

**DO NOT** apply by air.

**DO NOT** apply PURSUIT DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT DG.

Spray equipment used for PURSUIT DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

**DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.<sup>1</sup>

**DO NOT** make more than one application of PURSUIT DG per year.

Refer to the PURSUIT DG leaflet label for approved combinations. When PURSUIT DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive labeling restrictions and precautions. No label dosages should be exceeded. PURSUIT DG cannot be mixed with any product containing a label prohibiting such mixtures.

**RED KIDNEY TYPE DRY BEANS,  
WHITE LUPINS AND LIMA BEANS**

**DO NOT** apply PURSUIT DG after June 30.

**DO NOT** apply PURSUIT DG to white lupins grown on sand or loamy sand soils.

Allow at least 30 days between application and harvest of lima beans.

Allow at least 60 days between application and harvest of red kidney beans or white lupins.

**Preplant Incorporated Applications.** Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT DG may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after, or up to 3 days after planting. PURSUIT DG may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

**ENGLISH PEAS**

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT DG to English peas that have been or will be treated with trifluralin.

**Preemergence Applications.** Apply PURSUIT DG at the broadcast rate of 1.08 ounces per acre immediately after,

or up to 3 days after planting. **DO NOT** apply PURSUIT DG if English peas have been treated with trifluralin.

**Postemergence Applications.** Apply PURSUIT DG at 1.08 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**DO NOT APPLY PURSUIT DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.** For optimum weed control and crop safety, apply PURSUIT DG prior to 5 nodes before flowering. If English peas have been treated with trifluralin, PURSUIT DG may be applied ONLY at 0.72 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT DG application may increase the likelihood and severity of crop injury.

### ECO-PAK CONVERSION TABLE

Use Rate (oz/acre)	1 ECO-PAK Soluble Bag Treats
1.08	6-2/3 acres

### WEEDS CONTROLLED

PURSUIT DG applied at the broadcast rate of 1.08 ounces per acre preplant incorporated, preemergence, or early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 1.08 ounces per acre must be to weeds less than 2 inches tall for best results.

### ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT DG as recommended in this label:

- Anytime:
  - Soybeans
  - Peanuts
  - Edible bean and pea types listed on this label (Refer to heading on page 1).
  - IMI-CORN<sup>®</sup> seed hybrids

- Four months after PURSUIT DG application:
  - Barley
  - Edible bean and pea types not listed on this label (Refer to heading on page 1).
  - Rye
  - Wheat
- Eight and one-half months after PURSUIT DG application:
  - Field corn
  - Field corn grown for seed\*
- Nine and one-half months after PURSUIT DG application:
  - Tobacco
- Eighteen months after PURSUIT DG application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	
- Twenty-six months after PURSUIT DG application:
  - Potatoes
- Forty months after PURSUIT DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, red kidney beans, white lupins, lima beans, or English peas may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production:

4094

American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT DG shall be assumed by the user.

**CONSULT THE PURSUIT DG LEAFLET LABEL FOR OTHER PRECAUTIONS.**



**American Cyanamid Company**

North America Agricultural Products Division  
Crop Protection Products Department  
One Campus Drive, Parsippany, NJ 07054-0140

<sup>1</sup>Dash is a trademark of BASF Corporation

<sup>®</sup> Registered Trademarks of American Cyanamid Company

PE-17277 Rev. 2 09 08 58

