

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (75-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.
241-354

DATE OF ISSUANCE
~~1995~~

TERM OF ISSUANCE
Conditional

FEB 24 1995

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME OF PESTICIDE PRODUCT

AVENELSM SC with out herbicide

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

American Cyanamid Company
Agricultural Research Division
P.O. BOX 400
PRINCETON, NJ 08543-0400

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Section 3(a)(7)(C) of the Act requires that the registrant

1. Add the phrase "EPA Registration No. 241-354" to the label
2. Submit five copies of the label to the Registration Division

and provided that American Cyanamid Company, the manufacturer of the active ingredient of this product, submits the required data to continue the registration of Avenel SC with out herbicide under 18756 in the registration evaluation document for this active ingredient.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(c). Your request for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Enclosure

Joanne I. Miller
Product Manager (23)
Fungicide-herbicide Branch
Registration Division (75000)

BEST AVAILABLE COPY

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

Joanne I. Miller FOR JOANNE MILLER

FEB 24 1995

ACCEPTED
with COMMENTS
in EPA Letter Dated

2 of 11

AVENGE® SG
Wild Oat Herbicide

FEB 24 1995

FOR USE IN BARLEY AND WHEAT

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, this pesticide
registered under EPA Reg. No.

241-354

ACTIVE INGREDIENT:	
difenzoquat methyl sulfate	
1,2-dimethyl-3,5-diphenyl-1H-pyrazolium methyl sulfate.....	92.5%
INERT INGREDIENTS.....	7.5%
TOTAL	100.0%

(1 pound contains 0.64 lb. of difenzoquat cation)

EPA Reg. No. 241-

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

DANGER!/PELIGRO!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.
- IF SWALLOWED: Call a physician or poison control center. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.
- IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

See Inside For Additional Precautionary Statements.
See Directions For Use Inside

In case of an emergency endangering life or property involving this product, call collect, day or night, Area Code 201-835-3100.

Net Contents: (Company Logo)
American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1994

®Registered Trademark of American Cyanamid Company

BEST AVAILABLE COPY

3 7 11

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER!

**CORROSIVE: CAUSES IRREVERSIBLE EYE DAMAGE
MAY BE FATAL IF SWALLOWED OR INHALED**

DO NOT get in eyes, on skin, or on clothing. May be fatal if inhaled. **DO NOT** breathe dust, vapor or spray mist.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear:

- . long-sleeved shirt and long pants
- . waterproof gloves
- . shoes plus socks
- . protective eyewear
- . For exposures outdoors, dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C).

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- o Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- o Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- o Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

BEST AVAILABLE COPY

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. DO NOT apply when weather conditions favor run-off and/or drift from target area. DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DO NOT apply this product through any type of irrigation system.

This label must be in the possession of the user at the time of pesticide application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protection equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- . coveralls
- . waterproof gloves
- . shoes plus socks
- . protective eyewear.

STORAGE AND DISPOSAL

Stability of AVENGE SG is not affected by exposure to cold temperature.

DO NOT contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. DO NOT reuse empty container.

BEST AVAILABLE COPY

7/94

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL FOR A BAG ATTACHED IN A BOX: Completely empty contents of bag into application equipment. Then dispose of container in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR REMOVABLE BAG IN A BOX: Completely empty contents of bag into application equipment. Then dispose of bag in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Cardboard boxes may be offered for recycling if allowed by state and local authorities.

CONTAINER DISPOSAL FOR PLASTIC: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR BULK: Return empty container for reuse.

GENERAL INFORMATION

AVENGE SG is a selective herbicide for the postemergence control of wild oats in barley and wheat. See the section DIRECTIONS FOR USE for varietal restrictions.

If broadleaf weeds must also be controlled in these crops, AVENGE SG may be tank mixed with certain broadleaf herbicides which are listed under the HOW TO USE AVENGE SG PLUS BROADLEAF HERBICIDES section. AVENGE SG alone or in combination can be applied postemergence by ground equipment or aircraft.

Maximum control of wild oats in barley and wheat with AVENGE SG alone, or in combination, is obtained when temperature, moisture, fertility and cultural practices provide favorable conditions for plant growth. When applied under poor conditions for plant growth such as cold/wet or hot/dry weather or low soil fertility, AVENGE SG can produce yellowing or slight tip burn of barley or wheat. However, recovery is rapid when good growth conditions return. A slight reduction of yield may result under conditions described above.

Observe all cautions and limitations on this label and the labels of products used in tank mixtures with AVENGE SG. The use of AVENGE SG not consistent with this label can result in injury to crops, animals or persons. Keep containers closed to avoid spills and contamination.

BEST AVAILABLE COPY

WHERE TO USE AVENGE SG: AVENGE SG alone or in combination can be applied to:

BARLEY: All Varieties

WHEAT: SEE TABLE BELOW

WHEAT TYPE	YES	NO
Fall and Winter Seeded	All EXCEPT	Borah, WS 1809, WS 1877, WS 1859, Klasic, Probrand 771, Tadinia, and Mexican durums
Soft-White Spring Seeded	All	
Spring Seeded durums	All EXCEPT	Edmore, Fjord, Kyle, Laker, Lakota, Lloyd, Regal, Renville Vic, Wakooma, Wascona, Westbred 803, and Westbred 881
Hard Red Spring Seeded	Amidon, Apex, Baart, Benito, Bronze Chief, Buckshot, Butte, Butte 86, Canuck, Centa, Chester, Columbus, Copper, Coteau, Courtney, Era, Erik, Federation, Fjeld, Fortuna, Germain 444, Glenlea, Glennman, HY320, Katepwa, Kitt, Kodiak, Leader, Leif, McKay, Marberg, Marshall, Minnpro, Newana, Neepawa, NK(PB) 751, Norak, Norana, Olaf, Oslo, Pioneer 2369, Pioneer 2385, Pondera, Probrand 711, Probrand 715, Prodax, Rambo, Selkirk, Solar, Stoa, Success, Telemark, Vance, Victory, W 2502, Walera, Wheaton and Yecoro rojo	DO NOT apply to varieties of hard red spring wheat not listed

WHEN TO USE AVENGE SG: AVENGE SG or AVENGE SG/broadleaf tank mix combinations are to be applied when the majority of wild oat plants are in the 3- to 5-true leaf stage of growth (tillering). The timing of AVENGE SG application to wild oats frequently coincides with barley in the 2- to 7-leaf stage, spring-seeded wheat in the 5- to 6-leaf stage, and fall-seeded wheat in the 4-leaf to tillered stage of growth. To achieve maximum control, the growth stage of wild oats in each field to be treated should be examined frequently to determine the 3- to 5-true leaf stage (tillering) for proper timing of AVENGE SG application. Density of wild oats per square foot should also be determined to select the correct AVENGE SG application rate.

BEST AVAILABLE COPY

7 8 11

WILD OAT STAGES

(artwork)

(artwork)

(artwork)

3-true leaf stage

4-true leaf stage

5-true leaf stage

EQUIPMENT APPLICATION INSTRUCTIONS

The spray equipment must be clean and properly calibrated before treatments are applied. For best results, use flat fan nozzles, at pressures of 20 to 40 psi to achieve uniform spray distribution and minimize drift. DO NOT use flood jet nozzles. Keep the by-pass line on or near the bottom of the tank to minimize foaming. In ground applications, check nozzles for even distribution of spray droplets. In aerial applications, it is recommended that a flagman be located at each end of the field, or an automatic mechanical flagging unit be attached to the aircraft to ensure uniform aerial application and to avoid overlapping and possible crop injury.

AVENGE SG APPLICATION

DO NOT apply AVENGE SG when the flagleaf is exposed or crop injury may occur. DO NOT apply AVENGE SG if rain is predicted within 6 hours. DO NOT spray AVENGE SG when freezing temperatures are forecast or when plants are wet with dew or rain. DO NOT apply AVENGE SG or AVENGE SG tank mix combinations under severe conditions for plant growth such as drought, extreme temperature fluctuation, low soil fertility and plant disease.

Apply AVENGE SG alone and AVENGE SG plus broadleaf herbicide combinations in 5 to 20 gallons of water per acre by ground equipment or 3 to 10 gallons of water per acre by aircraft. DO NOT let spray drift onto adjacent crops.

HOW TO USE AVENGE SG ALONE

If wild oats are growing vigorously, select the proper application rate from the AVENGE SG Rate Recommendation Table (below) that corresponds to the population of wild oat plants per square foot in the field to be treated.

**ASSERT®/AVENGE SG TANK MIX OR
ASSERT® SG/AVENGE SG TANK MIX FOR WILD OAT CONTROL**

AVENGE SG may be tank mixed* with ASSERT or ASSERT SG herbicides for the control of wild oats only. Consult the ASSERT and ASSERT SG labels for use directions, application rates and timing, state restrictions, precautions, and ASSERT or ASSERT SG recropping/rotation limitations.

* DO NOT tank mix AVENGE SG with any product not registered in specific states of intended use.

BEST AVAILABLE COPY

8 7 11

HOW TO USE AVENGE SG PLUS BROADLEAF HERBICIDES

AVENGE SG may be tank mixed* with the following broadleaf herbicides:

2,4-D amine or ester	Express ¹
Ally ¹	Glean ¹
bromoxynil	Harmony ¹
bromoxynil plus MCPA	Harmony Extra ¹
Curtail ²	MCPA amine or ester
Curtail M ²	

Select the proper AVENGE SG application rate as described in the AVENGE SG Rate Recommendation Table (below). In order to assure optimum broadleaf weed control follow label recommendations of your broadleaf herbicide. Be sure that application timings coincide with application timings for AVENGE SG. DO NOT apply AVENGE SG tank mix combinations when the crop flagleaf is exposed or crop injury may occur. DO NOT apply AVENGE SG tank mix combinations under stress conditions.

Follow the most restrictive precautions, directions, and recropping/rotation limitations that appear on combination partner product labels.

* DO NOT tank mix AVENGE SG with any product not registered in specific states of intended use.

AVENGE SG Rate Recommendation Table

Wild Oat Population	Wild Oat Density Plants Per Square Foot	AVENGE SG Rate Per Acre (ounces)
Low	1-10	15 5/8 (15.625)
Moderate	11-25	18 3/4 (18.75)
High	More than 25	25

GENERAL MIXING INSTRUCTIONS

A NON-IONIC SURFACTANT CONTAINING AT LEAST 80% NON-IONIC SURFACTANT MUST BE USED WITH AVENGE SG HERBICIDE. Use a surfactant registered for application to growing crops. Add 4 pints non-ionic surfactant per 100 gallons (0.5% vol./vol.) of final spray tank volume. Use of crop oil concentrates is not recommended, except as may be described in separately published recommendations from American Cyanamid Company. DO NOT use liquid fertilizer as a substitute for surfactant.

9 of 11

VIGOROUS AGITATION OF THE ENTIRE TANK INCLUDING THE TANK BOTTOM IS REQUIRED TO RAPIDLY DISSOLVE THE WATER SOLUBLE GRANULES OF AVENGE SG HERBICIDE. Poor agitation and dead spots in the tank will retard rapid and thorough dissolving of AVENGE SG herbicide.

PREMIXING INSTRUCTIONS (if tank agitation is inadequate)

Premix the desired amount of AVENGE SG in clean water in a clean polyethylene plastic bucket. Use polyethylene plastic or stainless steel stirring rod or water pressure to mix until the granules of AVENGE SG dissolve. DO NOT allow premix solution to contact skin or clothing. Add premix solution into the spray tank as described in Mixing Instructions. Thoroughly rinse all of the AVENGE SG into the spray tank.

MIXING INSTRUCTIONS

HOW TO MIX AVENGE SG ALONE

1. Fill tank three-quarters full with clean, filtered water. Start vigorous agitation that thoroughly mixes the entire tank, including the tank bottom.
2. Add AVENGE SG to the partially filled tank while agitating.
3. Continue filling the tank with clean water while maintaining agitation.
4. After the AVENGE SG is dissolved and the tank is full, add the non-ionic surfactant to the tank at a rate of 4 pints surfactant per 100 gallons (0.5% vol./vol.) of final spray solution (See section above for brands requiring greater percentage surfactant).
5. Allow mixture to agitate and recycle for at least five minutes before spraying.
6. Agitation must be continued until spraying is complete.

HOW TO MIX AVENGE SG PLUS BROADLEAF HERBICIDES

1. Fill tank three-quarters full with clean, filtered water. Start vigorous agitation that thoroughly mixes the entire tank, including the tank bottom.
2. Add the broadleaf herbicide first to the partially filled tank while agitating. Add the broadleaf herbicide(s) before adding AVENGE SG when using the ASSERT/AVENGE SG or ASSERT SG/AVENGE SG tank mix. NOTE: TO PREVENT MIXING PROBLEMS, TANK MIX PARTNERS MUST BE THOROUGHLY MIXED BEFORE ADDING AVENGE SG.
3. When the broadleaf herbicide is thoroughly mixed, add the AVENGE SG to the partially filled tank while continuing agitation.
4. Continue filling the tank with clean water while maintaining agitation.
5. After the AVENGE SG is dissolved and thoroughly mixed with the tank mix partners, and the tank is full, add the non-ionic surfactant at a rate of 4 pints per 100 gallons (0.5% vol./vol.) final spray solution (See section above for brands requiring greater percentage surfactant).
6. Allow mixture to agitate and recycle for at least five minutes before spraying.
7. Agitation must be continued until spraying is completed.

HOW TO TANK MIX AVENGE SG PLUS ASSERT OR ASSERT SG HERBICIDES

Follow mixing instructions on the ASSERT or ASSEPT SG labels.

BEST AVAILABLE COPY

10 7 11

NOTE: DO NOT MAKE MORE THAN ONE APPLICATION OF AVENGE SG PER GROWING SEASON. DO NOT GRAZE TREATED FIELDS OR CUT TREATED FORAGE FOR SILAGE OR HAY. BARLEY OR WHEAT CAN BE REPLANTED IF LOSS OF CROP OCCURS. AFTER HARVEST, BARLEY AND WHEAT GRAIN AND STRAW FROM TREATED FIELDS CAN BE FED TO LIVESTOCK. ROTATION TO OTHER CROPS CAN BE MADE THE FOLLOWING YEAR, EXCEPT WHERE LIMITED BY TANK MIX PARTNER LABEL RESTRICTION.

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, herbicide resistant weed populations, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. **CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.**

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF AVENGE SG. In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

BEST AVAILABLE COPY

11 7 11

USES WITH OTHER PRODUCTS (TANK MIXES)

If this product is used in combination with any other product except as specifically recommended in writing by American Cyanamid Company, then American Cyanamid Company shall have no liability for any loss, damage, or injury arising out of its use in any such combination not so specifically recommended. If used in a combination recommended by American Cyanamid Company, the liability of American Cyanamid Company shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the American Cyanamid Company product in such combination use, and in any event shall be limited to return of the amount of the purchase price of the product.

For additional information regarding the use of AVENGE SG wild oat herbicide, call telephone number (800) 942-0500.

- 1 Ally, Express, Glean, Harmony, and Harmony Extra are trademarks of E.I. duPont de Nemours and Company, Inc.
- 2 Curtail and Curtail M are trademarks of DowElanco.

BEST AVAILABLE COPY