

# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Jeffery H. Birk, Ph.D. BASF 26 Davis Drive Research Triangle Park, NC 27709

MAR 6 2008

Dear Dr. Birk:

Subject: Label Notification(s) for Pesticide Registration Notice 2007-4

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated, February 7, 2008 for:

### EPA Registration 241-364 Cadre Herbicide

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identify the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself

If you have any questions, please contact me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P)

Office of Pesticide Programs



The Chemical Company

February 7, 2008

U.S. Environmental Protection Agency Office of Pesticide Programs (7505P) Document Processing Desk 7504P (NOTIF) Room S-4900 One Potomac Yard 2777 S. Crystal Drive Arlington, VA 22202 Attention: Mr. James Tompkins, (PM 25)

RE: Notification complying with PRN 2007-4:

Revised Cadre herbicide container disposal language

EPA Reg. No. 241-364

Dear Mr. Tompkins:

BASF is hereby submitting revised labeling for Cadre herbicide (EPA Reg. No. 241-364) to comply with the container disposal changes required by PRN 2007-4. No other substantive changes have been made to the labeling.

### Enclosed please find:

- Application form 8570-1
- CD containing electronic copy of the label
- Certification with Respect to Label Integrity
- · Cadre herbicide label
- Approved Cadre herbicide label

No PRIA fee is required for this notification.

Thank you for your assistance with this matter. If you should have any questions, please feel free to call me at (919) 547-2622.

Regards,

Jeffrey H. Birk, Ph.D. Regulatory Manager Phone 919-547-2622 Mobile: 919-225-9220 Fax: 919-547-2850

Email: jeffrey.birk@basf.com

Registered Trademark of BASF

3	/	
1	Ί,	<u>ب</u>

Please read instructions o	n reverse before compl	eting form.			Form Apr	roved.	OMB No.	2070-006	O. Approvel expires 2	2-28-9
<b>SEPA</b>	Environmenta	United States al Protecti nington, DC 20	•	icy		<b></b>	Registr Amend Other		OPP Identifier Numb	Def
		Applicati	on for P	esticio	le - Sect	tion	1			
1. Company/Product Num 241-364	ber				roduct Men Tompkins	-		3. Pr	oposed Classification	icted
4. Company/Product (Nam Cadre herbicide	10)			РМ# 25					<b>.</b>	
5. Name and Address of A BASF 26 Davis Drive Research Triangle		code)		(b)(i), m to:	y product i	s simi	lar or ider	ntical in co	FIFRA Section 3(c) imposition and labeli	
Check if to	his is a new address		·	Produc	t Name _				,	
			Secti	on - II				······································		
Amendment - Expl	sponse to Agency lette	r dated	<u>.</u>	<b>V</b>	Final printed Agency lette "Me Too" A Other - Expl	er date oplica	d tion.		NOTIFICATION MAR 6 2008	N
Notification of container dis of PR Notice 95-2 and EP/ product. I understand that consistent with violation of This notification is not subj	A regulations at 40 CFR it is a violation of 18 U.S FIFRA and I may be sub	152.46, and no i.C. Sec. 1001 to ject to enforcen	other change o willfully mal nent action a rk at 919-547	es have b ke any fa nd penali 7-2622 (p	een made to se statemen ies under se hone), 919-5	the lab t to EP clions	eling or the A. I further 12 and 14 c	confidentia understand fFIFRA.	I statement of formula of that if this notification is	of this
		<u> </u>	Section	on - II					- <del></del>	
1. Material This Product V	<del></del>		···							
Child-Resistant Packaging  Yes  ✓ No  * Certification must	Unit Packaging  Yes  No  If "Yes" Unit Packaging wgt	No. per		es lo	No. per container	·	2. Type o	f Container  Metal Plastic Glass Paper Other (S		
be submitted		1			l					
3. Location of Net Content    ✓ Lebel	s Information  Container	4. Size(s) Re	tail Containe 1.0 gall			5. Loc		bel Directio	nying product	
6. Manner in Which Label	is Affixed to Product	Lithog Paper Stenc	reph glued iled		Other					
			Section	n - IV						
1. Contact Point	te items directly below .	for identification	n of individu	uel to be	contacted, i	f nece	ssary, to p	rocess this	application.)	
Name Jeffrey H. Birk			Title Regulator	y Manag	er	· · · · · · · · · · · · · · · · · · ·		Telephone 919-547-2	No. (Include Area Co 2622	ide)
	tements i have made or any knowlinglly false or e law.		all attachm			•		• •	6. Date Application Received (Stamped)	
2. Signatura	/ / Sin	t-	3. Title Regulatory	Manage						
4. Typed Name			5. Date		<del></del>		<del></del>			
Jeffrey H. Birk				Febi	uary 7, 2	2008				

Please read instructions on	reverse before comple	ting form.		Form A	pproved.	OMB No. 2	070-006	O. Approval expires 2-28-9
<b>SEPA</b>	Environmenta	Inited States  I Protection  Ington, DC 204	•	,		Registra Amendn Other		OPP Identifier Number
<u> </u>		Application	n for Pest	ticide - Se	ction I	_ <del></del> , .		
1. Company/Product Numb 241-364			2. E	PA Product Ma	nager	<u> </u>	3. Pro	pposed Classification
4. Company/Product (Name Cadre herbicide	)		PM: 25	-	λ,			None Restricted
5. Name and Address of Ap BASF 26 Davis Drive Research Triangle P		de)	(b)( to: EP	i), my produci A Reg. No.	t is simil	lar or identi	cal in co	FIFRA Section 3(c)(3) mposition and labeling
Grack ii tiii	3 13 4 11511 5401 533			oduct Name				
Notification - Explain  Explanation: Use addition  Notification of container disposed of PR Notice 95-2 and EPA	n below.  nal page(s) if necessar  posal label changes for 0  regulations at 40 CFR 1	y. (For section Cadre herbicide 52.46, and no c	(241-364) in co	Final print Agency le "Me Too" Other - Ex II.) mpliance with Pave been made	Applicate Applicate Applicate Applicate Applicate Applicate Applicate Application Applicat	tion.  ow.  4. This notificaling or the c	cation is confidentia	onsistent with the provisions statement of formula of this that if this notification is not
consistent with violation of F		ect to enforcem	ent action and p k at 919-547-26	penalties under s 322 (phone), 919	sections 1	12 and 14 of I	FIFRA.	
			Section	- [[]				
1. Material This Product W Child-Resistant Packaging Yes No Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt.	No. per container	Water Solution  Yes  No  If "Yes"  Package wg	ole Packaging No. per t contain		2. Type of 0	Container  Metal Plastic Glass Paper Other (S	pecify)
3. Location of Net Contents	Container	4. Size(s) Ret		Oth		ation of Lab	el Directio	ns
			Section	- IV				
1. Contact Point   Complete	e items directly below t	for identificatio	n of individual	to be contacted	d, if nece	ssary, to pro	cess this	application.)
Name Jeffrey H. Birk			Title Regulatory M	lanager			Telaphon 919-547-	B No. (Include Area Code) 2622
•	ements I have made or ny knowlinglly false or a law.		all attachment		-		of the	6. Date Application Received (Stamped)
nature	1 36		3. Title Regulatory Ma	anager				
4. Typed Name Jeffrey H. Birk			5. Date	February 7	, 2008			

5/22

# Certification with Respect to Label Integrity

version: 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

	PROPOS	SED LABEL
EPA Registration #	Date Submitted to EPA	Electronic file name
241-364	2-7-08	000241-00364.20080207.NVA 2008-04-265-0027

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.

	02/07/2008
Signature	Date
Jeffrey H. Birk	
Name (typed)	•
Regulatory Manager	
Title	





FOR USE ONLY IN PEANUTS IN THE STATES OF ALABAMA, ARIZONA, ARKANSAS, FLORIDA, GEORGIA, MISSISSIPPI, NEW MEXICO, NORTH CAROLINA, OKLAHOMA, SOUTH CAROLINA, TEXAS, AND VIRGINIA

#### **Active Ingredient:**

Ammonium salt of imazapic [(±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-	
5-oxo-1 <i>H</i> -imidazol-2-yl]-5-methyl-3-pyridinecarboxylic acid]*	23.6%
Inert Ingredients:	76.4%
Total:	100.0%
*Equivalent to 22.2% (±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-	5-

methyl-3-pyridinecarboxylic acid (1 gallon contains 2.0 pounds of active ingredient as the free acid)

U.S. Patent No. 4,798,619 EPA Reg. No. 241-364

EPA Est. No.

## KEEP OUT OF REACH OF CHILDREN CAUTION!/iPRECAUCION!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See inside booklet for complete First Aid, Precautionary Statements, Directions For Use, and Conditions of Sale and Warranty.

In case of an emergency endangering life or property involving this product, call day or night, 1-800-832-HELP (4357)

#### **Net Contents:**

BASF Corporation, Crop Protection 26 Davis Drive, Research Triangle Park, NC 27709 NOTIFICATION

MAR 6 2008

	FIRST AID	
if Inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.     Call a poison control center or doctor for further treatment advice.	
lf on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutés. Call a poison control center or doctor for treatment advice.	
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center for treatment advice.	
	HOTLINE NUMBER	

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).

#### PRECAUTIONARY STATEMENTS

### **HAZARDS TO HUMANS CAUTION!**

Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

#### Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- · Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations:

Users Should:

- Wash hands before eating, chewing gum, using tobacco or using the toilet.
   Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
   Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### **ENVIRONMENTAL HAZARDS**

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark.

DO NOT contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This labeling must be in the possession of the user at the time of pesticide application.

Observe all cautions and limitations on this label and on the labels of products used in combination with Cadre® herbicide. DO NOT use Cadre other than in accordance with the instructions set forth on this label; this may help prevent crop response. Keep containers closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

DO NOT graze or feed treated peanut hay to livestock.

DO NOT apply more than 0.063 lbs. a.i./A imazapic (4.0 fluid oz./A of Cadre) per application or per use season.

Preharvest Interval: There must be an interval of at least 90 days between an application of Cadre and peanut harvest.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal assistance. It also contains specific instructions and exceptions pertaining to the statements on this laber about personal performance protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls

- chemical-resistant gloves made of any waterproof material
- shoes plus socks

#### STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: KEEP FROM FREEZING, DO NOT store below 20° F.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### CONTAINER DISPOSAL:

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity ≤ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse containers too large to shake (capacity > 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

#### GENERAL INFORMATION

Cadre® herbicide can be applied early postemergence in peanuts. Refer to the specific treatment under the "APPLICATION INSTRUCTIONS" section of the label.

The weed killing activity involves uptake of **Cadre** by weed roots and/or foliage and rapid translocation to the growing points. After **Cadre** application, susceptible weeds may show yellowing and weed growth will stop. Several days may be required before the complete death of susceptible weeds. Adequate soil moisture is important for optimum **Cadre** activity. When adequate soil moisture is present, **Cadre** will have residual activity on susceptible germinating weeds; activity on established weeds will depend on the weed species and the depth of its root system in the soil. If adequate rainfall is not received within 5 days, then irrigation at 0.75 inches/acre will serve to activate the **Cadre** soil residual activity. Also when adequate soil moisture is not present and irrigation is not an option then a timely cultivation made at least 14 days after **Cadre** application may improve general herbicide performance.

A Cadre application may result in some peanut yellowing and/or a reduction in vine growth.

Under adverse conditions (including but not limited to high pH  $\geq$  7.5, low nutrient availability, saline conditions, and/or hard-pans), **Cadre** application may induce an adverse crop response.

Naturally occurring biotypes\* of some of the weeds listed on this label may not be effectively controlled by this and/or other products with either the ALS/AHAS enzyme inhibiting mode of action. Other herbicides with the ALS/AHAS enzyme inhibiting mode of action include the sulfonylureas (e.g., Accent\*) Basis\*, Classic\*, Concer\*, Exceed\*, Permit\*, Pinnacle\* herbicides, etc.), the sulfonamides (e.g., Broadstrike\* herbicide, etc.) and the pyrimidyl benzoates (e.g. Staple\* herbicide, etc.)

naturally occurring ALS/AHAS resistant biotypes are present in a field, **Cadre® herbicide** and/or any other ALS/AHAS enzyme inhibiting mode of action herbicide should be tank-mixed or applied sequentially with an appropriate registered herbicide having a different mode of action to ensure control.

\*A weed biotype is a naturally occurring plant within a given species that has a slightly different, but distinct, genetic makeup from other plants.

Use of **Cadre** is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

Replanting: If replanting is necessary in a field previously treated with **Cadre**, the field may be replanted to peanuts. Rework the soil no deeper than 2 inches. **DO NOT** apply an additional treatment of **Cadre** or **PURSUIT®** herbicide.

#### MANAGING OFF-TARGET MOVEMENT

The following recommendations are general in nature. Refer to the **SPRAYING INSTRUCTIONS** section of this label for the specific application directions for **Cadre**.

Spray Drift: Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Spray drift from applying this product may result in damage to sensitive plants adjacent to the treatment area. Only apply this product when the potential for drift to these and other adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, or non-target crops) is minimal. Do not apply when the following conditions exist that increase the likelihood of spray drift from intended targets: high or gusty winds, high temperatures, low humidity, temperature inversions.

To minimize spray drift, the applicator should be familiar with and take into account the following drift reduction advisory information. Additional information may be available from state enforcement agencies or the Cooperative Extension on the application of this product.

The best drift management strategy and most effective way to reduce drift potential are to apply large droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperty, or under unfavorable environmental conditions (see **Wind, Temperature and Humidity, and Temperature Inversions**).

#### Controlling Droplet Size:

Volume - Use high flow rate nozzies to apply the highest practical spray volume. Nozzies with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

-Number of Nozzles - Use the minimum number of nozzles that provide uniform coverage.

 Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is recommended practice. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

#### MANAGING OFF-TARGET MOVEMENT (continued)

•Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift. **DO NOT** use nozzles producing a mist droplet spray.

**Application Height:** Making applications at the lowest possible height (ground driven spray boom) that is safe and practical reduces exposure of droplets to evaporation and wind.

**Swath Adjustment:** When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the application equipment upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droolets, etc.).

Wind: Drift potential is lowest between wind speeds of 3-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 3 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

**Temperature and Humidity:** When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud, which can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Wind Erosion: Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.

**Ground Application (Broadcast)**: Use 5 or more gallons of water per acre (the maximum recommended spray volume for **Cadre® herbicide** is 10 gallons per acre). The actual minimum spray volume per acre is determined by the spray equipment used. Use adequate spray volume to provide accurate and uniform distribution of spray particles over the treated area and to avoid spray drift.

#### **MIXING INSTRUCTIONS**

Fill the spray tank one-half to three-quarters full with clean water. Use a calibrated measuring device to measure the required amount and add the required amount of **Cadre** to the spray tank while agitating. Fill the remainder of the tank with water

Add a nonionic surfactant, organosilicate adjuvant or crop oil concentrate to the spray tank. Maintain agitation while spraying to ensure a uniform spray mixture. An antiforming agent may be added to the tank if needed.

When tank-mixing **Cadre** with recommended herbicides, add wettable powders, dispersible granules or other dry formulations first, then **ECs**, then **Cadre**, and then an adjuvant.

To avoid injury to sensitive crops, spray equipment used for **Cadre** applications must be drained and thoroughly cleaned with water before applying other products or spraying other crops.

#### **SPRAYING INSTRUCTIONS**

**DO NOT** apply if wind conditions, temperature inversion conditions, or other conditions may cause drift onto adjacent areas or sensitive crops. Sensitive crops include, but are not limited to, leafy vegetables and cotton.

DO NOT apply if rainfall is threatening. Rainfall within 3 hours after Cadre® herbicide application may reduce weed control.

#### **Ground Application:**

Uniformly apply with properly calibrated ground equipment in 10 or more gallons of water per acre. A spray pressure of 20 to 40 psi is recommended.

To ensure proper spray coverage, the sprayer must be calibrated to deliver the recommended spray volume and pressure and the spray boom height adjusted to ensure proper coverage of weed foliage (according the manufacturer's recommendation). Spray nozzle tips should be selected to provide an even and thorough distribution of the spray mixture. The use of boomless or flood type nozzles is not recommended and may result in decreased weed control.

DO NOT apply by helicopter, airplane, or any other aerial equipment.

Avoid overlaps when spraying.

#### **APPLICATION INSTRUCTIONS**

Cadre is active on many grass weeds, but a soil-active grass herbicide such as PROWL® or Sonalan™ herbicides should be applied according to label directions before Cadre use.

Apply **Cadre** at a rate of 4.0 ounces per acre (0.063 lbs. a.i./A) plus an approved spray adjuvant (refer to the **SPRAY ADJUVANTS** section of this label) early postemergence to control broadleaf and grass weeds in peanuts. Refer to the **WEEDS CONTROLLED** chart below for weed species controlled.

#### **WEEDS CONTROLLED**

An early posternergence application of **Cadre**, at a use rate of 4.0 ounces per acre, will control or suppress the weeds listed below.

BROADLEAF WEEDS CONTROLLED	MAXIMUM HEIGHT AT APPLICATION (inches)
Anoda, Spurred	2
Burgherkin	` 2
Carpetweed	2
Citronmelon	2
Cocklebur, Common	6
Crownbeard, Golden	2
Indigo, Hairy	. 2
Morningglory, ,Cypressvine ,Entireleaf ,Ivyleaf ,Pitted ,Smallflower ,Tall	3 3 3 3 3 3 3

WEEDS CONTROLLED (continued)

An early postemergence application of Cadre® herbicide, at a use rate of 4.0 ounces per acre®, will control or suppress the weeds listed below.

BROADLEAF WEEDS CONTROLLED	MAXIMUM HEIGHT AT APPLICATION (inches)
Pigweed	
Amaranth, Palmer	2
, Redroot	4
, Smooth	4
, Spiny	4
Poinsettia, Wild	2
Pusley, Florida	2
Radish, Wild	4 .
Redweed	4
Senna, Coffee	3
Sicklepod	3
Sida, Prickly	2
Spurge Spp.	2
Starbur, Bristly	2
Velvetleaf	2

WEEDS CONTROLLED (continued)

An early postemergence application of Cadre® herbicide, at a use rate of 4.0 ounces per acre®, will control or suppress the weeds listed below.

BROADLEAF WEEDS SUPPRESSED	MAXIMUM HEIGHT AT APPLICATION (inches)
Beggarweed, Florida	2
Lamsbsquarter, Common	2
Ragweed, Common	2

GRASS WEEDS CONTROLLED*	MAXIMUM HEIGHT AT APPLICATION (inches)
Crabgrass,	
, Large	4
, Smooth	4
Crowfootgrass	2
Johnsongrass,	
, Rhizome**	8 - 10
, Seedling	4
Panicum,	
, Fall	4
, Texas	2
Sandbur spp.	4
Signalgrass, Broadleaf	. 4

#### WEEDS CONTROLLED (continued)

An early postemergence application of **Cadre® herbicide**, at a use rate of 4.0 ounces per acre\*, will control or suppress the weeds listed below.

GRASS WEEDS SUPPRESSED	MAXIMUM HEIGHT AT APPLICATION (inches)
Goosegrass	2

SEDGES CONTROLLED	MAXIMUM HEIGHT AT APPLICATION (inches)
Nutsedge,	
,Purple	4
,Yellow	4

<sup>\*</sup>Cadre will control many grass weeds which escape from the application of a soil-applied grass herbicide. However, Cadre should be used as a component of a grass weed control program and following the application of a soil-applied grass herbicide. Grass weeds must be present at the time of application to obtain control.

#### SPRAY ADJUVANTS

#### West Texas, New Mexico and Oklahoma .

DO NOT use a nonionic surfactant as an adjuvant. Include either a crop oil concentrate, or methylated seed oil concentrate or blends of these containing an organosilicate-based surfactant at 1 quart per acre. Maintain agitation while spraying to ensure a uniform spray mixture.

#### Areas outside of West Texas, New Mexico and Oklahoma

Aways include a nonionic surfactant or crop oil concentrate with **Cadre** applications. Use a nonionic surfactant with at least 80% active ingredient at one quart of surfactant for each 100 gallons of spray solution. If using crop oil concentrate, add one quart per acre. Under adverse application conditions (dry weather, larger weeds), the use of a crop oil concentrate at one quart per acre, and fertilizer (spray grade ammonium sulfate at 2.5 lbs per acre or liquid fertilizer at 1-2 quarts per acre), is recommended.

<sup>\*\*</sup>For control of rhizome johnsongrass, weeds must be at least 8 to 10 inches tall at application. Smaller weeds do not generally have enough leaf surface area to take up enough Cadre for complete control.

#### **CULTIVATION**

The control of difficult weeds (such as Florida beggarweed) and weeds treated under dry conditions is often greatly enhanced by a timely cultivation. Cultivation should be done at least 14 days after **Cadre® herbicide** application.

**DO NOT** cultivate prior to 14 days after **Cadre** application since this timing is too early to take full advantage of the weed control activity offered by **Cadre**. In addition, cultivations should be shallow to avoid excessive movement of treated soil and to avoid exposing weed seed buried deep within the soil.

#### HERBICIDE COMBINATIONS

Cadre may be tank-mixed with other herbicides if the practice is not prohibited by the label of the tank-mix partner. When Cadre is tank-mixed with another herbicide, read each label carefully to determine use rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label. No labeled use rate may be exceeded. DO NOT mix Cadre with any product whose label prohibits tank-mixes.

Using Gramoxone™ Max or Classic® herbicides in tank-mixtures with **Cadre** may result in increased peanut injury.

Using Basagran® herbicide in tank-mixtures with Cadre may result in reduced broadleaf weed control.

Using a postemergence grass control herbícide or fungicide in tank-mixtures with Cadre may result in reduced weed control.

It is not recommended to use **Cadre** in combination with or following a **PURSUIT®** or Strongarm™ herbicide application due to the potential for herbicide resistance development and uncertainty regarding crop response.

#### **ROTATIONAL CROPS**

The following rotational crops may be planted after applying Cadre in peanuts:

1. Any interval after Cadre application:

Peanuts

2. Four months after Cadre application:

Bahiagrass

Rve

Wheat

3. Nine months after Cadre application:

Field Corn

Snapbeans

Southern Peas

Soybeans

Tobacco ·

#### **ROTATIONAL CROPS** (continued)

4. Eighteen months after a Cadre® herbicide application:

Barley

Cotton\*

Grain Sorghum

Oats

Onions\*\*

Sweet Corn

5. Twenty-six months after a Cadre application:

All crops not otherwise listed.

6. Forty months after Cadre application:

Canola

Potatoes

Red Table Beets

Sugar Beets

Use of **Cadre** in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

Application of products containing chlorimuron ethyl (e.g., Classic® herbicide), or imazethapyr (e.g., **PURSUIT® herbicide**) the same year as labeled rates of **Cadre** may increase the risk of injury to sensitive rotational crops. Consult labels for recommended uses of these products in combinations. Always follow the more restrictive label limitations and precautions.

\*For Arizona, Arkansas, New Mexico, Oklahoma, and Texas only: Cotton may be planted 18 months after **Cadre** application in the states of Arizona, Arkansas, New Mexico, Oklahoma, and Texas unless drought conditions develop the year of **Cadre** application (**DO NOT** rotate to cotton at 18 months after **Cadre** application if less than 15 inches of rainfall or irrigation is received from the time of **Cadre** application through November 1 of the same year). If drought conditions develop the year of **Cadre** application, cotton may be planted 26 months after **Cadre** application.

\*\*For Florida and Georgia only.

### WEED LIST

MEED FIST		
BROADLEAF WEEDS	SCIENTIFIC NAME	
Anoda, Spurred	(Anoda cristata)	
Beggarweed, Florida	(Desmodium tortuosum)	
Burgherkin	(Cucumis anguria)	
Carpetweed	(Mollugo verticillata)	
Citronmelon	(Citrullus lanatus var. citroides)	
Cocklebur, Common	(Xanthium strumarium)	
Crownbeard, Golden	(Verbesina encelioides)	
Indigo, Hairy	(Indigofera hirsuta)	
Lambsquarters, Common	(Chenopodium album)	
Morningglory,		
, Cypressvine	(Ipomoea quamoclit)	
, Entireleaf	(Ipomoea hederacea var. integriuscula)	
, lvyleaf	(Ipomoea hederacea)	
, Pitted	(Ipomoea lacunosa)	
, Smallflower	(Jacquemontia tamnifolia)	
, Tall	(Ipomoea purpurea)	
Pigweed spp.	•	
, Amaranth, Palmer	(Amaranthus palmeri)	
, Redroot	(Amaranthus retroflexus)	
, Smooth	(Amaranthus hybridus)	
, Spiny	(Amaranthus spinosus)	
Poinsettia, Wild	(Euphorbia heterophylla)	
Pusley, Florida	(Richardia scabra)	
Radish, Wild	(Raphanus raphanistrum)	
Ragweed, Common	(Ambrosia artemisiifolia)	

WEED LIST (continued) **BROADLEAF WEEDS** SCIENTIFIC NAME Redweed (Melochia corchorifolia) Senna, Coffee (Cassia occidentalis) Sicklepod (Cassia obtusifolia) Sida, Prickly (Sida spinosa) Spurge spp. (Euphorbia spp.) Starbur, Bristly (Acanthospermum hispidum) Velvetleaf (Abutilon theophrasti) **GRASS WEEDS** SCIENTIFIC NAME Crabgrass, (Digitaria sanguinalis) , Large , Smooth (Digitaria ischaemum) Crowfootgrass (Dactyloctenium aegyptmum) (Eleusine indica) Goosegrass (Sorghum halepense) Johnsongrass Panicum, , Fall (Panicum dichotomiflorum) ,Texas (Panicum texanum) (Cenchrus spp.) Sandbur spp. Signalgrass, Broadleaf (Brachiaria platyphylla) **SEDGES** SCIENTIFIC NAME Nutsedge,

(Cyperus rotundus)

(Cyperus esculentus)

, Purple

, Yellow

#### **Conditions of Sale and Warranty**

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT PERMITTED BY LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale** and **Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

## USES WITH OTHER PRODUCTS (TANK-MIXES)

If this product is used in combination with any other product except as specifically recommended in writing by BASF then BASF shall have no liability for any loss, damage, or injury arising out of its use in any such combination not so specifically recommended. If used in combination recommended by BASF, the liability of BASF shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the BASF product in such combination use, and in any event shall be limited to return of the amount of the purchase price of the BASF product.

Basagran, Cadre, Prowl and Pursuit are registered trademarks of BASF
Broadstrike, Sonalan and Strongarm are trademarks of Dow AgroSciences LLC
Accent, Basis, Classic, Concert, Pinnacle, and Staple are registered trademarks of E.I. du Pont de Nemours & Co., Inc.
Permit is a registered trademark of Nissan Chemical Industries, Ltd.
Exceed is a registered trademark of Novartis Crop Protection, Inc.
Gramoxone is a registered trademark of Syngenta Crop Protection, Inc.

© 2008 BASF Corporation
All rights reserved.
000241-00365.20080205.**NVA 2008-04-265-0027**Supersedes: **NV**A 2004-04-265-0245

BASF Corporation Crop Protection 26 Davis Drive Research Triangle Park, NC 27709



12) |22



### Electronic Labeling Record

LABEL INFORMATION

Record Id#: ELL002260

Attachment Name: 000241-00364.20080207.NVA2008-04-265-0027.

**Product** Active

Status:

Registration #: 000241-00364

Product: CADRE HERBICIDE

pdf

PC Code: 128943

RM Team: RM 25

**Submission** 02/07/2008

Date:

Receipt Date:

Version: Initial

CD Enclosed?: Yes

Comments: NVA2008-04-265-0027

Application Correspondence:

Attachment:

000241-00364.20080207.NVA2008-04-265-0027.pdf

#### SUBMISSION INFORMATION

Status: Pending

Submisssi 824533

on BC:

Action 332

Code:

**OPPIN Receipt Date:** 02/08/2008

Action Comments: Notification per PRN 2007-4: Revision of Cadre Herbicide container disposal

language

**Reviewer Comments:** 

User Reviews & Response Letters