

## **PLEASE NOTE**

**This image contains more than one label approved for this product on this date.**



# Supplemental Labeling

EPA Reg. No. 241-365

## FOR USE OF PLATEAU HERBICIDE ON CONSERVATION RESERVE PROGRAM (CRP) LAND

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PLATEAU HERBICIDE LEAFLET LABEL BEFORE USING. SEE THE PLATEAU HERBICIDE LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION AND WEEDS CONTROLLED.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application.

DO NOT apply this product through any type of irrigation system.

### GENERAL INFORMATION

PLATEAU herbicide is an aqueous solution to be mixed with water and an adjuvant and applied as a spray solution to provide weed control and/or turf height suppression on noncropland areas such as railroad, utility, pipeline and highway rights-of-way, railroad crossings, utility plant sites, petroleum tank farms, pumping installations, non-agricultural fence rows, storage areas, non-irrigation ditchbanks, Conservation Reserve Program (CRP) land (see USE OF PLATEAU HERBICIDE ON CONSERVATION RESERVE PROGRAM LAND section), prairie sites, airports, industrial turf, recreational and non-residential turf and other similar areas.

### USE OF PLATEAU HERBICIDE ON CONSERVATION RESERVE PROGRAM (CRP) LAND

PLATEAU herbicide may be used on Conservation Reserve Program (CRP) land at rates up to 4 oz. per acre per year (see minimum plant-back intervals below). See appropriate section of this label for specific instructions for the intended use. DO NOT use rates higher than 4 oz per acre per year on CRP land.

61: E D 1E TT 96

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

AUG 26 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act (See reverse side) as amended, for the pesticide registered under EPA Reg. No.

241-365

RECD EPA/OPP/DPD1

MINIMUM PLANT-BACK INTERVALS (months after PLATEAU herbicide application)				
4	9	18	26	40
Bahiagrass Rye Wheat	Field Corn Snapbeans Southern Peas Soybeans Tobacco	Barley Cotton* Grain Sorghum Oats Sweet Corn	All crops not otherwise listed	Canola Potatoes Red Table Beets Sugar Beets

**\*For New Mexico, Oklahoma, and Texas only:** Cotton may be planted 18 months after PLATEAU herbicide application in the states of New Mexico, Oklahoma, and Texas unless drought conditions develop the year of PLATEAU herbicide application. DO NOT rotate to cotton at 18 months after PLATEAU herbicide application if less than 15 inches of rainfall or irrigation is received from the time of PLATEAU herbicide application through November 1 of the same year. If drought conditions develop the year of PLATEAU herbicide application, cotton may be planted 26 months after PLATEAU herbicide application.

Use of PLATEAU herbicide in accordance with label directions is expected to result in normal growth of plant-back crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, plant-back crop injury is always possible.

### NATIVE PRAIRIEGRASS RENOVATION AND RESTORATION

PLATEAU herbicide may be applied at a rate of up to 4 oz per acre to Conservation Reserve Program (CRP) land for the establishment or release of big bluestem, little bluestem, Indiangrass, sideoats and blue grama (see NATIVE PRAIRIEGRASS RENOVATION AND RESTORATION section of the label for details).



**NEXT**

**LABEL**



# Supplemental Labeling

EPA Reg. No. 241-365

## AERIAL APPLICATION

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PLATEAU HERBICIDE LEAFLET LABEL BEFORE USING. SEE THE PLATEAU HERBICIDE LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION AND WEEDS CONTROLLED.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application.

DO NOT apply this product through any type of irrigation system.

REC'D EPA/OPP/DPD1  
96 JUL 31 P 3:23

### SPRAYING INSTRUCTIONS

All precautions should be taken to minimize or eliminate spray drift. Fixed wing aircraft and helicopters can be used to apply PLATEAU herbicide, however, DO NOT make applications by fixed wing aircraft unless appropriate buffer zones can be maintained to prevent spray drift out of the target area, or when treating open tracts of land, or where spray drift as a result of fixed wing aircraft application can be tolerated. Aerial equipment designed to minimize spray drift such as a helicopter equipped with a MICROFOIL boom, or THRU-VALVE boom or raindrop nozzles, must be used and calibrated. Except when applying with a MICROFOIL boom, a drift control agent may be added at the recommended label rate. To avoid drift, applications should not be made during inversion conditions, when winds are gusty, or under any other conditions which promote spray drift.

Uniformly apply recommended amount of PLATEAU herbicide in 5 or more gallons of water per acre, using enough volume to provide adequate coverage of target area or foliage. Include an adjuvant in the spray solution (See "Adjuvants" Section). A foam reducing agent may be added at the recommended rate, if needed.

**IMPORTANT:** Thoroughly clean application equipment, including landing gear, immediately after use of this product. Prolonged exposure of this product to uncoated steel (except stainless steel) surfaces may result in corrosion and failure of the exposed part. The maintenance of an organic coating (paint) may prevent corrosion.

ACCEPTED  
~~with COMMENTS~~  
In EPA Letter Dated  
AUG 26 1996



Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 241-365