



000241-00376-010599

Printed: 07:45:30 Tuesday, 08 Jun, 1999 # 12 / 3224

Systems Integration Group, Inc.

STEEL®
herbicide

Supplemental Labeling

EPA Reg. No. 241-376

FOR USE IN SOYBEANS
THIS LABEL EXPIRES DECEMBER 31, 1999

ROTATIONAL CROP RESTRICTIONS FOR USE AREAS IN CENTRAL NEBRASKA AND KANSAS

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS IN THE STEEL LABEL BEFORE USING. SEE THE STEEL LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION AND USES WITH OTHER PRODUCTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This labeling must be in the possession of the user at the time of herbicide application.

A single preplant incorporated or preemergence application of STEEL may be made at the 3 pint per acre rate following instructions for USE REGION 3 in the leaflet label. Consult the STEEL label for mixing and application information, weeds controlled, and use rates. When tank mixing, always follow the more restrictive label.

ROTATIONAL CROP RESTRICTIONS

Refer to the ROTATIONAL CROP GUIDELINES listed on the reverse side of this supplemental labeling for applications of STEEL in the counties listed below:

KANSAS		NEBRASKA		
BARBER	PHILLIPS	ADAMS	GARFIELD	MADISON
BARTON	PRATT	ANTELOPE	GOSPER	MERRICK
COMANCHE	RENO	BLAINE	GREELEY	NANCE
EDWARDS	RICE	BOONE	HALL	NUCKOLLS
ELLIS	ROOKS	BOYD	HAMILTON	PERKINS
HARPER	RUSH	BROWN	HARLAN	PHELPS
KINGMAN	RUSSELL	BUFFALO	HAYES	PIERCE
KIOWA	SMITH	CEDAR	HITCHCOCK	PLATTE
OSBORNE	STAFFORD	CHASE	HOLT	POLK
PAWNEE		CLAY	HOWARD	RED WILLOW
		CUSTER	KEARNEY	ROCK
		DAWSON	KEITH	SIERMAN
		DUNDY	KEYA PAHA	VALLEY
		FRANKLIN	KNOX	WEBSTER
		FRONTIER	LINCOLN	WHEELER
		FURNAS	LOUP	

ACCEPTED
JAN - 5 1999

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-376

(see other side)

ROTATIONAL CROP GUIDELINES

The following rotational crops may be planted after applying STEEL in soybeans:

CROP	ROTATIONAL INTERVAL
Soybeans	No restrictions
IMI-CORN® seed hybrids ¹	9.5 months
Field Corn (non IMI-CORN)	18 months ²
Wheat	4 months ³

¹ Contact your chemical dealer, seed supplier, or American Cyanamid to obtain information regarding the availability of imidazolinone tolerant field corn hybrids which are adapted to your area.

² Field corn (non-resistant) may be planted as a rotational crop 18 months following the application of STEEL unless extreme drought conditions develop (less than 15 inches of rainfall or irrigation is received from two weeks prior to the date of STEEL application through November 15 of the same year).

If the minimum rainfall requirement is not met only IMI-CORN field corn hybrids which possess

tolerance or resistance to STEEL and other imidazolinone herbicides may be planted the spring of the year following the 18-month crop rotation period.

³ Wheat may be planted as a rotational crop 4 months following the application of STEEL unless extreme drought conditions develop (less than 10 inches of rainfall or irrigation is received from two weeks prior to the date of STEEL application through September 15 of the same year). If the minimum rainfall requirement is not met, do not plant wheat until the fall of the year after the STEEL application.

For rotational guidelines to other crops, refer to the STEEL leaflet label.

Use of STEEL in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with use of this product and, therefore, rotational crop injury is always possible.



® Registered Trademarks of American Cyanamid Company

PE-49019 Rev. 1 11/30/98