


PM 25 241-379 4-28-98 10/2
5 2
RAPTORTM

herbicide

**Supplemental
Labeling**



EPA Reg. No. 241-379

FOR USE IN SOYBEANS

**TANK MIXTURES OF RAPTOR AND CLASSIC¹ APPLIED
POSTEMERGENCE IN SOYBEANS**

THIS LABEL EXPIRES APRIL 30, 1999

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE RAPTOR AND CLASSIC FULL LEAFLET LABELS BEFORE USING. SEE THE RAPTOR AND CLASSIC LEAFLET LABELS FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, WEEDS CONTROLLED, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. The use of RAPTOR herbicide not consistent with this labeling may result in injury to crops. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

GENERAL INFORMATION

RAPTOR provides postemergence control of many broadleaf and grass weeds in either conventional or no-till soybeans. RAPTOR may be applied to all soybean varieties including both Roundup Ready² and non-glyphosate-resistant soybeans.

Classic may be applied with RAPTOR in a tank-mixture for enhanced postemergence control of sicklepod, bristly starbur, Florida beggarweed, hemp sesbania, marestalk, pitted morningglory, and yellow nutsedge.

When tank-mixing herbicides, always follow the most restrictive label.

USE RATE


Apply 4 fluid ounces of RAPTOR herbicide per acre with 0.5 ounces per acre of Classic with a non-ionic surfactant when preceded by a full rate of a registered soil applied grass herbicide like PROWL[®] 3.3 EC herbicide

OR

Apply 5 fluid ounces of RAPTOR herbicide per acre with 0.5 ounces per acre of Classic with a nonionic surfactant in a total postemergence herbicide program.

Do not apply the tank-mixture with crop oil concentrate or methylated seed oil.

(see other side)



20/2

ROTATIONAL CROP INFORMATION

Use of RAPTOR herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible. **Always follow the most restrictive label.**

Refer to the RAPTOR and Classic leaflet labels for additional information regarding ROTATIONAL CROP INFORMATION, WEEDS CONTROLLED, MIXING AND SPRAYING INSTRUCTIONS, PRECAUTIONARY STATEMENTS AND WORKER PROTECTION INFORMATION.



American Cyanamid Company
North America Agricultural Products Division
Crop Protection Products Department
One Campus Drive
Parsippany, NJ 07054 ©1998

¹Trademark of E.I. duPont de Nemours & Co., Inc.

²Trademark of Monsanto Products Company

TMTrademark of American Cyanamid Company

PE-XXXXXX 04/08/98

