

RAPTOR[®] herbicide

Supplemental Labeling

EPA Reg. No. 241-379

FOLLOW CROP GUIDELINE FOR LAND ENROLLED IN THE CONSERVATION RESERVE PROGRAM

THIS LABEL EXPIRES APRIL 30, 2002

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE RAPTOR FULL LEAFLET LABEL BEFORE USING. SEE THE RAPTOR LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, WEEDS CONTROLLED, USES WITH OTHER PRODUCTS AND ADDITIONAL ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling. The use of RAPTOR herbicide not consistent with this labeling may result in injury to crops. **DO NOT** apply this product through any type of irrigation system. This labeling must be in the possession of the user at the time of herbicide application.

ROTATIONAL CROP INFORMATION

In addition to selected crops and rotational intervals listed on the RAPTOR leaflet label, the following rotational crops may be planted **nine months after applying up to 5 fluid ounces per acre of RAPTOR herbicide:**

Big Bluestem	Little Bluestem	Sand Bluestem
Smooth Bromegrass	Green Needlegrass	Indiangrass
Purple Prairieclover	Prairie Sandreed	Reed Canarygrass
Sideoats Grama	Sweetclover	Switchgrass
Crested Wheatgrass	Intermediate Wheatgrass	Tall Wheatgrass
Western Wheatgrass	Other native forbs and legumes	

Use of RAPTOR herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.



American Cyanamid Company
 North America Agricultural Products Division
 Crop Protection Products Department
 One Campus Drive
 Parsippany, NJ 07054 ©1999