

PM 25 241-386

2-10-94

10/3

Pursuit W
Herbicide

Supplemental Labeling

ACCEPTED

FEB 10 1999
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-386

EPA Reg. No. 241-386

FOR USE ON PEANUTS

IN ARIZONA (Yuma and La Paz Counties) ONLY

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT W DG LABELING BEFORE USING. SEE THE PURSUIT W DG LEAFLET LABEL FOR PRECAUTIONARY STATEMENTS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. The use of PURSUIT W DG herbicide not consistent with this labeling may result in injury to crops. **DO NOT** apply this product through any type of irrigation system. This labeling must be in the possession of the user at the time of herbicide application.

GENERAL INFORMATION

PURSUIT W DG kills weeds by root and/or foliage uptake and rapid translocation to the growing points. Adequate soil moisture is important for optimum PURSUIT W DG activity. When adequate soil moisture is present, PURSUIT W DG will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

PURSUIT W DG HERBICIDE USE RATE (1.44 OUNCES PER ACRE)

Apply PURSUIT W DG at a broadcast rate of 1.44 ounces per acre for all methods of application (except sequential - see below): preplant incorporated, preemergence, ground-cracking and postemergence. At this broadcast rate, one soluble packet of PURSUIT W DG will treat 2 acres of peanuts.

PURSUIT W DG may also be applied in a sequential application. Apply 0.72 ounces in a soil application (preplant incorporated or preemergence) followed by 0.72 ounces applied at ground-crack or postemergence. (At the 0.72 ounce rate, one packet will treat 4 acres).

NOTE: Do not apply more than 1.44 ounces of PURSUIT W DG during the growing season.

WEEDS CONTROLLED

When applied as directed, PURSUIT W DG will control or reduce competition from the weeds listed below. Refer to the MIXING INSTRUCTIONS section of the leaflet label for recommendations when weeds are at the maximum recommended growth stage, or are under stress.

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence).

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

SOIL APPLIEDAT-CRACKPOSTEMERGENCE 2015**BROADLEAF Weeds Controlled**Maximum
Leaf Stage Size (inches)

Lambsquarters, common	C*	C	R	1-2
Morningglory				
entireleaf	R	C	2	1-2
ivyleaf	R	C	2	1-2
pitted	R	C	2	1-2
smallflower	C	C	4	1-3
tall	R	C	2	1-2
Mustard sp.	C	C	4	1-3
Nightshade				
black	C	C	4	1-3
Eastern black	C	C	4	1-3
hairy	C	C	4	1-3
Pigweed				
redroot	C	C	8	1-8
smooth	C	C	8	1-8
spiny	C	C	8	1-8
Puncturevine	C	C		
Purslane, common	C	C		
Smartweed				
ladysthumb	C	C	4	1-3
Pennsylvania	C	C	4	1-3
Spurge				
prostrate	C	C	4	1-3
spotted	C	C	4	1-3
toothed	C	C		
Sunflower	C*	C	4	1-3

GRASS Weeds Controlled**

Barnyardgrass	R	R	3	1-3
Foxtail,				
giant	C	C	6	1-6
green	C	C	3	1-3
yellow	C	C	3	1-3
Johnsongrass,				
seedling	C	C	6	1-8
rhizome			R	6-12

SEDGES Controlled

Nutsedge				
purple	R	R	3	1-3
yellow	R	R	3	1-3

* When PURSUIT W DG is soil applied, these weeds are more consistently controlled by preplant incorporated treatments.

** When PURSUIT W DG is soil applied to grasses, more consistent control can be obtained from preplant incorporated treatments.

Do not count cotyledon leaves when determining weed stage of growth.

AERIAL APPLICATION

Do not apply to peanuts by aerial application.

PREPLANT INCORPORATED APPLICATIONS

When PURSUIT W DG is soil applied for control of nutsedge in peanuts, incorporate with two passes of the incorporation implement. Make the second pass at an offset angle to the first pass to minimize the potential fort streaking.

AT CRACK APPLICATION refers to the time when the soil cracks due to the emerging peanut seedling. This generally occurs from 10 to 14 days following planting. At this time weeds have generally not germinated, or are in the seedling stage. If weeds have more than 2 true leaves, refer to the **POSTEMERGENCE** weed control column for weeds controlled.

Wait until late cracking (most of the peanuts have emerged) before applying PURSUIT W DG.

PURSUIT W DG is active against many broadleaf and grass species. However, when heavy grass or common lambsquarters pressure is anticipated, PURSUIT W DG should be used in combination with a registered soil-applied grass herbicide.

PRECAUTIONS

Do not graze or feed peanut forage, vines, hay or straw to livestock.

There should be an interval of at least 85 days between an application of PURSUIT W DG and peanut harvest.

Do not apply PURSUIT PLUS® EC or PASSPORT® to peanuts the same year as PURSUIT W DG.

®Registered Trademarks of American Cyanamid Company

PE-XXXXXX 10/6/98

8279



American Cyanamid Company
North America Agricultural Products Division
Crop Protection Products Department
One Campus Drive
Parsippany, NJ 07054 ©1999

