PLEASE NOTE

This image contains more than one label approved for this product on this date.

ŧ.

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

241-386

(-

ł

3-2-2001

MAR 2 2001

Ms. Rebecca L Johnston BASF Corporation P.O. Box 400 Princeton, NJ 08543-0400

Dear Ms. Johnston:

SUBJECT: New Supplemental Label Adding Postemergence Use in Dry Edible Peas Pursuit® W DG Herbicide EPA Registration No. 241-386 Your Submission Dated February 7, 2001

The supplemental labeling referred to above and submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable, provided you correct the following error:

1. In the Rotational Crop Guidelines, delete <u>Rve</u> from the list of crops that may be planted eighteen months after application, since Rye may be planted four months after application of Pursuit W Herbicide. It already correctly appears with a four-month plant back interval on this label and the main product label.

At your next label printing, or within one year, whichever comes first, you must incorporate this supplemental labeling for postemergence use in dry edible peas into the main product labeling. A stamped copy of the label is enclosed for your records. Please submit one copy of your final printed label before you release the product for shipment.

Sincerely yours,

usa a. Stanton for

Daniel J. Rosenblatt Acting Product Manager (25) Herbicide Branch Registration Division (7505C)


Enclosure

RD:STANTON:PM Team 23:Rm. 239:CM-2:305-5218:Disk #14:S592706.LET

CONCURRENCES								
SYMBOL +	7505C							
SURNAME +	S. Stanton			1				
DATE •	Mar 2, 2001							
EPA Form 1320-1 (12-70)					OF	OFFICIAL FILE COPY		



Supplemental Labeling



EPA Reg. No. 241-386

FOR POSTEMERGENCE USE IN DRY EDIBLE PEAS

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF IDAHO, MONTANA, OREGON, AND WASHINGTON

This Label Expires April 30, 2002

OBSERVE ALL PRECAUTIONARY STATEMENTS, SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT W DG LEAFLET LABEL BEFORE USING.

DISCLAIMER

THIS PRODUCT WHEN USED ON EDIBLE LEGUME VEGETABLE CROPS MAY LEAD TO CROP INJURY, LOSS, OR DAMAGE. BASF CORPORATION RECOMMENDS THAT THE USER AND/OR GROWER TEST THIS PRODUCT IN ORDER TO DETERMINE ITS SUITABILITY FOR SUCH INTENDED USE. BASF MAKES THIS PRODUCT AVAILABLE TO THE USER AND/OR GROWER SOLELY TO THE EXTENT THE BENEFIT AND UTILITY, IN THE SOLE OPINION OF THE USER AND/OR GROWER, OUTWEIGH THE EXTENT OF POTENTIAL INJURY ASSOCIATED WITH THE USE OF THIS PRODUCT. THE DECISION TO USE OR NOT TO USE THIS HERBICIDE MUST BE MADE BY EACH INDIVIDUAL PURSUIT W DG USER AND/OR GROWER ON THE BASIS OF POSSIBLE CROP INJURY FROM PURSUIT W DG, THE SEVERITY OF WEED INFESTATION, THE COST OF ALTERNATIVE WEED CONTROLS, AND OTHER FACTORS. BASF INTENDS THAT BECAUSE OF THE RISK OF FAILURE TO PERFORM OR CROP DAMAGE THAT ALL SUCH USE IS AT THE USER'S AND/OR GROWER'S RISK.

BASF DISCLAIMS ANY LIABILITY FOR CLAIMS, CAUSES OF ACTION, FINES, PENALTIES, DAMAGES, INCLUDING CONSEQUENTIAL INCIDENTS AND DAMAGES, LOSSES, LIABILITIES, JUDGMENTS, AND EXPENSES ARISING OUT OF OR RELATING TO INJURY TO PERSONS, CROPS, OR PROPERTY

RESULTING FROM THE USE OF PURSUIT W DG HERBICIDE ON EDIBLE LEGUMES CONTRARY TO THE LABEL INSTRUCTIONS. BASF WARRANTS ONLY THAT THE MATERIAL CONTAINED HEREIN CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL AND JS REASONABLY FIT FOR THE USE THEREIN DESCRIBED WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE, SUBJECT TO THE RISKS REFERRED TO ABOVE.

BASF DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER, IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMUTED TO REPAYMENT OF THE PURCHASE PRICE OF PURSUIT W DG. In no case shall BASF or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

GENERAL INFORMATION

PURSUIT W DG may be applied POSTEMERGENCE to dry edible peas only. DO NOT apply PURSUIT W DG POSTEMERGENCE to dry edible peas in geographical regions outside those specifically defined in this label.

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT W DG if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT W DG if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT W DG postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT W DG ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall) PURSUIT W DG may cause injury to subsequent planted crops. Sugar beets are particularly sensitive to PURSUIT W DG residues in soil. PURSUIT W DG is not recommended in crop rotations that include sugar beets. See the ROTATIONAL CROP section of this supplemental label for rotations to other sensitive crops.

USE AREA

PURSUIT W DG may be used POSTEMERGENCE in Montana, Washington, Oregon, and Idaho only.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT W DG in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT W DG.

Spray equipment used for PURSUIT W DG application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash.¹

DO NOT make more than one application of PURSUIT W DG per year.

When PURSUIT W DG is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT W DG cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT feed treated forage, hay, or straw to livestock.

ž

Allow at least 60 days between application and harvest.

Postemergence Applications. Apply PURSUIT W DG at 0.72 ounces per acre (one ECO-PAK soluble bag treats 10 acres). A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Basagran¹ may be tank-mixed with PURSUIT W DG to control weeds not listed on the PURSUIT W DG label. Addition of Basagran may also cause antagonism, thereby reducing control of grass weeds. Nitrogen-based fertilizer may be included as a spray additive **ONLY** when PURSUIT W DG is tank-mixed with Basagran. Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions.

DO NOT APPLY PURSUIT W DG POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

WEEDS CONTROLLED

PURSUIT W DG applied at the broadcast rate of 0.72 ounces per acre will control:

wild mustard black nightshade* eastern black nightshade* hairy nightshade*

*suppression

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT W DG as recommended in this label: (Planting earlier than the recommended interval may result in crop injury).

- 1. Anytime: Soybeans Peas Lima beans Lentils Chickpeas CLEARFIELD* corn seed hybrids
- Four months after PURSUIT W DG application: Alfalfa Rye Edible bean and pea types not listed on this label Wheat
- Nine and one-half months after PURSUIT W DG application: Barley Field corn Field corn grown for seed*
- 4. Eighteen months after PURSUIT W DG application:

Cotton	Rye
Lettuce	Safflower
Oats	Sorghum
Popcorn	Sunflower
Sweet Corn	

- Twenty-six months after PURSUIT W DG application: Potatoes Flax
- 6. Forty months after PURSUIT W DG application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT W DG application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year. If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT W DG soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, BASF Corporation has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT W DG the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of BASF Corporation, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT W DG shall be assumed by the user.

CONSULT THE PURSUIT W DG LEAFLET LABEL FOR OTHER PRECAUTIONS.

BASF Corporation 26 Davis Drive Research Triangle Park, NC 27709



¹Dash is a trademark of BASF AG. [®]/* Trademarks of BASF.

©2001 BASF Corporation All rights reserved NVA 2001-04-130-035 01/26/01

i

NEXT

. .

LABEL.

÷

÷

241-386

ğ



Supplemental Labeling

EPA Reg. No. 241-386

FOR USE IN ALFALFA IN THE STATES OF ARIZONA AND CALIFORNIA TO AID IN THE SUPPRESSION OF DODDER

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT W DG AND TANK MIX PARTNER LEAFLET LABELS BEFORE USING. SEE PURSUIT W DG AND TANK MIX PARTNER LEAFLET LABELS FOR PRECAUTIONS, GENERAL INFORMATION, MIXING AND APPLICATION AND SPRAYING INSTRUCTIONS, USE RATES, USES WITH OTHER PRODUCTS, WEEDS CONTROLLED, ROTATIONAL CROP RESTRICTIONS AND DISCLAIMERS. OBSERVE THE MOST RESTRICTIVE INSTRUCTIONS ON ANY PRODUCT LABEL.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling. The use of PURSUIT W DG herbicide not consistent with this labeling may result in injury to crops. **DO NOT** apply this product through any type of irrigation system. This labeling must be in the possession of the user at the time of herbicide application.

For best results in suppressing dodder (*Cuscuta* spp.), apply PURSUIT W DG herbicide after dodder has emerged but prior to, or soon after attachment. PURSUIT W DG may be used on seedling or established alfalfa. Apply PURSUIT W DG when the seedling alfalfa is in the second (2^{nd}) trifoliate stage or larger. When tank-mixing herbicides, always follow the most restrictive label.

USE RATE

Apply PURSUIT W DG herbicide at a broadcast rate of 2.16 ounces per acre.

Note: A maximum of 2.16 ounces of PURSUIT W DG per year may be applied to alfalfa.



(See reverse side)

MIXING INSTRUCTIONS

Applications of PURSUIT W DG for dodder suppression require the addition of a petroleum or vegetable seed based oil concentrate. Use a crop oil concentrate at 2 pints per acre. A methylated seed oil may be used at the rate of 1.5 to 2 pints per acre.

SPRAYING INSTRUCTIONS

DO NOT apply when wind velocity is greater than 10 mph, or when spray may be carried to sensitive crops. Sensitive crops include leafy vegetables and sugar beets.

Uniformly apply with properly calibrated ground equipment in 20 or more gallons of water per acre. Applications of PURSUIT W DG for dodder suppression should be made using ground equipment only.

Adjust the boom height (according to the manufacturer's recommendation) to ensure proper coverage of dodder. Use flat-fan nozzle tips only. Avoid overlaps when spraying.

Refer to the PURSUIT W DG leaflet label for additional information regarding MIXING AND SPRAYING INSTRUCTIONS, PRECAUTIONARY STATEMENTS, WORKER PROTECTION INFORMATION AND ROTATIONAL CROP RESTRICTIONS.

BASF Corporation 26 Davis Drive Research Triangle Park, NC 27709

.....



®Registered trademark of BASF.

© 2001 BASF Corporation All rights reserved NVA 2001-04-130-033 02/07/01