

241-407

8/29/2001

1/2

Backdraft
HERBICIDE

Supplemental Labeling

EPA Reg. No. 241-407

FOR USE ON ROUNDUP¹ READY SOYBEANS

THIS LABEL EXPIRES DECEMBER 31, 2002

**FALL APPLICATION OF BACKDRAFT IN A PLANNED SEQUENTIAL PROGRAM
USING ROUNDUP¹ ULTRA ON ROUNDUP READY SOYBEANS**

**FOR DISTRIBUTION AND USE ONLY IN THE STATES OF DELAWARE, ILLINOIS,
INDIANA, IOWA, KANSAS, KENTUCKY, MARYLAND, MICHIGAN, MINNESOTA,
MISSOURI, NEBRASKA, OHIO, PENNSYLVANIA, VIRGINIA, WEST VIRGINIA
AND WISCONSIN (SEE USE AREA)**

**OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE BACKDRAFT AND ROUNDUP ULTRA
LABELING BEFORE USING. SEE THE BACKDRAFT LEAFLET LABEL FOR PRECAUTIONS,
WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, WEEDS
CONTROLLED, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.**

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

This labeling must be in the possession of the user at the time of herbicide application.

Apply BACKDRAFT after Fall Harvest and prior to ground freeze-up in the winter. Fall applications of BACKDRAFT will control existing weeds and provide residual control of winter annual weeds and early spring germinating weeds in soybeans. See the BACKDRAFT label for a complete list of weeds controlled.

Roundup Ready soybeans must be planted in the spring following the fall application of BACKDRAFT. If weeds emerge in-season, Roundup Ultra may be applied postemergence for weed control.

Refer to the BACKDRAFT product label for additional information on application instructions.

USE RATE

Apply BACKDRAFT at the rate of 4 pints per acre. If heavy grass or waterhemp pressure is expected, add PROWL[®] 3.3 EC herbicide to the spray mixture. See the PROWL 3.3 EC label for appropriate rates.

(See reverse side)

GENERAL INFORMATION

BACKDRAFT provides control of many broadleaf and grass weeds when applied preplant or preemergence. BACKDRAFT may be applied in the late fall for control of weeds in no-till or conventional soybeans, rather than in the spring prior to planting. Treated areas must be planted to Roundup Ready soybeans in the following spring, and, if required, Roundup Ultra applied in-season in a planned sequential program.

USE AREA

Fall applications of BACKDRAFT may be made only in the states of Delaware, Illinois, Indiana, Iowa, Kansas, Kentucky, Maryland, Michigan, Minnesota (south of state highway 210), Missouri, Nebraska, Ohio, Pennsylvania, Virginia, West Virginia and Wisconsin.

ROTATIONAL CROP INFORMATION

Refer to rotational crop restrictions listed in the BACKDRAFT product label.

NOTE: For fall applications of BACKDRAFT, adjust the rotational crop intervals by basing the interval on the date of soybean planting rather than the date of herbicide application.

Use of BACKDRAFT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709

BASF

™/®Trademarks of BASF
‡Trademarks of Monsanto Agricultural Products Company

©2001 BASF Corporation
All rights reserved

NVA 2001-04-107-xxxx
PE-62009 Rev. 1 07/09/01

