

241-418

3.30.2007

1/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Craig D. Kleppe, Ph.D.  
Registration Scientist  
BASF Corporation  
26 Davis Drive  
P.O. Box 13528  
Research Triangle Park, NC 27709

MAR 30 2007

**SUBJECT:** Application for Pesticide Notification – Revised Supplemental Labeling  
PROWL® H<sub>2</sub>O Herbicide  
EPA Reg. No. 241-418  
Application Dated November 13, 2006

Dear Dr. Kleppe:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the product above. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please me directly at 703-305-6249 or Terri Stowe of my staff at 703-305-6117.

Sincerely,

A handwritten signature in black ink, appearing to be "Linda Arrington".

Linda Arrington  
Notifications & Minor Formulations Team Leader  
Registration Division (7505P)  
Office of Pesticide Programs



2/12

	United States <b>Environmental Protection Agency</b> Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	---	--	-----------------------

### Application for Pesticide - Section I

1. Company/Product Number BASF Corporation / 241-418	2. EPA Product Manager James Tompkins	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) BASF Corporation / PROWL H2O herbicide	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) BASF Corporation 26 Davis Drive, PO Box 13528 Research Triangle Park, NC 27709  <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____  Product Name _____

### Section - II

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.
--	---

NOTIFICATION  
MAR 30 2007

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

NOTIFICATION to provide revised supplemental labeling for Prowl H2O herbicide on Fruiting Vegetables, Green Onions and Strawberries, in response to EPA-letter dated November 6, 2006.

No PRIA code/fee is applicable for this notification.

### Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt. _____ No. per container _____	If "Yes" Package wgt. _____ No. per container _____		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Craig D. Kleppe	Title Registration Scientist	Telephone No. (Include Area Code) 919 547 2000 ext 2615
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 		3. Title Registration Scientist
4. Typed Name Craig D. Kleppe		5. Date November 13, 2006

BASF Corporation

**BASF**

3/12

November 13, 2006

U.S. Environmental Protection Agency  
Office of Pesticide Programs (7505P)  
Document Processing Desk 7504P (NOTIF)  
Room S-4900  
One Potomac Yard (South Building)  
2777 South Crystal Drive  
Arlington, VA 22202 U.S.A.  
Attention: Mr. James A. Tompkins, PM-25

**NOTIFICATION**

**MAR 30 2007**

**RE: NOTIFICATION –**

**Revised Supplemental Labeling for PROWL® H2O Herbicide (EPA Reg.No. 241-418)  
In Response to EPA-letter Dated November 6, 2006**

Dear Mr. Tompkins:

With this letter as a notification in response to EPA letter dated November 6, 2006, BASF is providing revised supplemental labeling for the end use product **PROWL® H2O Herbicide** with **EPA Registration No. 241-418**. The following labels are attached:

1. **supplemental label** for use on **Fruiting Vegetables** –  
BASF has made the changes on the supplemental labeling as requested by EPA in letter to BASF dated November 6, 2006.
2. **supplemental label** for use on **Green Onions, Leeks, Spring Onions or Scallions, Japanese Bunching Onions, Green Shallots or Green Eschalots** –  
BASF has made the changes on the supplemental labeling as requested by EPA in letter to BASF dated November 6, 2006.
3. **supplemental label** for use on **Strawberries** –  
BASF has made the changes on the supplemental labeling as requested by EPA in letter to BASF dated November 6, 2006.

Please find enclosed the following documentation to support this notification:

1. Application Form 8570-1
2. **Exhibit 1** is the revised supplemental label for use on Fruiting Vegetables.
3. **Exhibit 2** is the revised supplemental label for use on Green Onions, etc.
4. **Exhibit 3** is the revised supplemental label for use on Strawberries.

Thank you for your continued assistance with **PROWL® H2O Herbicide**.

Regards,



Craig D. Kleppe, Ph.D.

Registration Scientist, email [craig.kleppe@basf.com](mailto:craig.kleppe@basf.com), Tel 919-547-2615, Fax 919-547-2850

® Registered Trademark of BASF

**NOTIFICATION**

MAR 30 2007

4/12



**FOR USE ON FRUITING VEGETABLES**

EPA Reg. No. 241- 418

Active ingredient:

pendimethalin (N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine)..... 38.7%

**INERT INGREDIENTS:** ..... 61.3%

**TOTAL** ..... 100.0%

OBSERVE ALL PRECAUTIONARY STATEMENTS, MIXING AND APPLICATION INSTRUCTIONS, AND FOLLOW CROP RESTRICTIONS ON THE PROWL® H<sub>2</sub>O HERBICIDE MAIN PRODUCT LABEL BEFORE USING.

**Directions For Use**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Prowl H<sub>2</sub>O controls most annual grasses and certain broadleaf weeds as they germinate. Uniformly apply Prowl H<sub>2</sub>O with properly calibrated ground equipment in 10 or more gallons of water or 20 or more gallons of liquid fertilizer per acre, or by aircraft in a minimum of 5 gallons of water per acre. Use sprayers equipped with nozzles that provide accurate and uniform application.

Adequate rainfall or irrigation after application prior to weed seedling emergence will provide the most effective weed control.

Always perform a mixing test to check the compatibility of Prowl H<sub>2</sub>O with all potential tank mix partners.

Follow all precautions and restrictions on the labels of all products applied in combination with Prowl H<sub>2</sub>O. Always follow the most restrictive label.

**Non-Target, Endangered Species**

If non-target plants occur in proximity to the application site, the following mitigation measures are required:

- If applied by ground, leave untreated buffer zone of 80 feet. The product must be applied using a low boom (20 inches

above the ground) and ASAE fine to medium/coarse nozzles.

- If applied by air, leave untreated buffer zone of 200 feet. Must use straight stream nozzles (such as D-6 to D-8). Wind can be no more than 8 mph and release height must be 15 feet or less.

If endangered species occur in proximity to the application site, the following mitigation measures are required:

- If applied by ground, leave untreated buffer zones of 200 feet. The product must be applied using a low boom (20 inches above the ground) and ASAE fine to medium/coarse nozzles.
- If applied by air, leave untreated buffer zone of 1,000 feet. Must use straight stream nozzles (such as D-6 to D-8). Wind can be no more than 8 mph and release height must be 15 feet or less

To determine whether your county has an endangered species, consult the website <http://www.epa.gov/espp/usa-map.htm>.

Endangered Species Bulletins may also be obtained from Extension offices or state pesticide agencies. If the bulletin is not available for your specific area, check with the appropriate local state agency to determine if known populations of endangered species occur in the area to be treated.

5/12

### Application Instructions

Prowl H<sub>2</sub>O may be applied to Fruiting Vegetables, such as:

- tomato
- eggplant
- pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper)
- pepino
- groundcherry (*Physalis* spp.)
- tomatillo.

Uniformly apply **PROWL H<sub>2</sub>O** as a broadcast preplant incorporated application or as a broadcast preplant surface application prior to transplanting fruiting vegetables or as a post-directed application to transplanted or established direct-seeded fruiting vegetables.

**DO NOT** apply prior to direct-seeded fruiting vegetables. **DO NOT** apply postemergence over the top of or to foliage of fruiting vegetables as severe injury may occur. **Prowl H<sub>2</sub>O** can be applied as a post-directed spray on the soil at the base of the plant beneath plants and between rows. Avoid direct contact with foliage. Be sure roots of transplants are established. Following the post-directed spray, incorporate at the time of blocking and thinning or at "lay-by." **Prowl H<sub>2</sub>O** should be applied prior to weed emergence. **EMERGED WEEDS WILL NOT BE CONTROLLED BY THIS TREATMENT.** Adequate rainfall or overhead irrigation within 7 days after application will provide the most consistent weed control.

Always perform a mixing test to check the compatibility of **Prowl H<sub>2</sub>O** with all potential tank mix partners.

Follow all precautions and restrictions on the labels of all products applied in combination with **Prowl H<sub>2</sub>O**. Always follow the most restrictive label.

#### Broadcast Rate of Prowl H<sub>2</sub>O

Soil Texture	Rate
Coarse*	1.0 to 1.5 pints/A
Medium*	1.5 to 2.0 pints/A
Fine*	1.5 to 3.0 pints/A

\* See the main **Prowl H<sub>2</sub>O** label for soil texture classifications.

### Restrictions and Limitations

**DO NOT** apply more than 3.0 pints per acre per season.

**DO NOT** apply within 70 days of harvest.

**DO NOT** use **Prowl H<sub>2</sub>O** on peat or muck soils.

### Weed Species Controlled

When applied as directed, **Prowl H<sub>2</sub>O** will control the following grasses and broadleaf weed species:

#### GRASSES

- |                          |                                     |
|--------------------------|-------------------------------------|
| Barnyardgrass            | ( <i>Echinochloa crus-galli</i> )   |
| Crabgrass                | ( <i>Digitaria</i> spp.)            |
| Crowfootgrass            | ( <i>Dactyloctenium aegyptium</i> ) |
| Fall panicum             | ( <i>Panicum dichotomiflorum</i> )  |
| Field sandbur            | ( <i>Cenchrus incertus</i> )        |
| Foxtails,                |                                     |
| giant                    | ( <i>Setaria faberi</i> )           |
| green                    | ( <i>Setaria viridis</i> )          |
| yellow                   | ( <i>Setaria glauca</i> )           |
| Goosegrass*              | ( <i>Eleusine indica</i> )          |
| Johnsongrass (from seed) | ( <i>Sorghum halepense</i> )        |
| Signalgrass*             | ( <i>Brachiaria platyphylla</i> )   |
| Texas panicum            | ( <i>Panicum texanum</i> )          |
| Witchgrass               | ( <i>Panicum capillare</i> )        |

#### BROADLEAF WEEDS

- |                |                                 |
|----------------|---------------------------------|
| Carpetweed     | ( <i>Mollugo verticillata</i> ) |
| Florida pusley | ( <i>Richardia scabra</i> )     |
| Kochia         | ( <i>Kochia scoparia</i> )      |
| Lambsquarters  | ( <i>Chenopodium album</i> )    |
| Pigweed        | ( <i>Amaranthus</i> spp.)       |
| Purslane       | ( <i>Portulaca oleracea</i> )   |
| Spurge, annual | ( <i>Euphorbia</i> spp.)        |

\*Suppression Only

6/12

**Conditions of Sale and Warranty**

The **Directions For Use** of this product reflects the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF"), or the Seller. To the extent permitted by law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT PERMITTED BY LAW, BASF OR THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

*Prowl* is a registered trademark of BASF.

© 2006 BASF Corporation.

All rights reserved.

000241-00418.20060517b.NVA 2006-04-195-0176

BASF Corporation  
Agricultural Products  
26 Davis Drive  
Research Triangle Park, NC 27709



The Chemical Company

MAR 30 2007

7/12

# PROWL H<sub>2</sub>O<sup>®</sup>

h e r b i c i d e

## Supplemental Labeling

### FOR USE ON GREEN ONIONS

(INCLUDING Leeks, Spring Onions or Scallions, Japanese Bunching Onions, Green Shallots or Green Eschalots)

#### EPA Reg. No. 241-418

Active Ingredient:

pendimethalin (N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine) ..... 38.7%

**INERT INGREDIENTS:** ..... 61.3%

**TOTAL** ..... 100.0%

OBSERVE ALL PRECAUTIONARY STATEMENTS, MIXING AND APPLICATION INSTRUCTIONS, AND FOLLOW CROP RESTRICTIONS ON THE PROWL<sup>®</sup> H<sub>2</sub>O HERBICIDE MAIN PRODUCT LABEL BEFORE USING

#### Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Prowl H<sub>2</sub>O controls most annual grasses and certain broadleaf weeds as they germinate. Uniformly apply Prowl H<sub>2</sub>O with properly calibrated ground equipment in 20 or more gallons of water. Use sprayers equipped with nozzles that provide accurate and uniform application.

Adequate rainfall or overhead irrigation after application will provide the most consistent weed control.

Always perform a mixing test to check the compatibility of Prowl H<sub>2</sub>O with all potential tank mix partners.

Follow all precautions and restrictions on the labels of all products applied in combination with Prowl H<sub>2</sub>O. Always follow the most restrictive label

#### Non-Target, Endangered Species

If non-target plants occur in proximity to the application site, the following mitigation measures are required:

- If applied by ground, leave untreated buffer zone of 80 feet. The product must be applied using a low boom (20 inches above the ground) and ASAE fine to medium/coarse nozzles.
- If applied by air, leave untreated buffer zone of 200 feet. Must use straight stream nozzles (such as D-6 to D-8). Wind can be no more than 8 mph and release height must be 15 feet or less.

If endangered species occur in proximity to the application site, the following mitigation measures are required:

- If applied by ground, leave untreated buffer zones of 200 feet. The product must be applied using a low boom (20 inches above the ground) and ASAE fine to medium/coarse nozzles.

- If applied by air, leave untreated buffer zone of 1,000 feet. Must use straight stream nozzles (such as D-6 to D-8). Wind can be no more than 8 mph and release height must be 15 feet or less

To determine whether your county has an endangered species, consult the website <http://www.epa.gov/espp/usa-map.htm>.

Endangered Species Bulletins may also be obtained from Extension offices or state pesticide agencies. If the bulletin is not available for your specific area, check with the appropriate local state agency to determine if known populations of endangered species occur in the area to be treated.

**Application Instructions**

Uniformly apply 2.0 pints per acre of **Prowl H<sub>2</sub>O** as a broadcast spray to the soil surface as preemergence spray or as a postemergence spray to the crop at the 2 to 3 true leaf stage at least 30 days before harvest. If **Prowl H<sub>2</sub>O** is to be applied sequentially as both a preemergence and postemergence spray then the preemergence spray must be applied 30 days prior to the postemergence spray.

**Chemigation** – **Prowl H<sub>2</sub>O** may be applied through a sprinkler system including solid set, center pivot, lateral move, end tow, side (wheel) roll, or hand move. Follow all label recommendations regarding rates per acre, timing of application, special instructions, and precautions.

**DO NOT** apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water.

The system must be properly calibrated (with water only) to ensure that the amount of **Prowl H<sub>2</sub>O** applied corresponds to the recommended rate.

Apply **Prowl H<sub>2</sub>O** in ½ to ¾ inches of water during the first sprinkler set. When application is complete, flush the system with water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Refer to the **Prowl H<sub>2</sub>O** main label for further information concerning Special Precautions for all Sprinkler Chemigation Applications.

**Restrictions and Limitations**

**DO NOT** apply more than 2.0 pints per acre per application.

**DO NOT** apply more than 4.0 pints per acre per season.

**DO NOT** apply within 30 days of harvest.

**DO NOT** feed forage or graze livestock in treated fields.

**DO NOT** use **Prowl H<sub>2</sub>O** on peat or muck soils.

Refer to the **Prowl H<sub>2</sub>O** main label for all crop plant-back restrictions.

9/12

### Conditions of Sale and Warranty

The **Directions For Use** of this product reflects the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent permitted by law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT PERMITTED BY LAW, BASF OR THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

*Prowl is a registered trademark of BASF.*

© 2006 BASF Corporation.

All rights reserved.

000241-00418.20060517b.NVA 2006-04-195-0178

BASF Corporation  
Agricultural Products  
26 Davis Drive  
Research Triangle Park, NC 27709

 **BASF**

The Chemical Company

NOTIFICATION

10/12

MAR 30 2007



**FOR USE ON STRAWBERRIES**

EPA Reg. No. 241-418

Active Ingredient:

pendimethalin (N-(1-ethylpropyl)-3,4-dimethyl-2,5-dinitrobenzenamine)..... 38.7%

**INERT INGREDIENTS:** ..... 61.3%

**TOTAL** ..... 100.0%

OBSERVE ALL PRECAUTIONARY STATEMENTS, MIXING AND APPLICATION INSTRUCTIONS, AND FOLLOW CROP RESTRICTIONS ON THE PROWL® H<sub>2</sub>O HERBICIDE MAIN PRODUCT LABEL BEFORE USING.

**Directions For Use**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Prowl H<sub>2</sub>O controls most annual grasses and certain broadleaf weeds as they germinate. Uniformly apply Prowl H<sub>2</sub>O with properly calibrated ground equipment in 10 or more gallons of water. Use sprayers equipped with nozzles that provide accurate and uniform application.

Adequate rainfall or overhead irrigation after application will provide the most consistent weed control.

Always perform a mixing test to check the compatibility of Prowl H<sub>2</sub>O with all potential tank mix partners.

Follow all precautions and restrictions on the labels of all products applied in combination with Prowl H<sub>2</sub>O. Always follow the most restrictive label.

**Non-Target, Endangered Species**

If non-target plants occur in proximity to the application site, the following mitigation measures are required:

- If applied by ground, leave untreated buffer zone of 80 feet. The product must be applied using a low boom (20 inches above the ground) and ASAE fine to medium/coarse nozzles.
- If applied by air, leave untreated buffer zone of 200 feet. Must use straight stream nozzles (such as D-6 to D-8). Wind can be no more than 8 mph and release height must be 15 feet or less.

If endangered species occur in proximity to the application site, the following mitigation measures are required:

- If applied by ground, leave untreated buffer zones of 200 feet. The product must be applied using a low boom (20 inches above the ground) and ASAE fine to medium/coarse nozzles.
- If applied by air, leave untreated buffer zone of 1,000 feet. Must use straight stream nozzles (such as D-6 to D-8). Wind can be no more than 8 mph and release height must be 15 feet or less.

To determine whether your county has an endangered species, consult the website <http://www.epa.gov/espp/usa-map.htm>.

Endangered Species Bulletins may also be obtained from Extension offices or state pesticide agencies. If the bulletin is not available for your specific area, check with the appropriate local state agency to determine if known populations of endangered species occur in the area to be treated.

### Application Instructions

Uniformly apply 3.0 pints per acre of **Prowl H<sub>2</sub>O** as a broadcast spray to the soil surface at pre-transplant time. A second application of 3.0 pints per acre of **Prowl H<sub>2</sub>O** may be applied in a band to the soil between crop rows 35 days before harvest but **DO NOT CONCENTRATE THE RATE** per acre into the treated area. The second application rate is based on per unit of treated area.

**Chemigation** – **Prowl H<sub>2</sub>O** may be applied at pre-transplant to strawberries through a sprinkler system including solid set, center pivot, lateral move, end tow, side (wheel) roll, or hand move. Follow all label recommendations regarding rates per acre, timing of application, special instructions, and precautions.

**DO NOT** apply this product through any other type of irrigation system.

**DO NOT** allow treated irrigation water to contact strawberry plants.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water.

The system must be properly calibrated (with water only) to ensure that the amount of **Prowl H<sub>2</sub>O** applied corresponds to the recommended rate.

Apply **Prowl H<sub>2</sub>O** in ½ to ¾ inches of water during the first sprinkler set. When application is complete, flush the system with water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Refer to the **Prowl H<sub>2</sub>O** main label for further information concerning Special Precautions for all Sprinkler Chemigation Applications.

### Restrictions and Limitations

**DO NOT** apply more than 3.0 pints per acre per application.

**DO NOT** apply more than 6.0 pints per acre per season.

**DO NOT** apply within 35 days of harvest.

**DO NOT** feed forage or graze livestock in treated fields.

**DO NOT** use **Prowl H<sub>2</sub>O** on peat or muck soils.

Refer to the **Prowl H<sub>2</sub>O** main label for all crop plant-back restrictions.

12/12

### Conditions of Sale and Warranty

The **Directions For Use** of this product reflects the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent permitted by law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT PERMITTED BY LAW, BASF OR THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF

*Prowl* is a registered trademark of BASF.

© 2006 BASF Corporation.

All rights reserved.

000241-00418.20060517b.NVA 2006-04-195-0179

BASF Corporation  
Agricultural Products  
26 Davis Drive  
Research Triangle Park, NC 27709

 **BASF**

The Chemical Company