

241-418

08/28/2012

1/13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Mr. Craig Kleppe  
BASF Corporation  
26 Davis Dr., PO Box 13528  
Research Triangle Park, NC 27709

AUG 28 2012

Subject: Prowl H2O Herbicide  
EPA Registration Numbers 241-418  
IR-4 Project application dated May 20, 2011  
Label resubmission by email dated 8/21/2012  
New Uses on Lettuce, leaf; Brassica, leafy greens, subgroup 5B; Turnip greens;  
Melon subgroup 9A; Vegetable, soybean, succulent; Small fruit vine climbing  
subgroup 13-07E, except grape  
Supplemental labels: For use on leaf lettuce, For use on leafy Brassica greens, For  
use between rows in melon production, For use on vegetable soybean (edamame),  
For use on kiwi and other small fruit vines

Dear Mr. Kleppe,

The supplemental labeling (5) referred to above and submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, are acceptable. They expire on September 1, 2015. At your next label printing, or within eighteen months of the date of this letter, whichever comes first, you must incorporate the supplemental labeling into the main product labeling.


A stamped copy of labeling is enclosed for your records. Products released for shipment after eighteen (18) months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions. If you have any questions, please contact Hope Johnson at 703-305-5410.

Sincerely,

A handwritten signature in black ink, appearing to read "Kable Bo Davis", written over a horizontal line.

Kable Bo Davis  
Product Manager 25  
Herbicide Branch  
Registration Division (7505P)

ACCEPTED  
AUG 28 2012  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-418



**PROWL<sup>®</sup> H<sub>2</sub>O**  
herbicide

**For use between rows in melon production**

This supplemental label expires September 1, 2015, and must not be used or distributed after this date.

<b>Active Ingredient*:</b>	
pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine .....	38.7%
<b>Other Ingredients:</b> .....	61.3%
<b>Total:</b> .....	100.0%

\*1 gallon contains 3.8 pounds of pendimethalin formulated as an aqueous capsule suspension.

**EPA Reg. No. 241-418**

**Directions For Use**

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Prowl<sup>®</sup> H<sub>2</sub>O herbicide** container label, EPA Reg. No. 241-418, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Prowl H<sub>2</sub>O** before applying.
- Use of **Prowl H<sub>2</sub>O** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Prowl H<sub>2</sub>O**.

**Product Information**

**Prowl H<sub>2</sub>O** is a selective herbicide for controlling most annual grasses and certain broadleaf weeds as they germinate. Refer to the **Prowl H<sub>2</sub>O** container label for a complete list of weeds controlled.

**Prowl H<sub>2</sub>O** may be applied in the following melons: cantaloupe, citron melon, muskmelon, watermelon.

**Application Instructions**

**Prowl H<sub>2</sub>O** may be applied only by ground.

**Use Methods and Timings**

**Prowl H<sub>2</sub>O** may be applied sequentially in melon production. Initially apply up to 2.1 pints per acre of **Prowl H<sub>2</sub>O** as a shielded application to row middles (either before melon transplanting or before a seeded crop has emerged) or between rows covered with plastic mulch (prior to holes being punched in plastic for melon planting). Make a second shielded application at up to 2.1 pints per acre of **Prowl H<sub>2</sub>O** to row middles or between plastic mulch prior to melon vine running. The interval between the sequential **Prowl H<sub>2</sub>O** applications must be at least 21 days. Avoid spray contact with melon foliage or running vines because crop injury could occur.

Apply **Prowl H<sub>2</sub>O** prior to weed germination. Emerged weeds will not be controlled by these treatments.

**Restrictions and Limitations**

- **DO NOT** apply more than 2.1 pints per acre in a single application or more than 4.2 pints per acre per season.
- **DO NOT** apply within 35 days before melon harvest.
- **DO NOT** feed forage or graze livestock in treated fields.

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Prowl is a registered trademark of BASF.*

©2012 BASF Corporation  
All rights reserved.


000241-00418.20110308b.NVA 2011-04-195-0058

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



The Chemical Company

413



## Supplemental Label

# PROWL<sup>H<sub>2</sub>O</sup>

herbicide

### For use on leaf lettuce

This supplemental label expires September 1, 2015, and must not be used or distributed after this date.

#### Active Ingredient\*

pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine ..... 38.7%

Other Ingredients: ..... 61.3%

Total: ..... 100.0%

\*1 gallon contains 3.8 pounds of pendimethalin formulated as an aqueous capsule suspension.

EPA Reg. No. 241-418

### Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Prowl<sup>H<sub>2</sub>O</sup> herbicide** container label, EPA Reg. No. 241-418, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Prowl H<sub>2</sub>O** before applying.
- Use of **Prowl H<sub>2</sub>O** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Prowl H<sub>2</sub>O**.

### Product Information

**Prowl H<sub>2</sub>O** is a selective herbicide for controlling most annual grasses and certain broadleaf weeds as they germinate. Refer to the **Prowl H<sub>2</sub>O** container label for a complete list of weeds controlled.

### Application Instructions

**Prowl H<sub>2</sub>O** may be applied by ground or air.

#### Use Methods and Timings

With a single application, uniformly apply up to 2.1 pints per acre of **Prowl H<sub>2</sub>O** as a broadcast foliar spray to either direct-seeded or transplanted leaf lettuce from the 3-leaf stage until 20 days prior to harvest.

**Prowl H<sub>2</sub>O** should be applied prior to weed germination. Emerged weeds will not be controlled by this treatment.

**Prowl H<sub>2</sub>O** may be used on peat and muck soils, but weed control may be inconsistent and/or reduced. Use maximum labeled use rate allowed in leaf lettuce.

### Restrictions and Limitations

- **DO NOT** apply preplant or pretransplant or preemergence (direct-seeded) to leaf lettuce because severe injury may occur.
- **DO NOT** foliar apply to leaf lettuce before the 3-leaf growth stage because severe injury may occur.
- **DO NOT** apply more than 2.1 pints per acre per season.
- **DO NOT** apply within 20 days before leaf harvest.

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Prowl is a registered trademark of BASF.*

©2012 BASF Corporation  
All rights reserved.


000241-00418.20110307b.NVA 2011-04-195-0056

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



The Chemical Company

6/13



## Supplemental Label

# PROWL H<sub>2</sub>O

herbicide

### For use on leafy Brassica greens

This supplemental label expires September 1, 2015, and must not be used or distributed after this date.

#### Active Ingredient\*

pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine ..... 38.7%

Other Ingredients: ..... 61.3%

Total: ..... 100.0%

\*1 gallon contains 3.8 pounds of pendimethalin formulated as an aqueous capsule suspension.

EPA Reg. No. 241-418

### Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Prowl H<sub>2</sub>O herbicide** container label, EPA Reg. No. 241-418, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Prowl H<sub>2</sub>O** before applying.
- Use of **Prowl H<sub>2</sub>O** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Prowl H<sub>2</sub>O**.

### Product Information

**Prowl H<sub>2</sub>O** is a selective herbicide for controlling most annual grasses and certain broadleaf weeds as they germinate. Refer to the **Prowl H<sub>2</sub>O** container label for a complete list of weeds controlled.

**Prowl H<sub>2</sub>O** may be applied in the following leafy Brassica greens:  
mustard greens, broccoli raab, Chinese cabbage (bok choy), collards, kale, mizuna, mustard spinach, rape greens and turnip greens (cultivars or varieties grown for leaves only).

### Application Instructions

**Prowl H<sub>2</sub>O** may be applied by ground or air.

### Use Methods and Timings

With a single application, uniformly apply up to 2.1 pints per acre of **Prowl H<sub>2</sub>O** as a broadcast foliar spray to either direct-seeded or transplanted leafy Brassica greens at the 4-leaf to 5-leaf stage.

**Prowl H<sub>2</sub>O** should be applied prior to weed germination. Emerged weeds will not be controlled by this treatment.

**Prowl H<sub>2</sub>O** may be used on peat and muck soils, but weed control may be inconsistent and/or reduced. Use maximum labeled use rate allowed in leafy Brassica greens.

### Restrictions and Limitations

- **DO NOT** apply preplant or pretransplant or preemergence (direct-seeded) to leafy Brassica greens because severe injury may occur.
- **DO NOT** foliar apply to leafy Brassica greens before the 4-leaf growth stage because severe injury may occur.
- **DO NOT** apply to turnip greens varieties grown for roots or to dual-purpose varieties grown for roots and tops.
- **DO NOT** use **Prowl H<sub>2</sub>O**-treated turnip greens roots for any feed or food purpose.
- **DO NOT** apply more than 2.1 pints per acre per season.
- **DO NOT** apply within 21 days before leafy greens harvest.

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Prowl is a registered trademark of BASF.*

©2012 BASF Corporation  
All rights reserved.


000241-00418.20110307b.NVA 2011-04-195-0057

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



The Chemical Company

AUG 28 2012  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act,  
 as amended, for the pesticide  
 registered under  
 EPA Reg. No. 241-418



**PROWL H<sub>2</sub>O**  
 herbicide

## For use on vegetable soybean (edamame)

This supplemental label expires September 1, 2015, and must not be used or distributed after this date.

### Active Ingredient\*:

pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine ..... 38.7%

Other Ingredients: ..... 61.3%

Total: ..... 100.0%

\*1 gallon contains 3.8 pounds of pendimethalin formulated as an aqueous capsule suspension.

EPA Reg. No. 241-418

### Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Prowl H<sub>2</sub>O herbicide** container label, EPA Reg. No. 241-418, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Prowl H<sub>2</sub>O** before applying.
- Use of **Prowl H<sub>2</sub>O** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Prowl H<sub>2</sub>O**.

### Product Information

**Prowl H<sub>2</sub>O** is a selective herbicide for controlling most annual grasses and certain broadleaf weeds as they germinate. Refer to the **Prowl H<sub>2</sub>O** container label for a complete list of weeds controlled:

### Application Instructions

**Prowl H<sub>2</sub>O** may only be applied by ground.

### Use Methods, Timings, and Rates

**Prowl H<sub>2</sub>O** may be applied in conventional, minimum, or no-till as a preplant surface, preplant incorporated, or preemergence application in vegetable soybean (edamame).

### Preplant Surface

Apply **Prowl H<sub>2</sub>O** up to 15 days prior to planting. **Prowl H<sub>2</sub>O** may be applied up to 45 days prior to planting when used in a tank mix or applied sequentially with postemergence-applied herbicides registered for use in vegetable soybean (edamame).

### Preplant Incorporated

Apply **Prowl H<sub>2</sub>O** up to 60 days prior to planting and incorporate.

### Preemergence

Apply **Prowl H<sub>2</sub>O** at planting or up to 2 days after planting. Apply to a firm seedbed, free of clods. **DO NOT** make applications of **Prowl H<sub>2</sub>O** preemergence north of Interstate 80, except in states of Indiana, Michigan and Ohio.



## Use Rates

### Preplant Surface or Preplant Incorporated

Soil Texture	<3% Organic Matter >3% (pts/A)	
Coarse	1.5	2.0
Medium	2.5*	3.0
Fine**	3.0	3.0
<p>* DO NOT exceed 2.1 pts for Southern states; see <b>Restrictions and Limitations</b> in the <b>Prowl® H<sub>2</sub>O herbicide</b> main container label for map of specific states.</p> <p>** For heavy clay soils, apply <b>Prowl H<sub>2</sub>O</b> at the broadcast rate of 3.2 pints per acre.</p>		

### Preemergence Applications

Soil Texture	<3% Organic Matter >3% (pts/A)	
Coarse	1.5	1.5
Medium	2.0	2.0
Fine	2.0	2.5

### Restrictions and Limitations

- Livestock can graze or be fed forage from treated vegetable soybean (edamame) fields.
- **DO NOT** apply within 85 days of harvest.
- **DO NOT** exceed one application per crop season at the highest rate per acre for any given soil type and application method.
- **DO NOT** use in California.

10/13

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Prowl is a registered trademark of BASF.*

©2012 BASF Corporation  
All rights reserved.

000241-00418.20110308b.NVA 2011-04-195-0054


BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



The Chemical Company

11/13

**ACCEPTED**  
AUG 28 2012  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticides registered under EPA Reg. No. **241-418**



**PROWL<sup>®</sup> H<sub>2</sub>O**  
herbicide

## For use on kiwi and other small fruit vines

This supplemental label expires September 1, 2015, and must not be used or distributed after this date.

### Active Ingredient\*:

pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine ..... 38.7%

Other Ingredients: ..... 61.3%

Total: ..... 100.0%

\*1 gallon contains 3.8 pounds of pendimethalin formulated as an aqueous capsule suspension.

**EPA Reg. No. 241-418**

### Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Prowl<sup>®</sup> H<sub>2</sub>O herbicide** container label, EPA Reg. No. 241-418, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Prowl H<sub>2</sub>O** before applying.
- Use of **Prowl H<sub>2</sub>O** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Prowl H<sub>2</sub>O**.

### Product Information

**Prowl H<sub>2</sub>O** is a selective herbicide for controlling most annual grasses and certain broadleaf weeds as they germinate. Refer to the **Prowl H<sub>2</sub>O** container label for a complete list of weeds controlled.

**Prowl H<sub>2</sub>O** may be applied in the following bearing small fruit vines climbing:  
Amur river grape, gooseberry, fuzzy kiwifruit, hardy kiwifruit, maypop, and Schisandra berry.

### Application Instructions

**Prowl H<sub>2</sub>O** may be only applied by ground, chemigation, or flood, flooded basin and gravity flow irrigation systems.

BASF Corporation  
26 Davis Drive, Research Triangle Park, NC 27709

### Use Methods and Timings

**Prowl H<sub>2</sub>O** may be applied either in a single application or sequentially with an interval of 30 days or more. Uniformly apply **Prowl H<sub>2</sub>O** in small fruit vines up to 4.2 quarts per acre depending on the grower's weed control program, level of weed infestation, and desired use strategy (see chart following).

#### Prowl H<sub>2</sub>O Use Rate per Acre

Low Use Rate	3.2 quarts
High Use Rate	4.2 quarts

**Prowl H<sub>2</sub>O** may be applied anytime after fall harvest, during winter dormancy, and in the spring.

### Ground Applications

**Prowl H<sub>2</sub>O** may be applied **surface incorporated or (surface) preemergence**.

Apply **Prowl H<sub>2</sub>O** as a broadcast or banded treatment using ground equipment before weed germination. Apply the spray directly to the ground beneath the small fruit vines and/or in areas between rows. **DO NOT** apply over the top of small fruit vines with leaves or buds or fruit. Contact with leaves, shoots, or buds by the spray mixture may cause injury.

 **BASF**  
The Chemical Company

## Chemigation Applications

**Prowl® H<sub>2</sub>O herbicide** may be applied through sprinkler irrigation and drip irrigation systems. Follow all recommendations, special instructions and precautions about chemigation in the **Spraying Instructions** section of the **Prowl H<sub>2</sub>O** container label. **DO NOT** apply **Prowl H<sub>2</sub>O**-treated irrigation water over the top of small fruit vines with leaves, or buds, or fruit.

## Flood, Flooded Basin, and Gravity Flow Irrigation Systems

**Prowl H<sub>2</sub>O** may be applied in flood, flooded basin, and gravity flow irrigation systems. Follow all recommendations, special instructions and precautions about flood, flooded basin, and gravity flow irrigation systems in the **Spraying Instructions** section of the **Prowl H<sub>2</sub>O** container label.

---

## Restrictions and Limitations

---

- **DO NOT** apply over the top of small fruit vines with leaves, or buds, or fruit.
- **DO NOT** apply by air.
- **DO NOT** apply more than 4.2 quarts per acre per year (a single growing season).
- **DO NOT** apply within 60 days before harvest of fruit.
- **DO NOT** feed forage or graze livestock in treated vines.
- **DO NOT** apply **Prowl H<sub>2</sub>O** when impregnated onto dry bulk fertilizer in small fruit climbing vines.

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Prowl is a registered trademark of BASF.*

©2012 BASF Corporation  
All rights reserved.

000241-00418.20110307b.NVA 2011-04-195-0055

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



The Chemical Company