



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

JAN - 7 2008

Jeffrey H. Birk, Ph.D.
Regulatory Manager
BASF
26 Davis Drive
Research Triangle Park, NC 27709

RE: Notification of Minor Changes for Arsenal Powerline Herbicide
EPA Registration Number: 241-431
Date of Submission: November 6, 2007

Dear Dr. Birk:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated November 6, 2007, for the product, Arsenal Powerline Herbicide. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

Linda Arrington
for Linda Arrington

Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
 Washington, DC 20460

Registration
 Amendment
 Other

**Registration
 Amendment
 Other**

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 241-431	2. EPA Product Manager James Tompkins	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Arsenal PowerLine herbicide	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) BASF 26 Davis Drive Research Triangle Park, NC 27709 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION
 JAN - 7 2008

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of minor label changes for Arsenal PowerLine herbicide (241-431). This notification is consistent with the provisions of PR Notice 95-2 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

notification is not subject to a fee under PRIA. Contact Jeff Birk at 919-547-2622 (phone), 919-547-2850 (fax) or by Email at jeffrey.birk@basf.com

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Unit Packaging wgt. No. per container	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Package wgt No. per container	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
--	---	--	--

*** Certification must be submitted**

3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> _____
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph Paper glued Stenciled <input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Jeffrey H. Birk	Title Regulatory Manager	Telephone No. (Include Area Code) 919-547-2622
-------------------------	-----------------------------	---

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature 	3. Title Regulatory Manager	6. Date Application Received (Stamped)
4. Typed Name Jeffrey H. Birk	5. Date November 6, 2007	

Certification with Respect to Label Integrity

version: 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

PROPOSED LABEL		
EPA Registration #	Date Submitted to EPA	Electronic file name
241-431	11-6-07	000241-00431.20071106.NVA 2007-04-295-0100

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.



 Signature

11/06/2007

 Date

Jeffrey H. Birk

 Name (typed)

Regulatory Manager

 Title

NOTIFICATION
JAN - 7 2008

4/21

NOTIFICATION

JAN - 7 2008

ARSENAL®

POWERline™

herbicide

For the control of undesirable vegetation in grass pasture, rangeland and industrial noncropland areas including railroad, utility, pipeline and highway rights-of-way, utility plant sites, petroleum tank farms, pumping installations, storage areas, fence rows, nonirrigation ditchbanks and for the establishment and maintenance of wildlife openings.

Active Ingredient:

Isopropylamine salt of imazapyr(2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid)* 26.7%

Other Ingredients: 73.3%

Total: 100.0%

*Equivalent to 21.8% 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid or 2-propanoic acid (not active).

EPA Reg. No. 241-431

U.S. Patent No. 4,798,619

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN. CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call day or night, 1-800-832-HELP (4357).

Net Contents:

400-1100000000
400-1100000000
400-1100000000
400-1100000000

BASF

The Chemical Company

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to by a poison control center or doctor. • DO NOT give anything to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).</p>	

PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS
CAUTION**

Harmful if swallowed, causes moderate eye irritation. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for **Category A** on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves made of any waterproof material.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:
<p>Users Should:</p> <ul style="list-style-type: none"> • Wash hands before eating, chewing gum, using tobacco or using the toilet. • Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

PHYSICAL AND CHEMICAL HAZARDS

Spray solutions of **Arsenal® PowerLine™ herbicide** should be mixed, stored and applied only in stainless steel, fiberglass, plastic and plastic-lined steel containers.

DO NOT mix, store or apply **Arsenal PowerLine** or spray solutions of **Arsenal PowerLine** in unlined steel (except stainless steel) containers or spray tanks.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Arsenal PowerLine may be used only in accordance with instructions on the leaflet label attached to the container. Keep containers closed to avoid spills and contamination.

4/21

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **48 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Protective eyewear
- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Noncrop weed control is not within the scope of the Worker Protection Standard. See the **GENERAL INFORMATION** section of this label for a description of noncrop sites.

DO NOT enter or allow others to enter treated areas until sprays have dried.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: DO NOT store below 10° F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL FOR 2.5 GALLON AND

30 GALLON: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in an approved sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR FIELD KEG, MINIBULK

AND BULK: When this container is empty, replace the cap and seal all openings that have been opened during use, and return the container to the point of purchase, or to a designated location. This container must only be refilled with the pesticide product. **DO NOT** reuse the container for any other purpose. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. **DO NOT** transport if this container is damaged or leaking. If the container is damaged or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling. Disposal of container must be in compliance with state and local regulations.

IMPORTANT

DO NOT use on food crops. Keep from contact with fertilizers, insecticides, fungicides and seeds. **DO NOT** drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. **DO NOT** use on lawns, walks, driveways, tennis courts, or similar areas where roots of desirable vegetation may extend and be exposed to potential injury and/or mortality from root uptake of **Arsenal® PowerLine™ herbicide** unless this risk is acceptable. **DO NOT** side trim desirable vegetation with this product unless severe injury or plant death can be tolerated. Prevent drift of spray to desirable plants.

Clean application equipment after using this product by thoroughly flushing with water.

GENERAL INFORMATION

Use Sites: Arsenal PowerLine is an aqueous solution to be mixed with water and a surfactant and applied as a spray solution to grass pasture and rangeland and non-cropland areas such as railroad, utility, and pipeline rights-of-way; utility plant sites, petroleum tank farms, pumping installations, storage areas, highway rights-of-way, fence rows, and nonirrigation ditchbanks including grazed or hayed areas within these sites. **Arsenal PowerLine** is recommended for the establishment and maintenance of wildlife openings. **Arsenal PowerLine** may also be used for the release of unimproved bermudagrass (see specific directions) and for use under certain paved surfaces (see specific directions).

Application Methods: Arsenal PowerLine will control most annual and perennial grasses and broadleaf weeds in addition to many brush and vine species, and **Arsenal PowerLine** will provide residual control of labeled weeds which germinate in the treated areas. This product may be applied either pre-emergence or postemergence to the weeds; however, postemergence application is the method of choice in most situations, particularly for perennial species. For maximum activity, weeds should be growing vigorously at the time of postemergence applica-

tion, and the spray solution should include a surfactant (see **ADJUVANT** section for specific recommendations). These solutions may be applied selectively by using low-volume techniques or may be applied broadcast by using ground equipment or aerial equipment. In addition, **Arsenal® PowerLine™ herbicide** may also be used for stump and cut stem treatments (see specific directions).

Herbicidal Activity: **Arsenal PowerLine** is readily absorbed through leaves, stems, and roots and is translocated rapidly throughout the plant, with accumulation in the meristematic regions. Treated plants stop growing soon after spray application. Chlorosis appears first in the newest leaves, and necrosis spreads from this point. In perennials, the herbicide is translocated into, and kills, underground storage organs which prevents regrowth. Chlorosis and tissue necrosis may not be apparent in some plant species until two weeks after application. Complete kill of plants may not occur for several weeks. Applications of **Arsenal PowerLine** are rainfast one hour after treatment.

PRECAUTIONS FOR AVOIDING INJURY TO NONTARGET PLANTS

Untreated trees can occasionally be affected by root uptake of **Arsenal PowerLine** through movement into the top soil. Injury or loss of desirable trees or other plants may result if **Arsenal PowerLine** is applied on or near desirable trees or other plants, on areas where their roots extend, or in locations where the treated soil may be washed or moved into contact with their roots.

SPRAY DRIFT REQUIREMENTS

Aerial Applications

- Applicators are required to use a coarse or coarser droplet size (ASABE S572) or, if specifically using a spinning atomizer nozzle, applicators are required to use a volume mean diameter (VMD) of 385 microns or greater for release heights below 10 feet. Applicators are required to use a very coarse or coarser droplet size or, if specifically using a spinning atomizer nozzle, applicators are required to use a VMD of 475 microns or greater for release heights above 10 feet. Applicators must consider the effects of nozzle orientation and flight speed when determining droplet size.
- Applicators are required to use upwind swath displacement.
- The boom length must not exceed 60% of the wingspan or 90% of the rotor blade diameter to reduce spray drift.
- Applications with wind speeds less than 3 mph and with wind speeds greater than 10 mph are prohibited.
- Applications into temperature inversions are prohibited.

Ground Boom Applications

- Applicators are required to use a nozzle height below 4 feet above the ground or plant canopy and coarse or coarser droplet size (ASABE S572) or, if specifically using a spinning atomizer nozzle, applicators are required to use a volume mean diameter (VMD) of 385 microns or greater.

- Applications with wind speeds greater than 10 mph are prohibited.
- Applications into temperature inversions are prohibited.

Wind Erosion: Avoid treating powdery, dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.

Aerial Application Methods and Equipment: Use 2 or more gallons of water per acre. The actual minimum spray volume per acre is determined by the spray equipment used. Use adequate spray volume to provide accurate and uniform distribution of spray particles over the treated area and to avoid spray drift.

Ground Application (Broadcast): Use 5 or more gallons of water per acre. The actual minimum spray volume per acre is determined by the spray equipment used. Use adequate spray volume to provide accurate and uniform distribution of spray particles over the treated area and to avoid spray drift.

ADJUVANTS

Postemergence applications of Arsenal PowerLine require the addition of a spray adjuvant for optimum herbicide performance.

Nonionic Surfactants: Use a nonionic surfactant at the rate 0.25% v/v or higher (see manufacturer's label) of the spray solution (0.25% v/v is equivalent to 1 quart in 100 gallons). For best results, select a nonionic surfactant with an HLB (hydrophilic to lipophilic balance) ratio between 12 and 17 with at least 70% surfactant in the formulated product (alcohols, fatty acids, oils, ethylene glycol or diethylene glycol should not be considered as surfactants to meet the above requirements).

Methylated Seed Oils or Vegetable Oil Concentrates: Instead of a surfactant, a methylated seed oil or vegetable-based seed oil concentrate may be used at the rate of 1.5 to 2 pints per acre. When using spray volumes greater than 30 gallons per acre, methylated seed oil or vegetable-based seed oil concentrates should be mixed at a rate of 1% of the total spray volume, or alternatively use a nonionic surfactant as described above. Research indicates that these oils may aid in **Arsenal PowerLine** deposition and uptake by plants under moisture or temperature stress.

Silicone-Based Surfactants: See manufacturer's label for specific rate recommendations. Silicone-based surfactants may reduce the surface tension of the spray droplet allowing greater spreading on the leaf surface as compared to conventional nonionic surfactants. However, some silicone-based surfactants may dry too quickly, limiting herbicide uptake.

Fertilizer/Surfactant Blends: Nitrogen-based liquid fertilizers, such as 28%N, 32%N, 10-34-0 or ammonium sulfate, may be added at the rate of 2 to 3 pints per acre in combination with the recommended rate of nonionic surfactant, methylated seed oil or vegetable/seed oil concen-

trate. The use of fertilizers in a tank mix without a nonionic surfactant, methylated seed oil or vegetable/seed oil concentrate is not recommended.

BRUSH CONTROL

AERIAL APPLICATIONS:

All precautions should be taken to minimize or eliminate spray drift. Fixed wing aircraft and helicopters can be used to apply **Arsenal® PowerLine™ herbicide**. However, **DO NOT** make applications by fixed wing aircraft unless appropriate buffer zones can be maintained to prevent spray drift out of the target area or, when treating open tracts of land, spray drift as a result of fixed wing aircraft application can be tolerated. Aerial equipment designed to minimize spray drift, such as a helicopter equipped with a **Microfoil™ boom, Thru-Valve™ boom** or raindrop nozzles, must be used and calibrated. Except when applying with a **Microfoil boom**, a drift control agent may be added at the recommended label rate. To avoid drift, applications should not be made during inversion conditions, when winds are gusty, or any other conditions which allow drift. Side trimming is not recommended with **Arsenal PowerLine** unless death of treated tree can be tolerated.

Uniformly apply the specified amount of **Arsenal PowerLine** in 5 to 30 gallons of water per acre; include in the spray solution a nonionic surfactant or methylated seed oil or manufacturer's label rate of a silicone-based surfactant (see the **ADJUVANT** section of this label for specific recommendations). A foam-reducing agent may be added at the recommended label rate, if needed.

IMPORTANT: Thoroughly clean application equipment, including landing gear, immediately after use of this product. Prolonged exposure of this product to uncoated steel (except stainless steel) surfaces may result in corrosion and failure of the exposed part. The maintenance of an organic coating (paint) may prevent corrosion.

GROUND APPLICATIONS:

IMPORTANT: To minimize spray drift, select proper nozzles to avoid spraying a fine mist. Use pressures less than 50 psi, and **DO NOT** spray under gusty or windy conditions. Add a foam-reducing agent, if needed, and a spray pattern indicator, if desired, at the recommended label rates. Clean application equipment after using this product by thoroughly flushing with water.

When making applications to rights-of-way corridors where desirable tree roots may extend, use 1 to 3 pints of **Arsenal PowerLine** per acre in combination with recommended tank mixes. Use rates higher than 3 pints per acre in these situations may cause injury or death of desirable trees when their roots extend into treated zones.

Side Trimming:

DO NOT side trim with **Arsenal PowerLine** unless severe injury or death of the treated tree can be tolerated. **Arsenal PowerLine** is readily translocated and can result in death of the entire tree.

Low Volume:

Use equipment calibrated to deliver 5 to 20 gallons of spray solution per acre. To prepare the spray solution, thoroughly mix in water 0.5 to 5% **Arsenal PowerLine** plus surfactant (see the **ADJUVANT** section of this label for specific recommendations). A foam-reducing agent may be applied at the recommended label rate, if needed. For control of difficult brush species (see **WEEDS CONTROLLED** section for relative susceptibility of weed species), use the higher concentrations of herbicide and/or spray volumes, but **DO NOT** apply more than 6 pints of **Arsenal PowerLine** per acre. Excessive wetting of foliage is not recommended. See the mixing guide below for some suggested volumes of **Arsenal PowerLine** and water.

9/24

TANK MIXES AND APPLICATION RATES*

Target Vegetation	Rate of Arsenal® PowerLine™ herbicide	Tank Mix
Mixed hardwoods without elm, locust, or pine	1.0 - 1.5% by volume	Surfactant
Mixed hardwoods containing elm, locust, and pine	0.5 - 1.0% by volume	Accord® at 2 - 3% by volume plus surfactant
Mixed hardwoods with locust and pine but no elm	0.5 - 1.0% by volume	Krenite® at 2 - 5% by volume plus surfactant
Mixed hardwoods with locust and elm but no pine	0.5 - 1.0% by volume	Escort® at 2 oz/Acre or 2.3 grams/gallon plus surfactant

*Tank mixes with 2,4-D or products containing 2,4-D have resulted in reduced efficacy of **Arsenal PowerLine**.

MIXING CHART

% Solution	Amount Arsenal PowerLine per Gallon of Mix (oz)	Amount Arsenal PowerLine per 4-Gallon Backpack (oz)
0.5%	0.6	2.6
1.0%	1.3	5.1
2.0%	2.6	10.2
3.0%	3.8	15.4
5.0%	6.4	25.6

MEASURING CHART

128 ounces	=	1 gallon
16 ounces	=	1 pint
8 pints	=	1 gallon
4 quarts	=	1 gallon
2 pints	=	1 quart

10/21

Application Instructions: For low volume, select proper nozzles so that herbicide is not overapplied. Best results are achieved when the spray covers the crown and approximately 70 percent of the plant. The use of an even flat fan tip with a spray angle of 40 degrees or less will aid in proper deposition.

Recommended tip sizes include 4004E or 1504E. For a straight stream and cone pattern, adjustable cone nozzles, such as 5500 X3 or 5500 X4, may be used. Attaching a roll-over valve onto a Spraying Systems Model 30 gunjet or other similar spray guns allows for the use of both a flat fan and cone tips on the same gun.

Proper Spray Pattern: Moisten but **DO NOT** drench target vegetation causing spray solution to run off.

Low Volume with Backpacks:

For brush up to 4-feet tall, spray down on the crown covering crown and penetrating approximately 70% of the plant.

For brush 4- to 8-feet tall, swipe the sides of target vegetation by directing spray to at least two sides of the plant in smooth vertical motions from the crown to the bottom. Make sure to cover the crown whenever possible.

For brush over 8-feet tall, lace sides of the brush by directing spray to at least two sides of the target in smooth zigzag motions from crown to bottom.

Low Volume with Hydraulic Handgun Application Equipment:

Use same technique as described above for **Low Volume with Backpacks**.

For broadcast applications, simulate a gentle rain near the top of target vegetation allowing spray to contact the crown and penetrate the target foliage without falling to the understory. Herbicide spray solution which contacts the understory may result in severe injury or death of plants in the understory.

SPRAY SOLUTION MIXING GUIDE FOR LOW-VOLUME APPLICATIONS

Amount Of Spray Solution Being Prepared (gallons)	Desired Concentration (fluid volume)				
	0.5%	0.75%	1%	1.5%	5%
	(amount of Arsenal® PowerLine™ herbicide to use)				
1	0.6 oz	0.9 oz	1.3 oz	1.9 oz	6.5 oz
3	1.9 oz	2.8 oz	3.8 oz	5.8 oz	1.2 pints
4	2.5 oz	3.8 oz	5.1 oz	7.7 oz	1.6 pints
5	3.2 oz	4.8 oz	6.5 oz	9.6 oz	2 pints
50	2 pints	3 pints	4 pints	6 pints	10 quarts
100	4 pints	6 pints	8 pints	6 quarts	5 gallons

2 tablespoons = 1 fluid ounce

High Volumes:

For optimum performance when spraying medium- to high-density brush, use equipment calibrated to deliver up to 100 gallons of spray solution per acre (GPA). Spray solutions exceeding 100 GPA may result in excessive spray run-off causing increased ground cover injury and injury to desirable species. To prepare the spray solution, thoroughly mix **Arsenal® PowerLine™ herbicide** at a rate of 2 to 6 pints per acre (see **IMPORTANT** section under **GROUND APPLICATIONS**) in water and add a surfactant (see **ADJUVANT** section for specific recommendations and rates of surfactants). A foam-reducing agent may be added at the recommended label rate, if needed. For control of difficult species (see **WEEDS CONTROLLED** section for relative susceptibility of weed species), use the higher concentrations of herbicide and/or spray volumes, but **DO NOT** apply more than 6 pints of **Arsenal PowerLine** per acre. Uniformly cover the foliage of the vegetation to be controlled, but **DO NOT** apply to run-off. Excessive wetting of foliage is not recommended.

TANK MIXES FOR BRUSH CONTROL:

Arsenal PowerLine may be tank mixed with **Accord®, Roundup®, Krenite®, Escort®, Telar®, Tordon® K, Garlon® 3A, Banvel® and Vanquish®** to provide control of **Arsenal PowerLine**-tolerant species.

Consult manufacturer's labels for specific rates and weeds controlled. Always follow the more restrictive label when making an application involving tank mixes. Tank mixing with 2,4-D or products which contain 2,4-D has resulted in reduced performance of **Arsenal PowerLine**.

INVERT EMULSIONS:

Arsenal PowerLine can be applied as an invert emulsion. The spray solution results in an invert (water-in-oil) spray emulsion designed to minimize spray drift and spray run-off resulting in more herbicide on the target foliage. The spray emulsion may be formed in a single tank (batch mixing) or injected (in-line mixing). Consult the invert chemical label for proper mixing directions.

CUT STUBBLE:

Arsenal PowerLine can be applied within 2 weeks after mechanical mowing or cutting of brush. To suppress or control resprouting, uniformly apply a spray solution of **Arsenal PowerLine** at the rate of 1 to 2 pints per acre to the cut area. **Arsenal PowerLine** may be tank mixed with **Tordon K** or picloram to aid in control or suppression of brush. The addition of 5% (v/v) or more of a penetrating agent can aid in uptake through the bark or exposed roots.

Cut stubble applications are made to the soil and cut brush stumps. This type of application may increase ground cover injury. However, vegetation will recover. Making applications of **Arsenal PowerLine** directly to the soil can increase potential root uptake causing injury or death of desirable trees.

Efficacy can be increased, and root uptake by desirable vegetation can be decreased, if the brush is allowed to regrow and the foliage is treated. See the **BRUSH CONTROLLED** section of this label.

STUMP AND CUT STEM TREATMENTS:

Arsenal PowerLine may be used to control undesirable woody vegetation on noncropland by applying the **Arsenal PowerLine** solution to the cambium area of freshly cut stump surfaces or to fresh cuts on the stem of the target woody vegetation. Applications can be made at any time of the year except during periods of heavy sap flow in the spring. **DO NOT** over apply solution causing runoff or puddling.

Mixing: Arsenal PowerLine may be mixed as either a concentrated or dilute solution for stump and cut stem treatments. The dilute solution may be used for applications to the surface of the stump or to cuts on the stem of the target woody vegetation. Concentrated solutions may be used for applications to cuts on the stem. Use of the concentrated solution permits application to fewer cuts on the stem, especially for large-diameter trees. Follow the application instructions to determine proper application techniques for each type of solution.

To prepare a dilute solution, mix 8 to 12 fluid ounces of **Arsenal PowerLine** with one gallon of water. If temperatures are such that freezing of the spray mixture may occur, antifreeze (ethylene glycol) may be used according to manufacturer's label to prevent freezing. The use of a surfactant or penetrating agent may improve uptake through partially callused cambiums. To prepare a concentrated solution, mix 2 quarts of **Arsenal PowerLine** with no more than 1 quart of water.

APPLICATION WITH DILUTE SOLUTIONS:

For cut stump treatments: Spray or brush the solution onto the cambium area of the freshly cut stump surface. Insure that the solution thoroughly wets the entire cambium area (the wood next to the bark of the stump).

For tree-injection treatments: Using standard injection equipment, apply 1 milliliter of solution at each injection site around the tree with no more than one-inch intervals between cut edges. Insure that the injector completely penetrates the bark at each injection site.

For frill or girdle treatments: Using a hatchet, machete, or similar device, make cuts through the bark at intervals around the tree with no more than two-inch intervals between cut edges. Spray or brush the solution into each cut until thoroughly wet.

APPLICATION WITH CONCENTRATED SOLUTIONS:

For tree injection treatments: Using standard injection equipment, apply 1 milliliter of solution at each injection site. Make at least one injection cut for every 3 inches of Diameter at Breast Height (DBH) on the target tree. For example, a 3-inch DBH tree will receive 1 injection cut, and a 6-inch DBH tree will receive 2 injection cuts. On trees requiring more than one injection site, place the injection cuts at approximately equal intervals around the tree.

For frill or girdle treatments: Using a hatchet, machete, or similar device, make cuts through the bark at approximately equal intervals around the tree. Make at least one cut for every 3 inches of DBH on the target tree. For example, a 3-inch DBH tree will receive 1 cut, and a 6-inch DBH tree will receive 2 cuts. Spray or brush the solution into each cut until thoroughly wet.

NOTE: Injury may occur to desirable woody plants if the shoots extend from the same root system or their root systems are grafted to those of the treated tree.

FOR CONTROL OF UNDESIRABLE WEEDS UNDER PAVED SURFACES

Arsenal® PowerLine™ herbicide can be used under asphalt, pond liners and other paved areas ONLY in industrial sites or where the pavement has a suitable barrier along the perimeter that prevents encroachment of roots of desirable plants.

Arsenal PowerLine should be used only where the area to be treated has been prepared according to good construction practices. If rhizomes, stolons, tubers or other vegetative plant parts are present in the site, they should be removed by scalping with a grader blade to a depth sufficient to insure their complete removal.

IMPORTANT: Paving should follow **Arsenal PowerLine** applications as soon as possible. **DO NOT** apply where the chemical may contact the roots of desirable trees or other plants.

The product is may not be used under pavement on residential properties such as driveways or parking lots, nor in recreational areas such as under bike or jogging paths, golf cart paths, or tennis courts, or where landscape plantings could be anticipated. Injury or death of desirable plants may result if this product is applied where roots are present or where they may extend into the treated area. Roots of trees and shrubs may extend a considerable distance beyond the branch extremities or so-called drip line.

APPLICATION DIRECTIONS FOR PAVED SURFACES:

Applications should be made to the soil surface only when final grade is established. **DO NOT** move soil following **Arsenal PowerLine** application.

Apply **Arsenal PowerLine** in sufficient water (at least 100 gal per acre) to ensure thorough and uniform wetting of the soil surface, including the shoulder areas. Add **Arsenal PowerLine** at a rate of 6 pints per acre (2.2 fluid ounce per 1000 square feet) to clean water in the spray tank during the filling operation. Agitate before spraying.

If the soil is not moist prior to treatment, incorporation of **Arsenal PowerLine** is needed for herbicide activation. **Arsenal PowerLine** can be incorporated into the soil to a depth of 4 to 6 inches using a rototiller or disc. Rainfall or irrigation of 1 inch will also provide uniform incorporation. **DO NOT** allow treated soil to wash or move into untreated areas.

FOR CONTROL OF UNDESIRABLE WEEDS IN UNIMPROVED BERMUDAGRASS AND BAHIAGRASS

Arsenal PowerLine may be used on unimproved bermudagrass and bahiagrass turf on roadsides, utility rights-of-way and other noncropland industrial sites. The application of **Arsenal PowerLine** on established common and coastal bermudagrass and bahiagrass provides control of labeled broadleaf and grass weeds. Competition from these weeds is eliminated, releasing the bermudagrass and bahiagrass. Treatment of bermudagrass with **Arsenal PowerLine** results in a compacted growth habit and seedhead inhibition.

Uniformly apply with properly calibrated ground equipment using at least 10 gallons of water per acre with a spray pressure 20 to 50 psi.

IMPORTANT: Temporary yellowing of grass may occur when treatment is made after growth commences.

DO NOT add surfactant in excess of the recommended rate (1 oz per 25 gallons of spray solution). **DO NOT APPLY** to grass during its first growing season. **DO NOT APPLY** to grass that is under stress from drought, disease, insects, or other causes.

DOSAGE RATES AND TIMING:

Bermudagrass - Apply **Arsenal PowerLine** at 6 to 12 oz per acre when the bermudagrass is dormant. Apply **Arsenal PowerLine** at 6 to 8 oz per acre after the bermudagrass has reached full green-up. Applications made during green-up will delay green-up. Include a surfactant in the spray solution (see **IMPORTANT** statement above).

For additional preemergence control of annual grasses and small seeded broadleaf weeds, add **Pendulum® herbicide** at the rate of 3.3 to 6.6 lbs per acre. Consult the **Pendulum** label for weeds controlled and for other use directions and precautions.

For control of Johnsongrass in bermudagrass turf, apply **Arsenal PowerLine** at 8 oz per acre plus **Roundup® herbicide** at 12 oz per acre plus surfactant. For additional control of broadleaves and vines, **Garlon® 3A** may be added to the above mix at the rate of 1-2 pints per acre. Observe all precautions and restrictions on the **Garlon 3A** and **Roundup** labels.

Bahiagrass - Apply **Arsenal PowerLine** at 4 to 8 oz per acre when the bahiagrass is dormant or after the grass has initiated green-up but has not exceeded 25% green-up. Include in the spray solution a surfactant (see **ADJUVANT** section for specific recommendations on surfactants).

WEEDS CONTROLLED:

- Bedstraw (*Galium* spp.)
- Bishopweed (*Ptilimnium capillaceum*)
- Buttercup (*Ranunculus parviflorus*)
- Carolina geranium (*Geranium carolinianum*)
- Fescue (*Festuca* spp.)
- Foxtail (*Setaria* spp.)
- Little barley (*Hordeum pusillum*)
- Seedling Johnsongrass (*Sorghum halepense*)
- White clover (*Trifolium repens*)
- Wild carrot (*Daucus carota*)
- Yellow woodsorrel (*Oxalis stricta*)

GRASS GROWTH AND SEEDHEAD SUPPRESSION

Arsenal® PowerLine™ herbicide may be used to suppress growth and seedhead development of certain turf-grass in unimproved areas. When applied to desirable turf, **Arsenal PowerLine** may result in temporary turf damage and/or discoloration. Effects to the desirable turf may vary with environmental conditions. For optimum performance, application should be made prior to culm elongation. Applications may be made before or after mowing. If applied prior to mowing, allow at least three days of active growth before mowing. If following a mowing, allow sufficient time for the grasses to recover before applying this product or injury may be amplified.

DO NOT APPLY to turf under stress (drought, cold, insect damaged, etc.) or severe injury or death may occur.

Bermudagrass - Apply **Arsenal PowerLine** at 6 to 8 oz per acre from early green-up to prior to seed head initiation. **DO NOT** add a surfactant for this application.

Cool Season Unimproved Turf - Apply **Arsenal PowerLine** at 2 oz per acre plus 0.25% nonionic surfactant. For increased suppression, **Arsenal PowerLine** may be tank mixed with such products as **Campaign®** (24 oz per acre) or **Embark®** (8 oz per acre).

Tank mixes may increase injury to desired turf. Consult each product label for recommended turf species and other use directions and precautions. Tank mixes with 2,4-D or products containing 2,4-D may decrease the effectiveness of **Arsenal PowerLine**.

TOTAL VEGETATION CONTROL WHERE BAREGROUND IS DESIRED

Arsenal PowerLine is an effective herbicide for preemergence or postemergence control of many annual and perennial broadleaf and grass weeds where bareground is desired. **Arsenal PowerLine** is particularly effective on hard-to-control perennial grasses. **Arsenal PowerLine** at 1.5 to 6 pints per acre can be used alone or in tank mix with **Roundup®, Finale®, MSMA, Karmex®, Pendulum®, simazine, Banvel®, Vanquish®,** or **Oust® herbicides**. The degree and duration of control are dependent on the rate of **Arsenal PowerLine** used, tank mix partner, the volume of carrier, soil texture, rainfall and other conditions.

Consult manufacturer's labels for specific rates and weeds controlled. Always follow the more restrictive label when making an application involving tank mixes.

TANK MIX INSTRUCTIONS FOR BAREGROUND

Herbicide Rates per Acre*

Arsenal PowerLine	Pendulum® WDG	Pendulum 3.3 EC	Diuron
Rate in pints	in lbs	in Quarts	in lbs ai
1.5 - 3	6.6	4.8	4 - 6
2 - 4	6.6	4.8	6 - 10
3 - 6	6.6	4.8	8 - 12

* Use higher rates for fall applications and in areas that have not been previously treated or that feature heavy infestations.

Applications of **Arsenal PowerLine** may be made anytime of the year. Use equipment calibrated to deliver desired gallons per acre spray volume and uniformly distribute the spray pattern over the treated area.

Postemergence Applications: Always use a spray adjuvant (see **ADJUVANT** section of this label) when making a postemergence application. For optimum performance on tough to control annual grasses, applications should be made at a total volume of 100 gallons per acre or less. For quicker burndown or brown-out of target weeds, **Arsenal PowerLine** may be tank mixed with products such as **Roundup®, Finale,** or **MSMA**. Tank mixes with 2,4-D or products containing 2,4-D have reduced performance of **Arsenal PowerLine**. Always follow the more restrictive label when tank mixing.

Spot Treatments: Arsenal PowerLine may be used as a follow-up treatment to control escapes or weed encroachment in a bareground situation. To prepare the spray solution, thoroughly mix in each gallon of water 0.5 to 5% **Arsenal PowerLine** plus an adjuvant. For increased burndown, include **Roundup®, Finale,** **MSMA,** or similar products. For added residual weed control or to increase the weed spectrum, add **Pendulum** or diuron. Always follow the more restrictive label when tank mixing.

FOR SPOT TREATMENT WEED CONTROL IN GRASS PASTURE AND RANGELAND

For the control of undesirable vegetation in grass pasture and rangeland, **Arsenal PowerLine** may be applied as a spot treatment at a rate of 2 to 48 fluid oz of product per treated acre using any of the described ground application methods. Spot applications to grass pasture and rangeland may not exceed more than one-tenth of the area to be grazed or cut for hay. See appropriate sections of this label for specific use directions for the application method and vegetation control desired. **DO NOT** apply more than 48 fluid oz per acre per year.

Grazing and Haying Restrictions: There are no grazing restrictions following **Arsenal PowerLine** application. **DO NOT** cut forage grass for hay for seven days after **Arsenal PowerLine** application.

17/21

INSTRUCTIONS FOR RANGELAND USE

Arsenal® PowerLine™ herbicide may be applied to rangeland for the control of undesirable vegetation in order to achieve one or more of the following vegetation management objectives:

1. The control of undesirable (non-native, invasive and noxious) plant species
2. The control of undesirable vegetation in order to aid in the establishment of desirable rangeland plant species.
3. The control of undesirable vegetation in order to aid in the establishment of desirable rangeland vegetation following a fire.
4. The control of undesirable vegetation for purposes of wildfire fuel reduction.
5. The release of existing desirable rangeland plant communities from the competitive pressure of undesirable plant species.
6. The control of undesirable vegetation for purposes of wildlife habitat improvement.

To ensure the protection of threatened and endangered plants when applying **Arsenal PowerLine** to rangeland:

1. Federal agencies must follow NEPA regulations to ensure protection of threatened and endangered plants.
2. State agencies must work with the Fish and Wildlife Service or the Service's designated state conservation agency to ensure protection of threatened and endangered plants.
3. Other organizations or individuals must operate under a Habitat Conservation Plan if threatened or endangered plants are known to be present on the land to be treated.

Please see the appropriate section(s) of this label for specific use directions for the desired rangeland vegetation management objective.

Arsenal PowerLine should only be applied to a given rangeland acre as specific weed problems arise. Long-term control of undesirable weed species ultimately depends on the successful use of land management practices that promote the growth and sustainability of desirable rangeland plant species.

ROTATIONAL CROP INSTRUCTIONS

Rotational crops may be planted twelve months after applying **Arsenal PowerLine** at the specified pasture and rangeland rate. Following twelve months after an **Arsenal PowerLine** application and before planting any crop, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted in the previously treated area in the grass pasture/rangeland and grown to maturity. The test strip should include low areas and knolls, and include variations in soil type and pH within the treated area. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

Use of **Arsenal PowerLine** in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environ-

mental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

WEEDS CONTROLLED BY ARSENAL POWERLINE

Arsenal PowerLine will provide preemergence or postemergence control with residual control of the following target vegetation species at the rates listed. Residual control refers to control of newly germinating seedlings in both annuals and perennials. In general, annual weeds may be controlled by preemergence or postemergence applications of **Arsenal PowerLine**. **For established biennials and perennials, postemergence applications of Arsenal PowerLine are recommended.** The rates shown below pertain to broadcast applications and indicate the relative sensitivity of these weeds. The relative sensitivity should be referenced when preparing low-volume spray solutions (see **Low-Volume** section of **GROUND APPLICATIONS**); low-volume applications may provide control of the target species with less **Arsenal PowerLine** per acre than is shown for the broadcast treatments. **Arsenal PowerLine** may be used only in accordance with the instructions on this label and the leaflet label.

Resistant Biotypes: Naturally occurring biotypes (a plant within a given species that has a slightly different but distinct genetic makeup from other plants of the same species) of some weeds listed on this label may not be effectively controlled by this and/or other herbicides (**Oust®**) with the ALS/AHAS enzyme inhibiting mode of action. If naturally occurring ALS/AHAS resistant biotypes are present in an area, **Arsenal PowerLine** should be tank mixed or applied sequentially with an appropriate registered herbicide having a different mode of action to ensure control.

15/24

GRASSES

COMMON NAME	SPECIES	GROWTH HABIT ²
Apply 2-3 pints per acre¹		
Annual bluegrass	(<i>Poa annua</i>)	A
Broadleaf signalgrass	(<i>Brachiaria platyphylla</i>)	A
Canada bluegrass	(<i>Poa compressa</i>)	P
Downy brome	(<i>Bromus tectorum</i>)	A
Fescue	(<i>Festuca</i> spp.)	A/P
Foxtail	(<i>Setaria</i> spp.)	A
Italian ryegrass	(<i>Lolium multiflorum</i>)	A
Johnsongrass	(<i>Sorghum halepense</i>)	P
Kentucky bluegrass	(<i>Poa pratensis</i>)	P
Lovegrass	(<i>Eragrostis</i> spp.)	A/P
Orchardgrass	(<i>Dactylis glomerata</i>)	P
Paragrass	(<i>Brachiaria mutica</i>)	P
Quackgrass	(<i>Agropyron repens</i>)	P
Sandbur	(<i>Cenchrus</i> spp.)	A
Sand dropseed	(<i>Sporobulus cryptandrus</i>)	A
Smooth brome	(<i>Bromus inermis</i>)	P
Vaseygrass	(<i>Paspalum urvillei</i>)	P
Wild oats	(<i>Avena fatua</i>)	A
Witchgrass	(<i>Panicum capillare</i>)	A
Apply 3-4 pints per acre¹		
Barnyardgrass ³	(<i>Echinochloa crus-galli</i>)	A
Beardgrass	(<i>Andropogon</i> spp.)	P
Bluegrass, annual ³	(<i>Poa annua</i>)	A
Cheat	(<i>Bromus secalinus</i>)	A
Crabgrass	(<i>Digitaria</i> spp.)	A
Crowfootgrass ³	(<i>Dactyloctenium aegyptium</i>)	A
Fall panicum	(<i>Panicum dichotomiflorum</i>)	A
Giant reed	(<i>Arundo donax</i>)	P
Goosegrass	(<i>Eleusine indica</i>)	A
Itchgrass ³	(<i>Rottboellia exaltata</i>)	A
Junglerice ³	(<i>Echinochloa colonum</i>)	A
Lovegrass ³	(<i>Eragrostis</i> spp.)	A
Maidencane	(<i>Panicum hemitomon</i>)	A
Panicum, browntop ³	(<i>Panicum fasciculatum</i>)	A
Panicum, Texas ³	(<i>Panicum texanum</i>)	A
Prairie threeawn	(<i>Aristida oligantha</i>)	P
Reed canarygrass	(<i>Phalaris arundinacea</i>)	P
Sandbur, field ³	(<i>Cenchrus incertus</i>)	A
Signalgrass ³	(<i>Brachiaria</i> spp.)	A
Torpedograss	(<i>Panicum repens</i>)	P
Wild barley	(<i>Hordeum</i> spp.)	A
Wooly cupgrass ³	(<i>Eriochloa villosa</i>)	A

16/21

GRASSES (continued)

COMMON NAME	SPECIES	GROWTH HABIT ²
Apply 4-6 pints per acre¹		
Bahiagrass	(<i>Paspalum notatum</i>)	P
Bermudagrass ⁴	(<i>Cynodon dactylon</i>)	P
Big bluestem	(<i>Andropogon gerardii</i>)	P
Cattail	(<i>Typha</i> spp.)	P
Cogongrass	(<i>Imperata cylindrica</i>)	P
Dallisgrass	(<i>Paspalum dilatatum</i>)	P
Feathertop	(<i>Pennisetum villosum</i>)	P
Guineagrass	(<i>Panicum maximum</i>)	P
Phragmites	(<i>Phragmites australis</i>)	P
Prairie cordgrass	(<i>Spartina pectinata</i>)	P
Saltgrass ⁴	(<i>Distichlis stricta</i>)	P
Sand dropseed	(<i>Sporobolus cryptandrus</i>)	P
Sprangletop ³	(<i>Leptochloa</i> spp.)	A
Timothy	(<i>Phleum pratense</i>)	P
Wirestem muhly	(<i>Muhlenbergia frondosa</i>)	P

BROADLEAF WEEDS

Apply 2-3 pints per acre¹

African rue	(<i>Peganum harmala</i>)	P
Alligatorweed	(<i>Alternanthera philoxeroides</i>)	A/P
Burdock	(<i>Arctium</i> spp.)	B
Carolina geranium	(<i>Geranium carolinianum</i>)	A
Carpetweed	(<i>Mollugo verticillata</i>)	A
Clover	(<i>Trifolium</i> spp.)	A/P
Common chickweed	(<i>Stellaria media</i>)	A
Common ragweed	(<i>Ambrosia artemisiifolia</i>)	A
Dandelion	(<i>Taraxacum officinale</i>)	P
Dogfennel	(<i>Eupatorium capillifolium</i>)	A
Filaree	(<i>Erodium</i> spp.)	A
Fleabane	(<i>Erigeron</i> spp.)	A
Hoary vervain	(<i>Verbena stricta</i>)	P
Indian mustard	(<i>Brassica juncea</i>)	A
Kochia ⁵	(<i>Kochia scoparia</i>)	A
Lambsquarters	(<i>Chenopodium album</i>)	A
Lespedeza	(<i>Lespedeza</i> spp.)	P
Miners lettuce	(<i>Montia perfoliata</i>)	A
Mullein	(<i>Verbascum</i> spp.)	B
Nettleleaf goosefoot	(<i>Chenopodium murale</i>)	A
Oxeye daisy	(<i>Chrysanthemum leucanthemum</i>)	P
Pepperweed	(<i>Lepidium</i> spp.)	A
Pigweed	(<i>Amaranthus</i> spp.)	A
Puncturevine	(<i>Tribulus terrestris</i>)	A
Russian thistle	(<i>Salsola kali</i>)	A
Smartweed	(<i>Polygonum</i> spp.)	A
Sorrell	(<i>Rumex</i> spp.)	P
Sunflower	(<i>Helianthus</i> spp.)	A
Sweet clover	(<i>Melilotus</i> spp.)	A
Tansymustard	(<i>Ambrosia psilostachya</i>)	P
Wild carrot	(<i>Daucus carota</i>)	B
Wild lettuce	(<i>Lactuca</i> spp.)	A/B
Wild parsnip	(<i>Pastinaca sativa</i>)	B

BROADLEAF WEEDS (continued)

COMMON NAME	SPECIES	GROWTH HABIT ²
Apply 2-3 pints per acre¹		
Wild turnip	(<i>Brassica campestris</i>)	B
Woollyleaf bursage	(<i>Franseria tomentosa</i>)	P
Yellow woodsorrel	(<i>Oxalis stricta</i>)	P
Apply 3-4 pints per acre¹		
Broom snakeweed ⁶	(<i>Gutierrezia sarothrae</i>)	P
Bull thistle	(<i>Cirsium vulgare</i>)	B
Burclover ³	(<i>Medicago</i> spp.)	A
Chickweed, mouseear ⁵	(<i>Cerastium vulgatum</i>)	A
Clover, hop ³	(<i>Trifolium procumbens</i>)	A
Cocklebur	(<i>Xanthium strumarium</i>)	A
Cudweed ³	(<i>Gnaphalium</i> spp.)	A
Desert camelthorn	(<i>Alhagi pseudalhagi</i>)	P
Diffuse knapweed	(<i>Centaurea diffusa</i>)	A
Dock	(<i>Rumex</i> spp.)	P
Fiddleneck ³	(<i>Amsinckia intermedia</i>)	A
Goldenrod	(<i>Solidago</i> spp.)	P
Henbit ³	(<i>Lamium aplexicaule</i>)	A
Knotweed, prostrate ³	(<i>Polygonum aviculare</i>)	A/P
Pokeweed	(<i>Phytolacca americana</i>)	P
Purple loosestrife ⁶	(<i>Lythrum salicaria</i>)	P
Purslane	(<i>Portulaca</i> spp.)	A
Pusley, Florida ³	(<i>Richardia scabra</i>)	A
Rocket, London ³	(<i>Sisymbrium irio</i>)	A
Rush skeletonweed ⁶	(<i>Chondrilla juncea</i>)	B
Saltbush	(<i>Atriplex</i> spp.)	A
Shepherdspurse ³	(<i>Capsella bursa-pastoris</i>)	A
Spurge, annual ³	(<i>Euphorbia</i> spp.)	A
Stinging nettle ⁶	(<i>Urtica dioica</i>)	P
Velvetleaf ³	(<i>Abutilon theophrasti</i>)	A
Yellow starthistle	(<i>Centaurea solstitialis</i>)	A
Apply 4-6 pints per acre¹		
Arrowwood	(<i>Pluchea sericea</i>)	A
Canada thistle	(<i>Cirsium arvense</i>)	P
Giant ragweed	(<i>Ambrosia trifida</i>)	A
Grey rabbitbrush	(<i>Chrysothamnus nauseosus</i>)	P
Japanese bamboo/knotweed	(<i>Polygonum cuspidatum</i>)	P
Little mallow	(<i>Malva parviflora</i>)	B
Milkweed	(<i>Asclepias</i> spp.)	P
Primrose	(<i>Oenothera kunthiana</i>)	P
Russian knapweed	(<i>Centaurea repens</i>)	P
Sago pondweed	(<i>Potamogeton pectinatus</i>)	P
Silverleaf nightshade	(<i>Solanum elaeagnifolium</i>)	P
Sowthistle	(<i>Sonchus</i> spp.)	A
Texas thistle	(<i>Cirsium texanum</i>)	P

VINES AND BRAMBLES

COMMON NAME	SPECIES	GROWTH HABIT ²
Apply 1 pint per acre		
Field bindweed	(<i>Convolvulus arvensis</i>)	P
Hedge bindweed	(<i>Calystegia sepium</i>)	A
Apply 2-3 pints per acre¹		
Wild buckwheat	(<i>Polygonum convolvulus</i>)	P
Apply 3-4 pints per acre¹		
Greenbriar	(<i>Smilax</i> spp.)	P
Honeysuckle	(<i>Lonicera</i> spp.)	P
Morningglory	(<i>Ipomoea</i> spp.)	A/P
Poison ivy	(<i>Rhus radicans</i>)	P
Redvine	(<i>Brunnichia cirrhosa</i>)	P
Wild rose	(<i>Rosa</i> spp.)	P
Including: Multiflora rose	(<i>Rosa multiflora</i>)	P
Macartney rose	(<i>Rosa bracteata</i>)	P
Apply 4-6 pints per acre¹		
Kudzu ⁴	(<i>Pueraria lobata</i>)	P
Trumpet creeper	(<i>Campsis radicans</i>)	P
Virginia creeper	(<i>Parthenocissus quinquefolia</i>)	P
Wild grape	(<i>Vitis</i> spp.)	P
BRUSH SPECIES		
Apply 4-6 pints per acre¹		
American beech	(<i>Fagus grandifolia</i>)	P
Ash	(<i>Fraxinus</i> spp.)	P
Bald cypress	(<i>Taxodium distichum</i>)	P
Bigleaf maple	(<i>Acer macrophyllum</i>)	P
Blackgum	(<i>Nyssa sylvatica</i>)	P
Black locust ⁷	(<i>Robinia pseudoacacia</i>)	P
Boxelder	(<i>Acer negundo</i>)	P
Brazilian peppertree	(<i>Schinus terebinthifolius</i>)	P
Cherry	(<i>Prunus</i> spp.)	P
Chinaberry	(<i>Melia azadarach</i>)	P
Chinese tallow-tree	(<i>Sapium sebiferum</i>)	P
Dogwood	(<i>Cornus</i> spp.)	P
Elm ⁸	(<i>Ulmus</i> spp.)	P
Hawthorn	(<i>Crataegus</i> spp.)	P
Hickory	(<i>Carya</i> spp.)	P
Honeylocust ⁹	(<i>Gleditsia triacanthos</i>)	P
Maple	(<i>Acer</i> spp.)	P
Melaleuca	(<i>Melaleuca quiquenervia</i>)	P
Mulberry	(<i>Morus</i> spp.)	P
Oak	(<i>Quercus</i> spp.)	P
Persimmon	(<i>Diospyros virginiana</i>)	P
Poplar	(<i>Populus</i> spp.)	P
Privet	(<i>Ligustrum vulgare</i>)	P
Red alder	(<i>Alnus rubra</i>)	P
Red maple	(<i>Acer rubrum</i>)	P
Russian olive	(<i>Eleagnus angustifolia</i>)	P
Saltcedar	(<i>Tamarix ramosissima</i>)	P
Sassafras	(<i>Sassafras albidum</i>)	P

BRUSH SPECIES (continued)

Apply 4-6 pints per acre¹

COMMON NAME	SPECIES	GROWTH HABIT ²
Sourwood	(<i>Oxydendrum arboreum</i>)	P
Sumac	(<i>Rhus</i> spp.)	P
Sweetgum	(<i>Liquidambar styraciflua</i>)	P
Willow	(<i>Salix</i> spp.)	P
Yellow poplar	(<i>Liriodendron tulipifera</i>)	P

¹ The higher rates should be used where heavy or well-established infestations occur.

² Growth Habit:- A = Annual, B = Biennial, P = Perennial

³ For preemergence control, tank mix with **Pendulum® herbicide**.

⁴ Use a minimum of 75 GPA - control of established stands may require repeat applications.

⁵ For preemergence control, tank mix with **Pendulum, Karmex®** or diuron.

⁶ For best results, early postemergence applications are required.

⁷ Tank mix with **Roundup®, Accord®, Escort®, Krenite®, Garlon® 3A** or **Tordon® K**.

⁸ Tank mix with **Roundup, Accord, or Escort**.

⁹ Tank mix with **Roundup, Accord, Garlon 3A** or **Tordon K**.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

807

Arsenal, Banvel and Pendulum are registered trademarks of BASF.

PowerLine is a trademark of BASF

Microfoil is a trademark of Rhone Poulenc Ag. Company.

Thru-Valve is a trademark of Waldrum Specialties.

Accord, Campaign and Roundup are registered trademarks of Monsanto Company.

Escort, Karmex, Krenite, Oust and Telar are registered trademarks of E. I. duPont de Nemours and Company.

Garlon and Tordon are registered trademarks of Dow AgroSciences Company.

Embark is a registered trademark of PBI/Gordon Corporation.

Finale is a registered trademark of Bayer.

Vanquish is a trademark of a Syngenta Group Company.

© 2007 BASF Corporation
All rights reserved.

000241-00431.20071106.NVA 2007-04-295-0100
Based on: 2006-04-295-0291

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709



The Chemical Company



The Chemical Company

21/24

November 6, 2007

U.S. Environmental Protection Agency
Office of Pesticide Programs (7505P)
Document Processing Desk 7504P (NOTIF)
Room S-4900
One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202
Attention: Mr. James Tompkins, (PM 25)

**RE: Notification of Label Changes for Arsenal PowerLine herbicide
EPA Reg. No. 241-431**

Dear Mr. Tompkins:

BASF is hereby submitting a notification for minor label changes for Arsenal PowerLine herbicide (EPA Reg. No. 241-431).

Enclosed please find:

- Application form 8570-1
- CD containing electronic copy of the label
- Certification with Respect to Label Integrity
- Arsenal PowerLine herbicide label
- Arsenal PowerLine herbicide label with annotated changes
- Approved Arsenal PowerLine herbicide label

No PRIA fee is required for this notification.

Thank you for your assistance with this matter. If you should have any questions, please feel free to call me at (919) 547-2622.

Regards,

A handwritten signature in black ink, appearing to read "J. Birk".

Jeffrey H. Birk, Ph.D.
Regulatory Manager
Phone 919-547-2622
Fax: 919-547-2850
Email: jeffrey.birk@basf.com

® Registered Trademark of BASF