

* Rinse empty container titoroughly with water and discard it. *

SAFE FOR SEPT

SANITIZING TOILET BOWLS:

1 — Saturate the swab mop wit

2 — Clean under the rim of box

3 — Clean bottom and top of si

4 — Flush, rinsing swab in the I

5 — To deodorize toilet bowls,

SANITIZING URINALS:

1 — Saturate swab mop with V

2 — Start at top of urinal and p

PREPARATIC FOR T

Remove residual water from b SHEEN with forty parts of wate be disinfected. Next swab with cleaned areas.

DANGER:

Keep Out of Reach

not get in eyes, on when handling. Har

FIRST AID:

NOTE TO

In case of contact, 15 minutes. For ey

fore reuse.

If swallowed, drink if these are not ava cian immediately.

Probable mucosal PHYSICIAN: against circulatory

CCEPTED DER THE FEDERAL INSECTIONS GIGIDE AND RODENTICIDE CT ECONOMIC POISON REGISTER. UNDER NO. SUBJECT ATTACHED COMMENTS.

OF CHILDREN

cautions

609

ard it. *

No

SAFE FOR SEPTIC TANKS — Do Not Use on Porcelain Surfaces DIRECTIONS FOR USE

SANITIZING TOILET BOWLS:

- 1 Saturate the swab mop with VITRO SHEEN.
- 2 Clean under the rim of bowl and progress downward.
- 3 Clean bottom and top of seat with damp swab mop.
- 4 Flush, rinsing swab in the running water.
- 5 To deodorize toilet bowls, add two ounces of VITRO SHEEN to water in bowl.

SANITIZING URINALS:

- 1 Saturate swab mop with VITRO SHEEN.
- 2 Start at top of urinal and purgress downward.

PREPARATION OF A DISINFECTANT SOLUTION FOR TOILET BOWLS AND URINALS

Remove residual water from bowl by expelling over trap. Dilute one part of VITRO SHEEN with forty parts of water. Apply this solution liberally with swab mop to areas to be disinfected. Next swab with a fresh solution of VITRO SHEEN for disinfection of cleaned areas.

DANGER: Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do

not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves

when handling. Harmful or 'atal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing be-

fore reuse.

If swallowed, drink prompily a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physi-

cian immediately.

NOTE TO Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed. PHYSICIAN: