

# American Fluoride Corporation

855 Avenue of the Americas  
New York, N. Y. 10001  
212-594-3820

RODENTICIDE, BAIT, ANTICOAGULANT PIVALYL

FOR CONTROL OF COMMENSAL RATS AND MICE

Stock No. 6840-753-4973

CAUTION: KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

2-Pivalyl-1,3-Indandione 0.025%

INERT INGREDIENTS:

Liquid Petrolatum	4.8 %	99.975%
Sucrose	4.8 %	
Dye*	0.05 %	
Breakfast Rolled Oats	90.325 %	

TOTAL 100.000%

\*FOR USDA USE ONLY: Flaming Red G305  
Manufacturer: OX COLOR WORKS  
New York, N. Y.

DIRECTIONS: Place 4 to 8 ounces of Rodenticide, Bait, Anticoagulant in locations where rats and mice feed, water, or travel. Bait placements along walls, in corners and in concealed locations where rats and mice can feed unobserved are desirable. Bait should be placed in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes. Contaminated or spoiled baits should be replaced promptly. Baiting should continue until all signs of feeding have stopped. Where a continuous source of infestation is present, permanent bait stations should be established and replenished as needed.

For Rats: Use at least four ounces of bait per station. An uninterrupted supply of bait should be maintained for at least 10 days.

For Mice: Use tablespoon amounts (1/4 - 1/2 ounce) of bait at 8 - 12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

ACCEPTED

- Continued -

MAR 13 1967

# American Fluoride Corporation

PAGE 2

855 Avenue of the Americas  
New York, N. Y. 10001  
212-594-3820

CAUTION: Keep away from humans, domestic animals, and pets. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

NET WEIGHT:

USDA REG. NO. 255-80

LOT NO.: