

WARNING: Keep out of
See additional directions and

DIRECTIONS. Lay down a paper-thin layer of dust along rat or mouse runways. Heavily dust rat burrows and under rat or mouse infested buildings. Apply dust with a dust pump or hand applicator. In some cases it may be necessary to repeat dusting of heavily used runways or renew dust applications that have been removed by rain, etc.

Control should be achieved in 10-14 days. One pound of AFC ANTICOAGULANT TRACKING POWDER should cover approximately 40 square feet of rat runway.


WARNING: The active ingredient of this product is an anticoagulant chemical. Avoid ingestion of the dust. Areas dusted with this product must be protected so that humans, domestic animals, wildlife and pets will not come in contact with this dust. Avoid inhalation. Wear dust respirator when applying or removing dust. Wash thoroughly after applying dust.

WARNING: Do not use in feed or food processing plants, or in any location where the possibility exists of contaminating feed or food items.

ANTIDOTE: In case of ingestion by humans, induce vomiting by giving a tablespoon of salt in a glass of warm water and repeating until vomit fluid is clear. Call a physician immediately. Treatment by physician should include the administration of oral or intravenous injection of Vitamin K₁. In critical situations a transfusion, with fresh whole blood should be given.

TERA

anticoagulant tr



kills rats & mice



WARNING: Keep out of reach of children
See additional directions and warnings on left panel.

TRAC[®]

coagulant tracking powder

kills rats & mice



FOR USE BY
PROFESSIONAL
PEST CONTROL
OPERATORS
ONLY

2.10%

| | |
|--|---------|
| Calcium Salt of 2-Isovaleryl-1,3-Indandione* | 2.18% |
| | 97.82% |
| | 100.00% |

*Equivalent to 2-Isovaleryl-1,3-Indandione 2.00%

WARNING: Keep away from children, wildlife and domestic animals. Avoid contamination of feed and foodstuffs.

American Fluoride Corporation
New Rochelle, New York 10801

NET WEIGHT