

American Fluoride Corporation

505 Avenue of the Americas
New York, N. Y. 10011
212-941-181

AFC DIPHACINONE 0.1%

Anticoagulant Concentrate Kills Rats and Mice

CAUTION: KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

Diphacinone (2-Diphenylacetyl-1,5-indandione).....0.1%

INERT INGREDIENTS:.....99.9%

TOTAL.....100.0%

DIRECTIONS

- a. Thoroughly mix one part (by weight) of rodenticide with nineteen parts of dry cereal bait.
- b. Place bait in locations where rats and mice feed, water or travel.
- c. Bait placements along walls, in corners, and in concealed locations where rats and mice feed are desirable.
- d. Place baits in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes.

FOR RAT CONTROL

- a. Place at least four ounces of bait at each bait location.
- b. Maintain an uninterrupted supply of bait in the stations for approximately 10 days.

FOR MOUSE CONTROL

- a. Place tablespoon amounts of bait at 8-10 foot intervals.
- b. Maintain an uninterrupted supply of bait for at least 10 days.

NOTE: If Bait molds, decomposes or becomes sour, replace with fresh material.

CAUTION

Keep ~~away~~ from humans, domestic animals and pets. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, the intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.



USDA REG. NO. 255-X

