

American Fluoride Corporation

ACCEPTED

355 Avenue of the Americas

New York, N. Y. 10001

212 594 3826

RODENTICIDE, BAIT, ANTICOAGULANT, WARFARIN

For Control of Commensal Rats and Mice

TO ATTENTION

CAUTION: KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS: 0.025%
Warfarin 5-(a-acetonylbenzyl)-4-hydroxycoumarin

INERT INGREDIENTS: 99.975%
Powdered Sugar 7.80%
Petrolatum 4.80%
Breakfast Rolled Oats 28.50%
Cracked Corn 19.00%
Corn Meal 42.875%
TOTAL 100.000%

GENERAL DIRECTIONS: Place bait in shallow containers in dry locations where rats and mice feed, water, or travel. It is desirable to place baits along walls, in corners, and in concealed locations, where rats and mice can feed unobserved. Contaminated or spoiled baits should be replaced promptly. Baiting should continue until all signs of feeding have stopped. Where a continuous source of infestation is present (nearby dumps, for example) permanent bait stations should be established and replenished as needed.

BAIT SHOULD BE PLACED IN LOCATIONS NOT ACCESSIBLE TO CHILDREN, PETS AND DOMESTIC ANIMALS, AS IN TAMPER-PROOF BAIT BOXES.

FOR RAT CONTROL: Use 1/4 - 1 pound per bait station, at points of greatest activity. Maintain an uninterrupted supply of bait for at least 10 days.

FOR MOUSE CONTROL: Use tablespoon amounts (1/4 - 1/2 ounce) per bait station. Bait should be placed at 8 - 12 foot intervals. Maintain an uninterrupted supply of bait for at least 15 days.

CAUTION: Keep away from humans, domestic animals and pets. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of dihydroxycoumarin.

Net Weight:

U.S. PAT. NO. 2,554,444