

MAY 04 1988

Cello Corp.
1354 Old Post Road
Havre de Grace, MD 21078

Attention: W. Thomas Ruckle

Gentlemen:

Subject: Fuldicide 14 Cleaner Disinfectant Deodorizer Fungicide
EPA Registration No. 257-262
Your Submission Dated November 19, 1987


The labeling referred to above, submitted in connection with the Label Improvement Program Notice for Antimicrobial Products Used as Food Contact Surface Sanitizers, is acceptable and a stamped copy is enclosed for your records.

However, at the printing of the next set of labels add the following misuse statement after the USE DIRECTIONS heading.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

If you have any questions concerning this letter, please call Mr. Robert Young at (703) 557-3966.

Sincerely yours,



John H. Lee
Product Manager (31)
Disinfectants Branch
Registration Division (TS-767C)

Enclosure

58284:I:J:EL 1:KENCO:04/29/88:05/05/88:EAE:jh

CONCURRENCES

SYMBOL	ORIGINATOR							
SURNAME								
DATE								

This product is a heavy duty cleaner and disinfectant designed for general purpose use where heavy duty cleaning is of prime importance. When used as directed, it will remove most dirt and grime in a single simple application. When used for industrial cleaning, follow the use directions shown below, for best results. Not recommended for pseudomonas.

DANGER: Keep Out of Reach of Children. Corrosive - causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and protect skin when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. Immediately call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call physician.

USE DIRECTIONS

For light duty cleaning, and deodorizing, use two ounces per gallon.

For heavy duty cleaning and disinfecting and fungicidal activity against pathogenic fungi, use 4.5 ounces per gallon.

fuldicide-14

**CLEANER • DISINFECTANT
DEODORIZER • FUNGICIDE**

DESIGNED FOR HOSPITAL AND INSTITUTIONAL USE

ACTIVE INGREDIENTS:

Alkyl (60% C14), 30% C16, 5% C12, 5% C18)	
dimethyl benzyl ammonium chlorides	0.8%
Alkyl (68% C12, 32% C14) dimethyl	
ethylbenzyl ammonium chlorides	0.8%

INERT INGREDIENTS:

98.4%
100.0%

ACCEPTED
with COMMENTS
EPA Label Dated:

EPA Reg. No. 257-262

EPA Est. 257-MD-1

MAY 04 1988

Keep Out Of Reach Of Children

DANGER

SEE FIRST AID STATEMENT AND OTHER
PRECAUTIONARY STATEMENTS ON SIDE PANEL

Wink the Federal...
Per...
as...
sept...
257-262

NET CONTENTS: GALLONS



CELLO CORP.
Havre de Grace, MD 21078-0366
301-939-1234 • 800-638-4850



Member of Grow Group, Inc.

Made for Industrial & Institutional Use Only



For best re...
mop and...
other hãr...
For extã...
cleaning u...
Do not mi...
solutions...
dilutions...
Rinse em...
with wate...

STORAGE

Store only...
container...
sible to ch...
water, loc...
disposal...
acutely ha...
disposal...
mixture, c...
Federal L...
be dispos...
label instr...
State Pes...
Control A...
waste rep...
EPA Reg...
Do not re...
Triple rin...
water Re...
reconditi...
dispose o...
and local...
container...
sanitary l...
allowed b...
burning. I...
smoke.

Contents, Con...

Some states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated, and appropriate actions to take in case of accidental exposure). When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: DANGER: This product is treated with TRIPHENYL TIN HYDROXIDE on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure see Statement of Practical Treatment. SUPER-TIN® 4L is a non-flammable, flowable fungicide.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or offer for sale and disposal in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

SUPER-TIN® 4L Fungicide can be applied as a ground or aerial spray to control fungicidal infections on listed crops. Application rates are for general use. The state agricultural extension or agricultural experiment station specialists should be consulted for specific application and timing recommendations. With any spray application thorough coverage is essential for good control.

Do not apply this product through any type of irrigation system.

PRECAUTIONS

SUPER-TIN® 4L Fungicide can be applied in combination with SUPER SIX™ for additional fungicidal control. Do not add surfactants, spreaders, stickers or buffers. Combinations with some pesticides, micro-nutrients, spreaders, stickers, surfactants or other ingredients can increase phytotoxicity. Phytotoxicity may be severe. Emulsifiable concentrate insecticides can be especially injurious in combination. We do not recommend combinations of the above unless testing or prior experience has shown the mixture to be non-phytotoxic to the crop. Do not graze dairy or meat animals in treated areas.

GENERAL INSTRUCTIONS FOR APPLICATION

CROP	DISEASE	RATE PER ACRE	DIRECTIONS
Sugar Beets	Cercospora Leafspot	4 - 8 fl oz	Ground Application: Apply in at least 15 gallons of water. Full coverage of the foliage is necessary for best results.
		4 - 8 fl oz	Aerial Application (helicopter or fixed wing aircraft): Apply in 5 - 10 gals of water. Diluted spray should be directed uniformly to all parts of the plant. Use lower gallonage when plants are small and increase volume with plant size. Use the lower rate for protective sprays and the higher rates later in the season or during high infection periods. Application should begin when leafspot conditions appear or when the disease is in the area and repeated at 10 to 14 day intervals up to a maximum of 3 applications per growing season. Do not treat within 14 days of harvest. Do not graze or feed beet tops to livestock.
Potatoes	Early and Late Blight, Suppression of Colorado Potato Beetle	4 - 8 fl oz	Ground Application: Apply in at least 15 gallons of water. Full coverage of the foliage is necessary for best results. A spray pressure of less than 200 pounds is recommended.
		4 - 8 fl oz	Aerial Application: Apply in 3 - 10 gallons of water. For helicopter application fly high enough so as not to whip the vines. Diluted spray should be directed uniformly to all parts of the plant and the gallonage increased according to the size of the plants. Application should begin with the appearance of blight weather conditions and continue on a 7 day schedule of up to a maximum of 3 applications per growing season. Do not treat within 7 days of harvest. The lower rate of application should be used early in the season and the high rate mid to late season or when blight infection is in the area.
Carrots	Cercospora Leafspot & Alternaria	4 - 8 fl oz	Use the lower rate when diseases are light, increasing rate when disease pressure increases. Apply in sufficient water for uniform, full coverage of foliage, 5 - 100 gallons per acre. Use lower volumes in early season when plants are small and increase spray volume as plant size increases. Applications should begin 6 weeks after planting or as soon as first signs of disease appear. Apply a maximum of 6 treatments during a single growing season at 7 day intervals or as needed for continued control. Do not apply within 14 days of harvest. Do not use tops as feed for livestock.
Pecans	Brown Leafspot, Powdery Mildew, Sooty Blotch, and Midrib Leaf Blotch	8 - 12 fl oz	Aerial Application: Apply in a minimum of 20 gals of water.
		8 - 12 fl oz	Ground Application: Apply in sufficient water to provide for full coverage. Diluted spray should be directed to all parts of the tree. Application should begin at pre-pollination stages when the young leaves are unfolding, and a second application made when the small nuts are forming. Apply a maximum of 10 treatments during a single growing season at 2 - 4 week intervals as needed to maintain control. Use the lower rate for the first two applications or until the disease becomes severe, or during dry weather. Use the higher rate during wet weather or during severe scab, powdery mildew or other disease infections. Do not apply after shucks have started to open.

WARRANTY STATEMENT

GRIFFIN CORPORATION warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GRIFFIN. In no case shall GRIFFIN be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. GRIFFIN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

lv

MAY 04 1988

Cello Corp.
1354 Old Post Road
Havre de Grace, MD 21078

Attention: W. Thomas Rockle

Gentlemen:

Subject: Fuldicide 14 Cleaner Disinfectant Deodoriser Fungicide
EPA Registration No. 257-262
Your Submission Dated November 19, 1987


The labeling referred to above, submitted in connection with the Label Improvement Program Notice for Antimicrobial Products Used as Food Contact Surface Sanitizers, is acceptable and a stamped copy is enclosed for your records.

However, at the printing of the next set of labels add the following misuse statement after the USE DIRECTIONS heading.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

If you have any questions concerning this letter, please call Mr. Robert Young at (703) 557-3966.

Sincerely yours,



John N. Lee
Product Manager (31)
Disinfectants Branch
Registration Division (TS-767C)

Enclosure

58284:I:Lee:LL-1:KENCO:04/29/88:05/05/88:EAE:jh

CONCURRENCES

SYMBOL	ORIGINATOR							
SURNAME								
DATE								