

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TO OPERATE: SHAKE WELL BEFORE USE. Hold sprayer 18 inches from foliage and spray with circular motion. Direct spray so that upper and lower leaf surfaces are contacted. Use caution on new growth and tender foliage. Repeat treatment as necessary. When using outdoors, spray when air is still.

This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestations. May be applied on day of harvest.

INDOORS, AS A SPACE SPRAY

To control flies, fruit flies, mosquitoes, gnats and small flying moths, apply using a good space sprayer adjusted to deliver a fine mist. Cover or remove exposed food and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of the enclosure filling the room with mist. Use 4 to 5 ounces per 1,000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

OUTDOORS, ADULT MOSQUITO CONTROL

Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest, particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant ponds and ponds. Apply at a rate of 5 to 6 quarts per acre (4 ounces per 1,000 square feet).

To control mosquito larvae, apply at the rate of 7.75 gallons per acre or 18 gallons per 1,000 square feet. Apply as a coarse wetting spray to the border or on shallow waters of wetlands, swamps and marshes where mosquito larvae is known to breed.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION Harmful if swallowed or absorbed through skin. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. Remove pets, birds and cover fish aquaria before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources.



GARDEN & PATIO SPRAY

A Concentrated Micro-Emulsion, Clear, Stable, Effective Insecticide for Use on Vegetable and Ornamental Pests.

Apply with Conventional Mechanical Compressed Air Equipment

ACTIVE INGREDIENTS

Pyrethrins 0.02 %

*Piperonyl Butoxide, Technical 0.20 %

Petroleum Distillate 0.08 %

INERT INGREDIENTS: 99.70 %

100.00 %

*Equivalent to 0.1% (butoxylate) (6-propyl piperonyl ether and 0.01% related compound)

EPA Reg. No. 257 EPA Est. No. 257-1101

Keep Out Of Reach Of Children

CAUTION

See side panel for additional precautions

NET CONTENTS: 5 GALLONS

Manufactured by *257-337*
STALFORD CONSUMER PRODUCTS DIV

a GRCY GROUP member



CELLO CORP.

HAVRE DE GRACE, MD 21078

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TO OPERATE: SHAKE WELL BEFORE USE. Hold sprayer 18 inches from foliage and spray with circular motion. Direct spray so that upper and lower leaf surfaces are contacted. Use caution on new growth and tender foliage. Repeat treatment as necessary. When using outdoors, spray when air is still.

This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestations. May be applied on day of harvest.

INDOORS, AS A SPACE SPRAY

To control flies, fruit flies, mosquitoes, gnats and small flying moths, apply using a good space sprayer adjusted to deliver a fine mist. Cover or remove exposed food and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of the enclosure filling the room with mist. Use 4 to 5 ounces per 1,000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

OUTDOORS, ADULT MOSQUITO CONTROL

Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest, particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant ponds and ponds. Apply at a rate of 5 to 6 quarts per acre (4 ounces per 1,000 square feet).

To control mosquito larvae, apply at the rate of 7.75 gallons per acre or 18 gallons per 1,000 square feet. Apply as a coarse wetting spray to the border or on shallow waters of wetlands, swamps and marshes where mosquito larvae is known to breed.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION Harmful if swallowed or absorbed through skin. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. Remove pets, birds and cover fish aquaria before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources.

STORAGE & DISPOSAL

STORAGE Keep from freezing. If freezing occurs, thaw and agitate vigorously. Store in a cool dry area away from heat or open flame.

DISPOSAL Do not reuse empty container. Wrap container and dispose in trash collection.

Buyer assumes all risks of use, storage or handling of this product in strict accordance with directions given herewith.

44 03

<p>U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDE REGISTRATION REGISTRATION DIVISION, WASHINGTON, D.C. 20460</p> <p>NOTICE OF PESTICIDE.</p> <p><i>Unlabeled Use of Pesticide</i></p>	<p>REGISTRATION NUMBER</p> <p>DATE OF ISSUANCE</p>	<p>DATE OF ISSUANCE</p>
<p>NAME AND ADDRESS OF REGISTRANT (SEE ZIP CODE)</p> <p style="text-align: right;">92906</p>		
<p>NOTE: Changes in label formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p>		
<p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p>		
<p>Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, in his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p>		
<p><input type="checkbox"/> ATTACHMENT IS APPLICABLE</p>		
<p>SIGNATURE OF APPROVING OFFICIAL</p>	<p>DATE</p>	