

NET CONTENTS

# Weed Wilt

## Liquid Concentrate

Active Ingredients  
Dimethylamine salt of 2, 4-dichlorophenoxyacetic acid \_\_\_\_\_ 32.6%  
(Equivalent to 2, 4-dichlorophenoxyacetic acid—27% or 2.7 lbs. per gallon.)  
Inert Ingredients \_\_\_\_\_ 67.4%  
USDA Registration No. 259-6

For use in killing annual and perennial weeds in established turf, lawns, golf courses, pastures. (Do not graze treated areas by dairy animals within 7 days after treatment), fence rows, right-of-ways, ditch banks.

For killing such weeds as Canada Thistle, Perennial Sow Thistle, Bindweed, White Top, Russian Knapweed, Cornet Weed, Cockspur, Lamb's Quarters, Mustards, Pigweed, Wild Morning Glory, and other broad leaf weeds.

### CAUTION: Keep Out of Reach of Children

Avoid Contamination of Food. Harmful if Swallowed.

Avoid Contact with Skin, Eyes or clothing.

### DIRECTIONS FOR USE


Dilute MKC WEED WILT, approximately 1 oz. to two gallons of water. Use two gallons of solution per 1,000 square feet, approximately. Best results are obtained by treatment when weeds are growing vigorously. Apply during periods of active growth in Spring and Fall. Some hard-to-kill weeds may not be controlled by one application, but top growth and reseeding usually will be prevented. Repeated treatments generally give good control. Dosage of Weed Wilt should be in the range of 1 to 2 pounds per acre.

### CAUTION

Do not apply to newly seeded turf or pastures. Under no circumstances should this herbicide product or any 2,4-D or 2,4,5-T weed killer be used in the vicinity of cotton, tomatoes, garden crops, grapes, ornamentals, or other susceptible crops, or severe damage may result. Do not apply on a windy day. Extreme care must be exercised to prevent drifting during application. (Coarse sprays are less likely to drift.)

Do not re-use this container. Do not use equipment used in applying this product or any 2,4-D or 2,4,5-T weed killer to apply insecticides, fungicides, or other material to susceptible crops. Do not store near fertilizers, seeds, insecticides, fungicides, or foodstuffs.

Application to grassy areas may seriously damage or destroy clovers or bent grasses if they are present. Avoid contamination of water supplies that may be used to irrigate or water susceptible crops, or used for domestic purposes. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops.



# MOKAN

missouri • kansas chemical co.



# Weed Wilt

## Liquid Concentrate

Active Ingredient: Dimethylamine salt of 2, 4-dichlorophenoxyacetic acid 32.6%  
 (Equivalent to 2, 4-dichlorophenoxyacetic acid—27% or 2.7 lbs. per gallon.)  
 Inert Ingredients 67.4%

USDA Registration No. 259-6

For use in killing annual and perennial weeds in established turf, lawns, golf courses  
 pastures (Do not graze treated areas by dairy animals within 7 days after treatment), fence  
 rows, right-of-ways, ditch banks.

For killing such weeds as Canada Thistle, Perennial Sow Thistle, Bindweed, White Top,  
 Russian Knotweed, Carpet Weed, Cockspur, Lamb's Quarters, Mustards, Pigweed, Wild  
 Morning Glory, and other broad leaf weeds.

**CAUTION: Keep Out of Reach of Children**  
 Avoid Contamination of Feed. Harmful if Swallowed.  
 Avoid Contact with Skin, Eyes or Clothing.

### DIRECTIONS FOR USE

Dilute MKC WEED WILT, approximately 1 oz. to two gallons of water. Use two gallons  
 of solution per 1,000 square feet, approximately. Best results are obtained by treatment  
 when weeds are growing vigorously. Apply during periods of active growth in Spring and  
 Fall. Some hard-to-kill weeds may not be controlled by one application, but top growth  
 and reseeding usually will be prevented. Repeated treatments generally give good control.  
 Dosage of Weed Wilt should be in the range of 1 to 2 pounds per acre.

### CAUTION

Do not apply to newly seeded turf or pastures.  
 Under no circumstances should this herbicide product or any 2,4-D or 2,4,5-T weed  
 killer be used in the vicinity of cotton, tomatoes, garden crops, grapes, ornamentals, or  
 other susceptible crops, or severe damage may result. Do not apply on a windy day.  
 Extreme care must be exercised to prevent drifting during application (Coarse sprays are  
 less likely to drift).

Do not re-use this container. Do not use equipment used in applying this product or  
 any 2,4-D or 2,4,5-T weed killer to apply insecticides, fungicides, or other material to  
 susceptible crops. Do not store near fertilizers, seeds, insecticides, fungicides, or foodstuffs.

Application to grassy areas may seriously damage or destroy clovers or bent grasses  
 if they are present. Avoid contamination of water supplies that may be used to irrigate or  
 water susceptible crops, or used for domestic purposes. Misuse as to quantity, timing or  
 method of application can cause damage or injury to animals, persons, property or crops.

CONTAINED

**MOKAN**

missouri • kansas chemical co.