

ACCEPTED

259-51

OCT 9 1974

Net Contents
1 Gallon

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROENTICIDE ACT
OR ECONOMIC POISON REGISTER
UNDER 259-51

Minto-Cide II



Manufactured Only by **MOKAN**

missouri • kansas chemical co.

1708-16 CAMPBELL STREET • KANSAS CITY, MISSOURI 64108

Minto-Cide

BACTERICIDE AND DISINFECTANT

Active Ingredients—Total	37.5%
Isopropanol	20.0%
Soap	5.5%
o-benzyl-p-chlorophenol	4.5%
o-phenylphenol	4.0%
Methyl Salicylate	2.0%
Para-tertiary-amyphenol	1.5%
Inert Ingredient	
Water	62.5%

Phenol Coefficient 3 A.O.A.C.
EPA Registration No. 259-51 EPA EST. 259-MC 1

FOR HOSPITALS

DIRECTIONS FOR USE

For the SICK ROOM: All utensils, such as bedpans and urinals should be pre-cleaned before disinfecting. For disinfecting, wash with a solution of one ounce of this disinfectant to one-half gallon of water.

WATER CLOSETS and URINALS: Use 3 ounces of this disinfectant to a gallon of water. Apply freely to all surfaces to be disinfected and flush the drain pipes with the solution for deodorization.

This disinfectant is highly recommended for general hospital sanitation: Floors, walls, and for the cleaning and disinfecting of sick-room utensils.

GENERAL DISINFECTING

For Floors, Walls, Tables, Counters, Bed Frames, Window and Door Ledges, Venetian Blinds, Desks, Telephones, Toilet and Bathroom Fixtures.

Remove dirt from the premises and use a solution of two parts of this disinfectant to 200 parts of water.

As a disinfectant cleaning solution, use two parts of this disinfectant mixed with 70 parts of water.

To deodorize Water Closets and Urinals. Use 3 ounces of this disinfectant to a gallon of water. Apply freely to all surfaces to be deodorized and flush the drain pipes with the solution.

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

Rinse empty container thoroughly with water and discard it. Do not use, pour, spill or store near heat or open flame.