

GENERAL INFORMATION

WEEDONE LV4 is a 2,4-D herbicide for use on broadleaf weeds in crops and non-crop areas. It is a low volatility formulation. For more information on this product, contact your local Union Carbide Agricultural Products Company representative. **NOTE:** Do not use on crops or areas where the crop is to be harvested within 30 days of application. **NOTICE:** Do not use on areas where the crop is to be harvested within 30 days of application.

RECOMMENDATIONS FOR THE APPLICATION OF WEEDONE LV4 Broadleaf Postemergence Herbicide

WEEDS IN CROPS	PH/5 WEEDONE LV4 PER ACRE	DIRECTIONS
Wheat, Barley, Rye, Winter Grain Annual and biennial weeds Perennial broadleaf weeds	1.0 pint 1.0 pint	Apply when weeds are 2 to 4 inches tall. Apply when weeds are 2 to 4 inches tall.
Spring Grain Annual broadleaf weeds Perennial broadleaf weeds	1.0 pint 1.0 pint	Apply after grain has emerged. Apply when weeds are 2 to 4 inches tall.
Wild onion or garlic	1.0 pint	Apply when weeds are 2 to 4 inches tall. Apply when weeds are 2 to 4 inches tall.
Emergency Weed Control in Wheat Perennial broadleaf weeds	1.0 pint	Apply when weeds are 2 to 4 inches tall. Apply when weeds are 2 to 4 inches tall.
Field Corn Pre-emergence	1.0 pint	To control broadleaf weeds and suppress annual grasses, apply 3 to 5 days after planting but before corn emergence. Use higher rate in soil with high moisture or when soil moisture is inadequate for normal weed growth.
Post-emergence Annual broadleaf weeds	1.0 pint	Apply when weeds are 2 to 4 inches tall. Apply when weeds are 2 to 4 inches tall.
Perennial broadleaf weeds	1.0 pint	Spray when weeds are in the bud to break stage. Do not spray when weeds are in the bud to break stage. Do not spray when weeds are in the bud to break stage.
Grain Sorghum Annual broadleaf weeds Perennial broadleaf weeds	1.0 pint 1.0 pint	Treat when weeds are 2 to 4 inches tall. Do not spray on treated grain fields within 30 days after treatment with this product. Do not feed treated straw to livestock.

For aerial application on grain crops, WEEDONE LV4 Broadleaf Postemergence Herbicide should be applied at a rate of 1.0 pint per acre. Do not mix with oil for crop uses. Do not graze or graze grain fields within two weeks after treatment with this product. Do not feed treated straw to livestock.

WEEDS IN NON-CROP AREAS (lawns, roadsides, drainage ditch-banks)

Annual broadleaf weeds	1.0 pint	Apply when weeds are 2 to 4 inches tall.
Perennial broadleaf weeds	1.0 pint	Apply when weeds are 2 to 4 inches tall.
Gutters and Tules	1.0 pint	Apply when weeds are 2 to 4 inches tall.

For aerial application to lawns, roadsides, drainage ditch-banks, use 1.0 pint per acre. For ground application to lawns, roadsides, drainage ditch-banks, use 1.0 pint per acre.

WEEDS IN PASTURES, OTHER TURF AREAS (parks, golf courses, cemeteries)


Annual broadleaf weeds	1.0 pint	Apply in spring or fall when weeds are 2 to 4 inches tall.
Perennial broadleaf weeds	1.0 pint	Apply in spring or fall when weeds are 2 to 4 inches tall.

Apply to newly seeded areas, golf courses, and pastures. Do not graze or graze pastures within two weeks after treatment with this product. Do not feed treated straw to livestock.

Uses in Forest Management
CONIFER RELEASE: WEEDONE LV4 Broadleaf Postemergence Herbicide is used to control broadleaf weeds in conifer plantations. It is applied in the summer months when weeds are 2 to 4 inches tall. Do not apply to areas where conifers are to be planted within 30 days of application. Do not graze or graze pastures within two weeks after treatment with this product. Do not feed treated straw to livestock.

SITE PREPARATION: Broadcast Spray
Apply when weeds are 2 to 4 inches tall. Do not apply to areas where conifers are to be planted within 30 days of application.

Emergence Spray
Apply when weeds are 2 to 4 inches tall. Do not apply to areas where conifers are to be planted within 30 days of application.



BROADLEAF POSTEMERGENCE

The 2,4-D Low Volatile Formulation
Agricultural Weed Control
Conifer Release and Site

CAUTION: Keep out of reach of children on right panel.


ACTIVE INGREDIENT:
2,4-Dichlorophenoxyacetic acid, butoxyethyl
INERT INGREDIENTS:
Isomer specific by AOAC method No. 6
*2,4-Dichlorophenoxyacetic acid equivalent
3.8 pounds per gallon.

UNION CARBIDE AGRICULTURAL PRODUCTS COMPANY

AMBLER, PA. 19002

MADE IN U.S.A.

WEEDONE



POSTEMERGENCE HERBICIDE

Low Volatile Ester for
Post-emergence Weed Control,
Pre-planting and Site Preparation

Keep out of reach of children. See other cautions

CONTENTS:
2,4-Dichlorophenoxyacetic acid, butoxyethyl ester* 60.8%
2,4-Dichlorophenoxyacetic acid 39.2%
* by AOAC method No. 6 D01-5
2,4-Dichlorophenoxyacetic acid equivalent 41.9% by weight or
per gallon.

MADE IN U.S.A.

2,4-DICHLOROPHENOXYACETIC ACID
AGRICULTURAL
HERBICIDE
AMBIER COMPANY, INC.

AMBIER, PA 19002

St. Joseph, MO Fremont, CA

PRECAUTIONARY STATEMENTS Hazard to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Avoid breathing vapors or spray mist. Remove and wash contaminated clothing before reuse. Wash thoroughly before eating or smoking.

Statement of Practical Treatment

In case of contact wash skin with soap and water. For eyes flush with water for at least 15 minutes and get medical attention. Never give anything by mouth to an unconscious person. Get medical attention.

Environmental Hazards

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply when weather conditions favor drift from treated areas.

Do not use the same spray equipment for other purposes unless thoroughly cleaned.

Do not use in or near a greenhouse.

Do not contaminate water used for irrigation or domestic purposes.

Do not apply WEEDONE LV4 Broadleaf Postemergence Herbicide directly to or permit spray mist to drift onto cotton, okra, grapes, tomatoes, fruit trees, vegetables, flowers or other desirable crop or ornamental plants which are susceptible to 2,4-D herbicide. Do not apply near susceptible plants since very small quantities of the 2,4-D will cause severe injury during the growing or dormant periods. Crops contacted by WEEDONE LV4 Broadleaf Postemergence Herbicide sprays or spray drift may be killed or suffer significant stand loss with extensive quality and yield reduction.

Do not apply when a temperature air inversion exists. Such a condition is characterized by little or no air movement and an increase in air temperature with an increase in height. In humid regions a fog or mist may form. An inversion may be detected by producing a smoke column and checking for a layering effect. If questions exist pertaining to the existence of an inversion consult with local weather services before making an application.

Use coarse nozzles to minimize drift. Do not apply with hollow cone type insecticide or other nozzles that produce fine spray droplets.

Drift from aerial or ground application may be reduced by (1) applying as near to the target as possible in order to obtain coverage, (2) by increasing the volume of spray mix per acre, (3) by decreasing the poundage of pressure at the nozzle tips, (4) by using nozzles which produce a coarse spray pattern, (5) by not applying when wind is blowing toward susceptible crops or valuable plants.

At high air or ground surface temperatures vapors from this product may injure susceptible plants.

Storage and Disposal Statements

Do not contaminate or store near food, feedstuff, fertilizers, seeds, insecticides, fungicides or other pesticides. Do not burn. Open dumping is prohibited.

Container Disposal

Triple rinse and offer for recycling, reconditioning or dispose of in an approved landfill area. Puncture and bury in a safe place in non-croplands away from water supplies.

Local conditions may affect the use of herbicides. Consult your State Agriculture Experiment Station, Farm Advisors or Extension Weed Specialists for advice in selecting treatment from this label to best fit local conditions.

Be sure that use of this product conforms to all applicable laws, rules and regulations. Certain states have restrictions pertaining to application distances from susceptible crops. The applicator should become familiar with these laws, rules or regulations and follow them exactly. Apply this product only as specified on this label.

WARRANTY

The manufacturer warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. The manufacturer therefore makes no other express or implied warranty, and no agent of the manufacturer is authorized to do so. Buyer agrees, in purchasing this product, to assume the risks and in the event of damages arising from a breach of the warranty to accept refund of the purchase price of the product or to bear the charge of the manufacturer's liability.