

DIRECTIONS FOR USE


CROP	Amount WEEDAR MCPA per Acre	DIRECTIONS
FLAX	2/3 - 3/4 pint	Where susceptible weeds such as mustard are the main problem.
	1 pint	Where weeds such as pigweed, lambsquarters, stinkweed, ragweed, cocklebur are the problem. Apply when flax is 2 to 8 inches tall but before the bud stage. Do not spray after flax is in the bud stage.
SMALL GRAINS (wheat, oats, barley, rye) Spring planted (not seeded to legumes)	1 pint	Treat in spring when grain is fully tillered and 8 to 10 inches tall. Do not spray grain in the boot to dough stage. Use about 8 gallons of water per acre.
	1 pint	Apply in early spring when annual broadleaf weeds are small and grain is fully tillered, but before grain is in the boot stage. Use about 8 gallons of water per acre.
SMALL GRAINS Underseeded with mixed legumes (alfalfa, alsike, red and ladino clover, trefoil)	1 pint	Treat in early spring when grain is in the tiller to boot stage and legume seedlings are 2 to 3 inches tall. Do not spray when grain is in the boot to dough stage. Use 5-6 gallons of water per acre since higher volumes may result in injury to legumes.
OATS	1 to 2 pints	Spray after oats are fully tillered (4 to 6 inches tall) but before they are in the boot to dough stage. Apply the lower rate where annual weeds are the major problem, the higher rate for perennial weeds. Use 8-10 gallons of water per acre.
RICE	3 to 5 pints	Treat 7 to 10 weeks after planting when rice is fully tillered and 6 to 8 inches above water. Do not spray after rice is in the boot stage. Use the higher rate where curly indigo, sedge or bulrush are the major problems. Use lower rate for arrow head, water plantain and redstem. Do not spray when temperatures are over 90°F.
PEAS	1 to 1½ pints	Treat when peas are 4 to 6 inches tall using 15 to 30 gallons of water per acre. Peas may show slight injury but they usually recover a few days after treatment. Use the higher rate before buds form on Canada thistle to prevent thistle from going to seed. Do not spray peas that are under stress from lack of moisture or when temperatures are over 90°F. Do not forage or graze treated pea fields. Do not feed treated vines to livestock.

To convert local recommendations into terms of WEEDAR MCPA, use the following table:

MCPA acid	1¼ lb. (20 oz.)	1 lb. (16 oz.)	¾ lb. (12 oz.)	½ lb. (8 oz.)	3/8 lb. (6 oz.)	¼ lb. (4 oz.)
WEEDAR MCPA	5 pt.	4 pt.	3 pt.	2 pt.	1½ pt.	1 pt.

WEEDAR[®]

MCPA



Kills weeds in flax, rice, oats, peas, and small grains.

CAUTION: Keep out of reach of children.

See other cautions on right side panel.

ACTIVE INGREDIENT:

*Dimethylamine salt of 2-methyl-4-chlorophenoxyacetic acid . . . 27.6%

INERT INGREDIENTS: 72.4%

*2-methyl-4-chlorophenoxyacetic acid equivalent 22.5%
by weight or two pounds per gallon.




AMCHEM PRODUCTS, INC

First Name in Herbicide Research
AMBLER, PA ■ Clinton, Iowa ■ St. Joseph, Mo ■ Fairport, N.Y.

EPA Reg. No. 264-1-1A

WEEDAR[®]

MCPA



Kills weeds in flax, rice, oats, peas, and small grains.

CAUTION: Keep out of reach of children.
See other cautions on right side panel.

ACTIVE INGREDIENT:

*Dimethylamine salt of 2-methyl-4-chlorophenoxyacetic acid . . . 27.6%

INERT INGREDIENTS: 72.4%

*2-methyl-4-chlorophenoxyacetic acid equivalent 22.5%
by weight or two pounds per gallon.



AMCHEM PRODUCTS, INC.

First Name in Herbicide Research
AMBLER, PA ■ Clinton, Iowa ■ St. Joseph, Mo ■ Fremont, Calif

EPA Reg. No. 264-16 AA

GENERAL INFORMATION

Tests in this country and in Europe show that several crops such as flax, rice, and small grains underseeded with legumes are more tolerant of MCPA than they are of 2,4-D. WEEDAR MCPA is effective on a large number of broadleaf weeds (mustard, thistle, and pigweed for example) and is useful for controlling these weeds in certain crops. Use MCPA on oats because oats are more tolerant of MCPA than they are of 2,4-D.

WEEDAR MCPA is an amine form of MCPA that is easily dissolved in water, even in the hard waters found in some parts of this country. WEEDAR MCPA can be sprayed in either high or low volumes of water, according to the recommendations of local agricultural authorities. Mix thoroughly while diluting this material and mix again if allowed to stand long before spraying.

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, or clothing.

Avoid spray drift to susceptible plants such as cotton, beans, peas, and ornamentals, as injury may result. (Coarse sprays are less likely to drift.) Do not use same sprayer for other purposes. Do not store near fertilizers, seeds, insecticides, or fungicides. Under very high temperature of use, vapors from this product may injure susceptible plants in the immediate vicinity. Do not contaminate water used for irrigation or domestic purposes. Do not use in a greenhouse.

Do not reuse container. Destroy in a safe place.

Do not use this product for purposes other than those recommended on this label.

WARRANTY

Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. Amchem, therefore, makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees, in purchasing this product, to assume the risks and in the event of damages arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of Amchem's liability.