

DIRECTIONS

WEEDONE 638 is a special formulation containing 2,4-D acid. WEEDONE 638 is more effective than 2,4-D amines for controlling hard-to-kill weeds such as field bindweed (morningglory), Russian knapweed, Canada thistle, leafy spurge, cattails, tules and nutgrass.

ROADSIDES, FENCEROWS, DRAINAGE DITCH BANKS

Spray when daytime temperatures are lower than 80 to 85 degrees. Spraying when temperatures are higher may cause rapid foliage burn and decrease the effectiveness of WEEDONE 638. Treat at the stage of growth recommended below. Wet weeds thoroughly, applying about 150 to 300 gallons of spray mixture per acre, depending on the stand of weeds and spray equipment used. Where necessary, repeat applications should be made on active regrowth when it develops.

WEEDS	quarts WEEDONE 638 per acre	WHEN TO SPRAY
field bindweed (morningglory), hoary cress,	2	bud to full bloom
leafy spurge, Russian knapweed	2	full bloom
Canada thistle, camellthorn, lotus	2	bud to early bloom
cattails, tules	3	late spring to early summer when 4 to 6 feet tall; repeat in fall and for next spring
tansy ragwort	3	bud to early bloom
nutgrass	3	6 to 8 inches high, before bloom; repeat when re- growth is 6 to 8 inches tall

PASTURES, FALLOW AREAS, TURF

	Amounts per Acre		REMARKS
	quarts WEEDONE 638	gallons WATER	
field bindweed (morningglory)	1½	50	for each spraying*
	1½	10	for annual spraying*

Fallow land treated with WEEDONE 638 in the spring or summer can be returned to crops in the fall or following spring. Do not graze livestock for seven days after spraying pastures with this product. Do not use on newly seeded turf until grass has been cut several times. WEEDONE 638 may injure bentgrass, St. Augustine grass, Dichondra, carpetgrass, and clover lawns.

CROPPED AREAS

CROP	Amount WEEDONE 638 per acre	WHEN TO SPRAY
Wheat*, Barley		
annual broadleaf weeds	1 pint	Apply after grains are fully tillered (4 to 6 inches tall) but before they are in the boot stage.
perennial broadleaf weeds	2 pints	Apply when weeds are in the bud stage if possible, but do not spray after grains are in the boot to dough stage. *Treat fall-planted wheat in the spring after grains are fully tillered but before they are in the boot stage.
Grain Sorghum		
annual broadleaf weeds	1 pint	Treat once sorghum is 6 to 10 inches tall.
perennial broadleaf weeds	1½ pints	

Do not top-dress or graze beef or dairy fields within two weeks after treatment. Do not use straw or hay from treated fields as bedding for livestock.

Among the weeds controlled are

Annuals: lambsquarters, pigweed, mustard, kochia and Russian thistle.

Perennials: bindweed (morningglory), Canada thistle, blueweed, bur ragweed, alkali mallow.

Use low pressure sprayers (25 to 40 psi) and apply a minimum of 5 to 10 gallons of water per acre.

WEEDONE

638 BROADLEAF POST- EMERGENCE HERBICIDE

ACCEPTED

MAY 14 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 149-1475

A special formulation containing 2,4-D acid.

CAUTION: Keep out of reach of children.
See other cautions on right panel.

ACTIVE INGREDIENTS:

*2,4 Dichlorophenoxyacetic acid 14.8%

*Butoxyethanol ester of 2,4-dichlorophenoxyacetic acid 26.0%

INERT INGREDIENTS: 59.2%

*Total 2,4-dichlorophenoxyacetic acid equivalent 32.8% by weight
or 3.0 pounds per gallon.



AMCHEM

PRODUCTS, INC.

10000 W. 10th Street, Omaha, Nebraska 68144

10000 W. 10th Street, Omaha, Nebraska 68144

EPA Reg. No. 149-1475 37

PA 149-1475 1