

## DIRECTIONS

WEEDONE 638 broadleaf herbicide is a special formulation containing 2,4-D acid. WEEDONE 638 is more effective than 2,4-D amines for controlling hard-to-kill weeds such as field bindweed (morningglory), Russian knapweed, Canada thistle, leafy spurge, cattails, tules and nutgrass.

### ROADSIDES, FENCEROWS, DRAINAGE DITCH BANKS

Spray when daytime temperatures are lower than 80 to 85 degrees. Spraying when temperatures are higher may cause rapid foliage burn and decrease the effectiveness of WEEDONE 638 broadleaf herbicide. Treat at the stage of growth recommended below. Wet weeds thoroughly, applying about 150 to 300 gallons of spray mixture per acre, depending on the stand of weeds and spray equipment used. Where necessary, repeat applications should be made on active regrowth when it develops.

WEEDS	quarts WEEDONE 638 per acre	WHEN TO SPRAY
field bindweed (morningglory), hoary cress	2	bud to full bloom
leafy spurge, Russian knapweed	2	full bloom
Canada thistle, cornelthorn, lotus	2	bud to early bloom
cattails, tules	3	late spring to early summer when 4 to 5 feet tall, repeat in fall and following spring
tansy ragwort	3	rosette stage
nutgrass	3	6 to 8 inches high, before bloom, repeat when regrowth is 6 to 8 inches high

### PASTURES, FALLOW AREAS, TURF

	Amounts per Acre		REMARKS
	quarts WEEDONE 638	gallons WATER	
field bindweed (morningglory)	1½	50	for boom spraying*
	1½	10	for aerial spraying*
			*Treat during early to full bloom

Fallow land treated with WEEDONE 638 broadleaf herbicide in the spring or early summer can be returned to crops in the fall or following spring. Do not graze livestock for seven days after spraying pastures with this product. Do not use on newly seeded turf until grass has been cut several times. WEEDONE 638 broadleaf herbicide may injure bentgrass, St. Augustine grass, Dichondra, carpetgrass, and clover lawns.

### CROPPED AREAS

CROP	Amount WEEDONE 638 per acre	WHEN TO SPRAY
Wheat,* Barley		
annual broadleaf weeds	1 pint	Apply after grains are fully tillered (4 to 6 inches tall) but before they are in the boot stage.
perennial broadleaf weeds	2 pints	Apply when weeds are in the bud stage if possible, but do not spray after grains are in the boot to dough stage. *Treat fall planted wheat in the spring after grains are fully tillered but before they are in the boot stage.
Grain Sorghum		
annual broadleaf weeds	1 pint	Treat when sorghum is 6 to 10 inches tall.
perennial broadleaf weeds	1½ pints	

Do not forage or graze treated grain fields within two weeks after treatment with this product. Do not feed treated straw to livestock.

Among the weeds controlled are:

Annuals: Lambsquarters, pigweed, mustard, Kochia and Russian thistle.

Perennials: Field bindweed (morningglory), Canada thistle, blueweed, bur, ragweed, alkali marshes.

Use for: pressure sprayers (25 to 40 psi) and apply in a minimum of 5 to 10 gallons of water per acre.

# WEEDONE<sup>®</sup>

## 638 BROADLEAF HERBICIDE

A special formulation containing  
2,4-D acid.

**CAUTION:** Keep out of reach of children.  
See other cautions on right panel.

### ACTIVE INGREDIENTS:

\*2,4-dichlorophenoxyacetic acid .....  
\*2,4-dichlorophenoxyacetic acid, butoxyethyl ester .....

### INERT INGREDIENTS: .....

\*Isomer specific by AOAC method No. 6.D01-5

\*Total 2,4-dichlorophenoxyacetic acid equivalent 30  
or 2.8 pounds per gallon.

IN CASE OF EMERGENCY, TELEPHONE COLLECT  
(24 HOURS A DAY) IN THE USA (304) 744-3427  
Made in U.S.A.

AMERICAN CARBIDE AGRICULTURAL PRODUCTS CO.

AMBLER, PA. 19002


Clinton, IA. ■ St. Joseph, MO. ■ Fremont, CA

265 EPA Reg. No. 264-37

EPA Est. 26

# WEEDONE®

## BROADLEAF HERBICIDE



Formulation containing

Keep out of reach of children.  
See instructions on right panel.

3: oxyacetic acid ..... 13.8%  
oxyacetic acid, butoxyethyl ester ..... 24.5%  
..... 61.7%

by AOAC method No. 6.D01-5

phenoxyacetic acid equivalent 30.8% by weight  
gallon.

OF EMERGENCY, TELEPHONE COLLECT  
24 HOURS A DAY) IN THE USA (304) 744-3487  
Made in U.S.A.

AGRICULTURAL PRODUCTS COMPANY, INC.  
AMBLER, PA 19002

A. ■ St. Joseph, MO ■ Fremont, CA

64-37

EPA Est. 264-PA-1

### GENERAL INFORMATION

#### WEED CONTROL IN CORN

One thorough shallow cultivation is recommended when corn is 10-12 inches high.

Apply spray when corn is 18 to 24 inches high.

Use drop pipes to direct spray away from corn leaves and onto weeds at the base of the corn.

Use 1 pint WEEDONE 638 broadleaf herbicide in 30 gallons of water for each acre sprayed.

2,4-D may cause brittleness to corn. Winds or cultivations may cause stalk breakage while brittle.

Certain single cross corn hybrids may be more susceptible to 2,4-D injury than others.

#### Combination with nitrogen solutions:

Use 14 fluid ounces (7/8 pint) WEEDONE 638 broadleaf herbicide per acre, mixed with the amount of nitrogen solution required to apply the pounds of nitrogen you want per acre.

Pour liquid nitrogen into the spray tank, then add WEEDONE 638 broadleaf herbicide. Mix thoroughly before spraying and maintain good by-pass agitation while spraying.

#### MIXING SPRAYS

Add the required amount of water to the spray tank. Start agitation. While agitating, slowly add WEEDONE 638 broadleaf herbicide. Mix thoroughly. Whenever practical, maintain continuous agitation and use spray material as soon as possible after mixing.

### PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

#### CAUTION

Harmful if swallowed. Avoid contact with skin, eyes or clothing.

#### Statement of Practical Treatment

In case of contact wash skin with soap and water for eyes flush with water for at least 15 minutes and get medical attention.

#### Environmental Hazards

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not contaminate water used for irrigation or domestic purposes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Avoid spray drift to susceptible plants such as cotton, beans, peas, ornamentals, and most vegetables, as injury may occur. Coarse sprays are less likely to drift.

Do not store near fertilizers, seeds, insecticides, or fungicides. Under very high temperature vapors from this product may injure susceptible plants in the immediate vicinity. Do not use in or near a greenhouse.

Do not use this product for purposes other than those recommended on this label.

### STORAGE AND DISPOSAL STATEMENT

Do not store near fertilizers, seeds, insecticides, or fungicides. Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixtures, or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place. Do not burn. Consult Federal, state or local disposal authorities for approved alternate procedures.

### LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants that this product conforms to the chemical description on the label and that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions, and that the directions, warnings and other statements on the label are based upon responsible experts' evaluation of reasonable tests of effectiveness of toxicity to laboratory animals and to plants, and of residue, and upon reports of field experience. Tests have not been made in all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

This warranty does not extend to, and the Buyer shall be solely responsible for, any and all loss or damage which results from the use of this product in any manner which is inconsistent with the label, directions, warnings or cautions.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT SUCH CLAIMS, DAMAGES, OR INJURIES ARE CAUSED IN WHOLE OR IN PART BY THE NEGLIGENCE OF THE MANUFACTURER OR SELLER, SHALL BE LIMITED TO THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITHIN THE RESPECTIVE WHICH CLAIMS ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

200