


pm 23

264-47

1 of 13

WEEDAR®  
 MCPA Concentrate  
 Broadleaf Postemergence Herbicide



Controls weeds in flax, rice, peas, and small grains

ACTIVE INGREDIENT:

Dimethylamine salt of 2-methyl-4-chlorophenoxyacetic acid\* . . . . . 52.1%  
 INERT INGREDIENTS: . . . . . 47.9%

\*2-Methyl-4-chlorophenoxyacetic acid equivalent  
42.6% by weight or four pounds per gallon.

E.P.A. Reg. No. 264-47 AA

E.P.A. Est. No. 264-MO-01

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

PRACTICAL TREATMENT

IF IN EYES: Flush with water for at least 15 minutes and get medical attention.

PRECAUTIONARY STATEMENTS: See inside booklet

PRECAUCION AL USARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

For Emergency Information ONLY Call 24 Hours A Day 1-800-521-8235

For Product Handling, Use and General Information Call 1-800-334-9745

RHONE-POULENC AG COMPANY  
 P. O. Box 12014 T. W. Alexander Drive  
 Research Triangle Park, N. C. 27709

WEEDAR® is a registered trademark of RHONE-POULENC NEDERLAND B.V.  
Made in U.S.A.

5 of 13

## PRECAUTIONARY STATEMENTS

### DANGER

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage. Harmful if swallowed. May be fatal if absorbed through skin. Avoid contact with eyes, skin or clothing. Avoid breathing vapors or spray mist. Remove and wash contaminated clothing before reuse. Wash thoroughly before eating or smoking.

#### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: If patient is conscious and alert, give 2-3 glasses of water to drink. Do not induce vomiting. Get medical attention.

IF ON SKIN: Wash skin with plenty of soap and water, if available, while removing contaminated clothing and shoes. Wash clothing separately before reuse.

IF IN EYES: Hold eyelids open and flush with a steady gentle stream of water for 15 minutes. Get medical attention preferably an ophthalmologist.

If INHALED: Remove victim to fresh air. If not breathing, give artificial respiration. Administer oxygen if necessary. GET MEDICAL ATTENTION.

#### NOTE TO PHYSICIAN

No specific antidote is available.



All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

This product is irritating to mucous membranes. If large amounts (greater than 1 ml/kg body weight) of the product have been ingested, the stomach should be evacuated by gastric intubation with the aid of a cuffed endotracheal tube to prevent exposure of the esophagus. After removal of stomach contents, wash stomach by instilling 30-50 g. of activated charcoal in 3-4 ounces of water through the stomach tube and again remove stomach contents. Avoid oily laxatives.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water except as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes.

Avoid spray drift to susceptible plants such as cotton, beans tomatoes, and ornamentals. (coarse sprays are less likely to drift).

Do not apply when weather conditions favor drift from treated areas. Avoid use of small diameter spray nozzles. At high air or ground surface temperatures, vapors from this product may injure susceptible plants.

Do not contaminate water used for irrigation or domestic purposes.

Do not use the same spray equipment for other purposes unless thoroughly cleaned.

Do not use in or near a greenhouse.

Do not apply WEEDAR® MCPA Concentrate through any type of irrigation system.

GENERAL WORKER PROTECTION STATEMENTS

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in treated area or in an area about to be treated with this product. Advise workers to stay out of fields during application and until sprays have dried. Regular long-sleeved work clothing should be worn when working in treated fields. See PRECAUTIONARY STATEMENTS, STATEMENT OF PRACTICAL TREATMENT and NOTE TO PHYSICIAN for information on accidental exposures. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: Appropriate signal word (DANGER), area treated with WEEDAR® MCPA Concentrate Broadleaf Postemergence Herbicide, date of application, appropriate protective clothing, and reentry interval (i.e. until sprays have dried).



DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply WEEDAR MCPA Concentrate through any type of irrigation system.

GENERAL INFORMATION

WEEDAR® MCPA Concentrate Broadleaf Postemergence Herbicide is effective on a large number of broadleaf weeds and is useful for controlling these weeds in certain crops. Several crops, such as flax, rice, oats and small grains underseeded to legumes, are more tolerant of MCPA than they are of 2,4-D.

WEEDAR® MCPA Concentrate Broadleaf Postemergence Herbicide is the amine form of MCPA that is easily dissolved in water, even in the hard waters found in some areas. Mix thoroughly while diluting this material and mix again if allowed to stand long before spraying.

COMPATIBILITY: If WEEDAR MCPA Concentrate is to be tank mixed with other pesticides, compatibility should be tested prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 qt) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5-15 minutes after mixing. Read and follow all directions and precautions on this label and on the labels of any products for which a tank mixture is being considered.

Weeds controlled by properly applied treatments of WEEDAR MCPA Concentrate include the following:

- |                                |                             |
|--------------------------------|-----------------------------|
| Amaranthus species (pigweeds)  | plantains                   |
| common burdock                 | purslanes                   |
| common cocklebur               | ragweed (common and giant)  |
| common lambsquarters           | shepherdspurse              |
| dandelion                      | thistles (bull and Russian) |
| field pennycress               | yellow rocket               |
| mustards (except blue mustard) |                             |
| pepperweeds (except perennial) |                             |



To convert locally recommended application rates into amount of WEEDAR® MCPA Concentrate, use the following table.


MCPA acid	1 1/2 lb	1 1/4 lb	1 lb	3/4 lb	1/2 lb	3/8 lb	1/4 lb
WEEDAR® MCPA Concentrate	3 pt.	2 1/2 pts	2 pts	1 1/2 pts	1 pt	3/4 pt	1/2 pt

INJURY TO CROPS MAY OCCUR FROM THIS HERBICIDE. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT.

Crop varieties vary in response to MCPA, and some may be easily injured. Apply WEEDAR MCPA Concentrate only to varieties know to be tolerant to MCPA. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to MCPA, contact your seed company or state Agricultural Extension Service for advice.

RECOMMENDATIONS FOR APPLYING WEEDAR MCPA CONCENTRATE BROADLEAF POSTEMERGENCE HERBICIDE

CROP	AMOUNT WEEDAR® MCPA CONCENTRATE PER ACRE	DIRECTIONS
<b>CEREAL GRAINS</b>		
Wheat, Barley, Oats and Rye (not underseeded with legumes)		
Annual Weeds	0.5 pint	Apply after grain is in the 4 leaf stage but not forming joints in the stem. <u>Do not spray in the boot to dough stage.</u>
Annual and Biennial weeds	1-2 pints*	Apply after grain is fully tillered (usually about 4-8" high), but not forming joints in the stem. <u>Do not apply in the boot to dough stage.</u>
Perennial broadleaf weed	1.5-2 pints*	
Emergency control in wheat		Apply when weeds are approaching bud stage, <u>but do not spray grain in the boot to dough stage.</u> The 3 pint (1.5 pound acid equivalent) per acre application <u>injury</u> to wheat. Balance the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the effect of crop injury.
perennial broadleaf weeds	3 pints	



CROP	AMOUNT WEEDAR® MCPA CONCENTRATE PER ACRE	DIRECTIONS
WHEAT, BARLEY, OATS and RYE Underseeded with legumes (alfalfa, alsike, red and ladino clovers, trefoil)	0.5 pint	Treat after grain is tillered and in the 4 leaf stage and when legumes are 2 to 3 inches tall. Small grains should form a protective canopy over the legume seedlings. Do not use more than 6 gallons of water per acre, higher volumes may result in injury to legumes. Do not apply to grain underseeded with vetch or sweet clover. Do not spray when grain is in the boot to dough stage. ✓

\*Use the lower rate if small annual and biennial weeds are the major problem. Use the higher rate if perennial weeds or annual and biennial weeds are present which are in the hard-to-kill categories as determined by local experience. The higher rates increase the risk of grain injury and should be used only where the weed control problem justifies the risk of grain damage.

RESTRICTIONS AND LIMITATIONS FOR USE ON SMALL GRAINS:

- Use a minimum of 10 gallons of water per acre for ground application and 2 gallons of water per acre for aerial application, unless specific site directions listed above indicate otherwise.
- Do not allow livestock to forage or graze treated areas within seven days of slaughter.

CROP	AMOUNT WEEDAR® MCPA CONCENTRATE PER ACRE	DIRECTIONS
FLAX	1/3-1/2 pint	Apply when flax is 2 to 8 inches tall. Do not spray once flax has reached the bud stage. Use at least 8 to 10 gallons of water per acre for ground application and at least 3 to 5 gallons of water per acre for aerial application.

7 of 3

CROP	AMOUNT WEEDAR® MCPA CONCENTRATE PER ACRE	DIRECTIONS
RICE	1 1/2 - 2 1/2 pints	Treat 7 to 10 weeks after planting when rice is fully tillered and 6 to 8 inches above water. Do not spray rice has reached the boot stage. Use the higher rate where sedges, or bulrushes are the major problem. Use lower rate for control of arrowhead, water plantain, and redstem. Do not spray when temperatures are over 90°F. Use at least 3 to 5 gallons of water per acre for aerial application. ✓
Rice (California only)	1 - 1 1/2 pints	Treat when rice is in the 3 to 4 leaf stage, and has well established root systems, usually 18 to 24 days after planting. Applications made during hot weather or at the higher rate may cause temporary stunting of the rice. For best results weeds should be exposed at time of application. If retreatment is necessary, usually 35 to 65 days after planting, the total amount per acre of WEEDAR MCPA Concentrate applied should not exceed 2 1/2 pints in one season. Use 8 to 10 gallons of water per acre for aerial application.

39511-65-

8 of 13

**RESTRICTION AND LIMITATIONS FOR USE ON FLAX AND RICE:**

Do not allow livestock animals to forage or graze treated areas within seven days of slaughter.

Do not grow crayfish or catfish in treated rice fields.

PEAS

1/2 to 3/4  
pint

Treat when peas are 4 to 6 inches tall. Do not spray while peas are in blossom. Peas may show slight injury, but they usually recover a few days after treatment. Use the higher rate before buds form on Canada thistle to prevent from going to seed. Do not spray peas that are stressed from lack of moisture or when temperatures are over 90°F. Apply in 15 to 30 gallons of water per acre for ground application and 10 gallons of water per acre for aerial application.

**RESTRICTIONS AND LIMITATIONS FOR USE ON PEAS:**

Do not allow livestock to forage or graze treated pea fields.

Do not feed treated vines to livestock.

**STORAGE AND DISPOSAL**

Do not store near food, feedstuffs, fertilizers, seeds, insecticides or fungicides. Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL**

Open dumping is prohibited. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL**

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.



9-8-83

**LIMITED WARRANTY AND DISCLAIMER**

The manufacturer warrants that this product conforms to the chemical description on the label, that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions, and that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ✓

# Weedar®

## MCPA - CONCENTRATE

**Broadleaf Postemergence Herbicide**  
Controls weeds in flax, rice, peas, and small grains

**ACTIVE INGREDIENT:**  
Dimethylamine salt of 2-methyl-4-chlorophenoxyacetic acid\* ..... 52.1%  
**INERT INGREDIENTS** ..... 47.9%  
\*2-Methyl-4-chlorophenoxyacetic acid equivalent  
42.6% by weight or four pounds per gallon.

E.P.A. Reg. No. 264-47 AA      E.P.A. Est. No. 264-MO-01

**KEEP OUT OF REACH OF CHILDREN**  
**DANGER PELIGRO**

**PRECAUCION AL USUARIO:** Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(**TRANSLATION: TO THE USER:** If you cannot read English do not use this product until the label has been fully explained to you.)

**PRECAUTIONARY STATEMENTS: See inside booklet.**

For **PRODUCT** Handling, **USE** and **General** Information Call 1-800-334-9745.

For **EMERGENCY** Information **ONLY** Call 24 Hours A Day 1-800-334-7577.

**RHÔNE-POULENC AG COMPANY**  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, N.C. 27709

WEEDAR is a registered trademark of RHÔNE-POULENC.

**CONTENTS 2.5 GAL.**      RP-2200-146-97  
**DO NOT DETACH BOOKLET FROM CONTAINER**

**BEST AVAILABLE COPY**

11513

## PRECAUTIONARY STATEMENTS

### DANGER

#### HAZARD TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage. Harmful if swallowed. May be fatal if absorbed through skin. Avoid contact with eyes, skin or clothing. Avoid breathing vapors or spray mist. Remove and wash contaminated clothing before reuse. Wash thoroughly before eating or smoking.

#### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED, drink large quantities of water. Do not induce vomiting. Get medical attention. IN CASE OF SKIN CONTACT, wash skin with plenty of soap and water. IN CASE OF EYE CONTACT, flush with large amounts of water for 15 minutes. Get medical attention. Never give anything by mouth to an unconscious person.

**NOTE TO PHYSICIAN:** Aspiration hazard may contraindicate the use of gastric lavage.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water except as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes.

Avoid spray drift to susceptible plants such as cotton, beans, tomatoes, and ornamentals. (Coarse sprays are less likely to drift).

Do not apply when weather conditions favor drift from treated areas. Avoid use of smaller diameter spray nozzles. At high air or ground surface temperatures, vapors from this product may injure susceptible plants.

Do not contaminate water used for irrigation or domestic purposes.

Do not apply WEEDAR® MCPA Concentrate through any type of irrigation system.

Do not use the same spray equipment for other purposes unless thoroughly cleaned.

Do not use in or near a greenhouse.

#### GENERAL INFORMATION

Tests in this country and in Europe show that several crops such as flax, clover, and small grains underseeded with legumes are more tolerant of MCPA than they are of 2,4-D.

WEEDAR® MCPA CONCENTRATE Broadleaf Postemergence Herbicide is effective on a large number of broadleaf weeds (mustards, thistles, pigweed, ragweed, burdock, plantain, dandelion, cocklebur, shepherdspurse, stinkweed, purslane, and many others) and is useful for controlling these weeds in certain crops. Use MCPA on oats because oats are more tolerant of MCPA than they are of 2,4-D.

WEEDAR® MCPA Concentrate Broadleaf Postemergence Herbicide is the amine form of MCPA that is easily dissolved in water, even in the hard waters found in some parts of this country. Mix thoroughly while diluting this material and mix again if allowed to stand long before spraying.

## STORAGE AND DISPOSAL

### STORAGE

Do not store near food, feedstuff, fertilizers, seeds, insecticides, or fungicides. Do not contaminate water, food or feed by storage or disposal.

### PESTICIDE DISPOSAL

Open dumping is prohibited. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If container is burned, stay out of smoke.

The manufacturer warrants that this product conforms to the chemical description on the label; that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

To convert local recommendations into terms of WEEDAR® MCPA Concentrate, use the following table.

MCPA acid	1 ¼ lb.	1 lb.	¾ lb.	½ lb.	⅓ lb.	¼ lb.
WEEDAR® MCPA CONCENTRATE	2 ½ pt.	2 pt.	1 ½ pt.	1 pt.	¾ pt.	½ pt.

### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

### GENERAL WORKER PROTECTION STATEMENTS

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Advise workers to stay out of fields during application and until sprays have dried. Regular long-sleeved work clothing should be worn when working in treated fields. See PRECAUTIONARY STATEMENTS, STATEMENT OF PRACTICAL TREATMENT and NOTE TO PHYSICIAN for information on accidental exposures. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: Appropriate signal word (DANGER), area treated with WEEDAR® MCPA-Concentrate Broadleaf Postemergence Herbicide, date of application, appropriate protective clothing, and reentry interval (i.e. until sprays have dried).

CROP	Amount WEEDAR® MCPA CONCENTRATE Per Acre	DIRECTIONS
SMALL GRAINS (wheat, barley, rye, not seeded to legumes) Spring planted	½ pint	Treat in spring when grain is fully tillered and 8 to 10 inches tall. Do not spray grain in the boot to dough stage. Use about 8 gallons of water per acre.
Fall planted	½ pint	Apply in early spring when annual broadleaf weeds are small and grain is fully tillered, but before grain is in the jointed stage. Use about 8 gallons of water per acre.
Emergency treatment for perennial broadleaf weeds in wheat (all producing areas)	3 pints	Treat in spring after grain is tillered and at least 8 to 10 inches tall. Apply when weeds are approaching bud stage, but do not spray grain in the boot to dough stage. The 3 pint (1.5 pound acid equivalent) per acre application of any MCPA product can produce injury to wheat. Balance the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the effect of crop injury. Use at least 8 to 10 gallons of water per acre for ground application and at least 3 to 5 gallons of water per acre for aerial application.

**SMALL GRAINS**  
Underseeded with mixed  
legumes (alfalfa, clover,  
red and ladino clovers,  
trefoil)  
(All producing areas)

½  
pint

Treat in early spring when grain  
is in the tiller to boot stage and 8  
to 10 inches tall, and when  
legume seedlings are 2 to 3  
inches tall. Do not spray when  
grain is in the boot to dough  
stage. Use 5-6 gallons of water  
per acre since higher volumes  
may result in injury to legumes.

<b>an.</b> CROP	<b>Amount WEEDAR® MCPA CONCENTRATE Per Acre</b>	<b>DIRECTIONS</b>
OATS	1½ to 1 pint	Spray after oats are fully tillered (4 to 6 inches tall) but before they are in the boot to dough stage. Apply the lower rate where annual weeds are the major problem, the higher rate for perennial weeds. Use at least 8 to 10 gallons of water per acre for ground application and at least 3 to 5 gallons of water per acre for aerial application.

13 of 13

CROP	Amount WEEDAR® MCPA CONCENTRATE Per Acre	DIRECTIONS
FLAX	1/2-3/4 pint 1/2 pint	Where susceptible weeds such as mustard are the main problem. Where weeds such as pigweed, lambsquarters, stinkweed, ragweed, cocklebur are the problem. Apply when flax is 2 to 8 inches tall but before bud stage. Do not spray after flax is in the bud stage. Use at least 8 to 10 gallons of water per acre for ground application and at least 3 to 5 gallons of water per acre for aerial application.
RICE	1 1/2-2 1/2 pint	Treat 7 to 10 weeks after planting when rice is fully tillered and 6 to 8 inches above water. Do not spray after rice is in the boot stage. Use the higher rate where curly indigo, sedge or bulrush are the major problems. Use lower rate for arrowhead, water plantain, and redstem. Do not spray when temperatures are over 90°F. Use at least 3 to 5 gallons of water per acre for aerial application.
PEAS	1/2-3/4 pint	Treat when peas are 4 to 6 inches tall using 15 to 30 gallons of water per acre for ground application and 10 gallons of water per acre for aerial application. Do not spray while peas are in blossom. Peas may show slight injury, but they usually recover a few days after treatment. Use the higher rate before buds form on Canada thistle to prevent from going to seed. Do not spray peas that are stressed from lack of moisture or when temperatures are below 90°F.

Do not forage or graze meat animals on treated areas within seven days of slaughter.

Do not grow crayfish or catfish in treated rice fields.

Do not feed treated vines to livestock.

Do not forage or graze meat animals in treated pea fields.

# Weedar®

## MCPA - CONCENTRATE

### Broadleaf Postemergence Herbicide

Controls weeds in flax, rice, peas, and small grains

#### ACTIVE INGREDIENT:

Dimethylamine salt of 2-methyl-4-chlorophenoxyacetic acid\* ..... 52.1%

INERT INGREDIENTS ..... 47.9%

\*2-Methyl-4-chlorophenoxyacetic acid equivalent  
42.6% by weight or four pounds per gallon.

E.P.A. Reg. No. 264-47 AA      E.P.A. Est. No. 264-MO-01

**KEEP OUT OF REACH OF CHILDREN  
DANGER PELIGRO**

**PRECAUCION AL USARIO:** Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**TRANSLATION: TO THE USER:** If you cannot read English do not use this product until the label has been fully explained to you.)

**PRECAUTIONARY STATEMENTS: See inside booklet.**

For PRODUCT Handling, USE and General Information Call 1-800-334-9745

For EMERGENCY Information ONLY Call 24 Hours A Day 1-800-334-7577.

#### **RHÔNE-POULENC AG COMPANY**

P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, N.C. 27709

WEEDAR is a registered trademark of RHÔNE-POULENC.

**CONTENTS 2.5 GAL.**      RP-2200-146-97  
**DO NOT DETACH BOOKLET FROM CONTAINER**