

DIRECTIONS

FOLIAGE AND STEM SPRAY

This is the standard method for water-borne sprays. Apply spray directly to both foliage and stems of all plants. Use this method as a "first spray" on thick brush composed primarily of root suckering species such as locust, sumac, elderberry, and brambles. Spray from the time foliage reaches full size until plants begin to go dormant. For effective control *all leaves must be drenched, and stems and suckers must be thoroughly wet to ground line.*

Add 4 quarts of WEEDAR AMINE BK to 100 gallons of water. Mix thoroughly before spraying. Apply with power equipment.

FRILL METHOD

Make a frill by using an axe to cut overlapping notches in a continuous ring around the trunk near its base. Cut through the bark, but do not remove chips. This method is recommended for cull trees 5 to 6 inches in diameter and over. Treat freshly cut frills at any time of the year.

Mix 2 gallons WEEDAR AMINE BK in 50 gallons of water. Pour in enough material to fill notches without wasting chemical.

TREE INJECTION

Make injections as near the root collar as possible, using one injection per inch of trunk diameter at breast height (4½ feet). For resistant species such as hickory, injections should overlap. For best results, injections should be made during the growing season — May 15 to October 15.

Mix 1 gallon of WEEDAR AMINE BK in 19 gallons of water.

Form No 1571D 9 71 SED 1M

Printed in U. S. A.

AMCHEM PRODUCTS, Inc.

FIRST NAME IN HERBICIDE RESEARCH

AMBLER, PA ■ Clinton, Iowa ■ St. Joseph, Mo ■ Fremont, Calif

DIRECTIONS

FOLIAGE AND STEM SPRAY

This is the standard method for water-borne sprays. Apply spray directly to both foliage and stems of all plants. Use this method as a "first spray" on thick brush composed primarily of root suckering species such as locust, sumac, elderberry, and brambles. Spray from the time foliage reaches full size until plants begin to go dormant. For effective control *all leaves must be drenched, and stems and suckers must be thoroughly wet to ground line.*

Add 4 quarts of WEEDAR AMINE BK to 100 gallons of water. Mix thoroughly before spraying. Apply with power equipment.

FRILL METHOD

Make a frill by using an axe to cut overlapping notches in a continuous ring around the trunk near its base. Cut through the bark but do not remove chips. This method is recommended for cut trees 5 to 6 inches in diameter and over. Treat freshly cut frills at any time of the year.

Mix 2 gallons WEEDAR AMINE BK in 50 gallons of water. Pour in enough material to fill notches without wasting chemical.

TREE INJECTION

Make injections as near the root collar as possible, using one injection per inch of trunk diameter at breast height (4 1/2 feet). For resistant species such as hickory, injections should be made. For best results, injections should be made during the growing season - May 15 to October 15.

Mix 1 quart of WEEDAR AMINE BK in 19 gallons of water.

AMCHEM PRODUCTS, Inc.

FIRST NAME IN HERBICIDE RESEARCH

AMBLER, PA. ■ Clinton, Iowa ■ St. Joseph, Mo. ■ Fremont, Calif.



AR

JECTS, Inc.

RESEARCH

Mo. ■ Fremont, Calif

AA