

UNION
CARBIDE

Weedar[®]

Sodium MCPA

Controls weeds in rice, small grains, and peas.

ACTIVE INGREDIENT
Sodium salt of 2-methyl-4-chlorophenoxyacetic acid* 24.0%

INERT INGREDIENTS 76.0%
2-Methyl-4-chlorophenoxyacetic acid equivalent 21.7% or 2 pounds per gallon

E.P.A. Reg. No. 264-55AA

E.P.A. Est. No. 264-MO-01

with comment
OCT 20 1983
264-55

WARNING: Keep out of reach of children.
PRECAUTIONARY STATEMENTS: See inside booklet.

**IN CASE OF EMERGENCY, TELEPHONE COLLECT
(24 HOURS A DAY) IN U.S.A. (304) 744-3487**

UNION CARBIDE AGRICULTURAL PRODUCTS COMPANY INC
P.O. Box 12014 T.W. Alexander Drive Research Triangle Park, NC 27709

WEEDAR and UNION CARBIDE are registered trademarks

MADE IN U.S.A.

APC-7301-42

DO NOT DETACH BOOKLET FROM CONTAINER

BELOW CONTAINER IS A...

WARNINGS

Harmful if swallowed. Avoid contact with eyes, nose, mouth, and skin. Wash thoroughly after handling.
Irritation may occur if the product is in contact with the eyes, nose, mouth, or skin. If contact occurs, flush with plenty of water. In case of eye contact, rinse thoroughly with water. In case of skin contact, wash with plenty of soap and water.

Environmental Hazards:
Avoid spray drift to desirable plants such as cut flowers, ornamentals, and shrubs. Do not use on lawns or other ornamentals. Do not use on lawns or other ornamentals. Do not use on lawns or other ornamentals. Do not use on lawns or other ornamentals.

Storage and Disposal Statements:
Do not store near fertilizers, seeds, insecticides, or fungicides. Do not contaminate water, food, or feed by storage or disposal.
Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
Container Disposal:
Triple Rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants that this product conforms to the chemical description on the label, that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions, and that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.
This warranty does not extend to, and the Buyer shall be solely responsible for, any and all loss or damage which results from the use of this product in any manner which is inconsistent with the label directions, warnings or cautions. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

GENERAL INFORMATION

Among the weeds controlled are:

Susceptible

Lamb's quarters
marestail
radweed
stinkweed
wild radish
yellow rocket
winter cress
purslane
peppercress
cockle

Moderately Resistant

buttercup
Canada thistle
dandelion
plantain
hemlock
stinging nettle
sowthistle
vetch
kochia
shepherdspurse
whiteweed (hoary cress)

DIRECTIONS

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CROP	AMOUNT PER ACRE	REMARKS
SMALL GRAINS (Wheat, barley, rye, oats)		Treat in the spring when grain is fully tillered and 8 to 10 inches tall Do not spray when grain is in the boot to dough stage
Not underseeded with mixed legumes Broadleaf weeds	1 to 3 pints	

Emergency treatment
of perennial cereal
weeds in wheat

1/2 pint

Apply 1/2 pint per acre when weeds are approaching boot stage but not past grain. Do not apply in the boot stage. Do not apply 1/2 pint and emergency per acre application of 1/2 MCFE product can produce injury to wheat. Balance the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered spot treatment is suggested to minimize the extent of crop injury. Treat in the spring when grain is fully tillered and 8 to 10 inches tall.

Underseeded with
mixed legumes
alfalfa, alsike,
red and ladino
clovers, etc.

1/2 pint

Treat in early spring when legumes are 2 to 3 inches tall when grain is fully tillered and 8 to 10 inches tall. Do not spray when grain is in the boot to dough stage. Use 5 to 6 gallons of water per acre since higher volumes may cause injury to legumes.

RICE

3 to 5 pints

Controls arrowhead, water plantain and redstem. Controls sedge and bulrush. Treat 55 to 65 days after planting when rice is fully tillered and 6 to 8 inches above water. Water should be at least 2-3 inches deep. Do not spray after rice is in the boot stage or when temperatures are over 90°F.

For airplane applications, use 10 gallons of water per acre. For ground application use a conventional low pressure, low volume sprayer that delivers 10 to 40 gallons of water per acre.

When diluting the material must be mixed thoroughly before using. If the material is allowed to stand for any length of time after dilution, it should be stirred again before using.

PEAS

1/2 to 1 1/2 pint

Treat when peas are 4 to 6 inches tall using 15 to 30 gallons of water per acre. Do not spray while peas are in blossom. Peas may show slight injury, but they usually recover a few days after treatment. Use the higher rate before buds form on Canada thistle to prevent from going to seed.

Do not feed treated vines to livestock. Do not forage or graze treated pea fields. Do not spray peas that are stressed from lack of moisture or when temperatures are over 90°F.

BEST DOCUMENT AVAILABLE

OCT 10 1983

Dr. Warren Z. Davis
Union Carbide Agricultural Products
Company, Inc.
P.O. Box 12014
2110 Alexander Drive
Research Triangle Park, NC 27709

Dear Dr. Davis:

Subject: Acute Toxicity Data
Eccentrodin 1007
LHA Registration No. 204-55
Your Application dated July 6, 1983

The amendment referred to above, submitted in connection with registration under FIFRA, is acceptable, provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing amended labeling:
 - a. On the front panel and in the precautionary statement, change GMBHC to L1031 based on the results of the rabbit eye irritation study. (Note: The eye irritation study using two cycloplegic regimens is considered supplementary data since at least six animals were not used.)
 - b. Where reference to fish and aquatic animals, change "may cause eye irritation" to "corrosive, causes irreversible eye damage."
 - c. Lastly, Environmental Hazards Statement to read: "Do not apply directly to water except as specified on this label."
2. Submit also (3) copies of your final printed labeling when you release the product for shipment.

CORRECTIONS							
SYMBOL							
SURNAME							
DATE							

-2-

A stamped copy of the label is enclosed for your records. We are accepting this amendment based on your submission of data pursuant to 40 CFR 162.8(b). The precautionary text is prescribed in the regulations at 162.10(h).

Sincerely yours,

Richard P. Mountfort *RFM*
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure

TS-767: MOUNTFORT: DCR-25929: HANG-1004C: vgr: Raven: 479-2013: 6/10/83