

## DIRECTIONS

### FOLIAGE STEM SPRAY

This is the standard method for water-borne sprays. Apply spray directly to both foliage and stems of all plants. Use this method as a "first spray" on thick brush composed primarily of root suckering species such as locust, sumac, elderberry and brambles. Spray from the time foliage reaches full size until plants begin to go dormant. For effective control, **all leaves, stems and suckers must be thoroughly wet to ground line.**

Add 4 quarts of WEEDAR 2,4,5-T to 100 gallons of water. Mix thoroughly before spraying. Apply with powder equipment.

### FRILL METHOD

Make a frill by using an axe to cut overlapping notches in a continuous ring around the trunk near its base. Cut through the bark, but do not remove chips. This method is recommended for cull trees 5 to 6 inches

in diameter and over. Treat freshly cut frills at any time of year.

Mix 2 gallons WEEDAR 2,4,5-T in 50 gallons of water. Pour in enough material to fill notches without wasting chemical.

### TREE INJECTION

Make injections as near the root collar as possible, using one injection per inch of trunk diameter at breast height (4½ ft.). For resistant species such as hickory, injections should overlap. For best results, injections should be made during the growing season.

### CONCENTRATE INJECTORS

Use 1—2 ml of concentrate WEEDAR 2,4,5-T per injection. The injector bit must penetrate inner bark.

### DILUTE INJECTORS

Mix 1 gallon of WEEDAR 2,4,5-T in 19 gallons of water.

# WEEDAR<sup>(R)</sup>

# 2,4,5-T

*Jan 13, 1975*

*264-62*

For the Control of Woody Plants on utility rights-of-way, along highways, fencelines or pipelines.

**CAUTION:** Keep out of reach of children. See other cautions on right panel.

#### ACTIVE INGREDIENTS:

\*Triethylamine salt of 2,4,5-Trichlorophenoxyacetic acid . . . . . 56.9%

INERT INGREDIENTS: . . . . . 43.1%

\*2,4,5-Trichlorophenoxyacetic acid equivalent 40.8% by weight  
or 4 pounds per gallon



A RORER-AMCHEM COMPANY



## AMCHEM PRODUCTS, INC.

First Name in Herbicide Research

AMBLER, PA. ■ Clinton, Iowa ■ St. Joseph, Mo. ■ Fremont, Calif.

EPA Reg. No. 264-62-ZA      EPA Est. 264-PA-1

# WEEDAR<sup>®</sup> 2,4,5-T

*Jan 13, 1975  
264-62*

For the Control of Woody Plants on utility rights-of-way, along highways, fencelines or pipelines.

**CAUTION:** Keep out of reach of children. See other cautions on right panel.

**ACTIVE INGREDIENTS:**

\*Triethylamine salt of 2,4,5-Trichlorophenoxyacetic acid . . . . . 56.9%

**INERT INGREDIENTS:** . . . . . 43.1%

\*2,4,5-Trichlorophenoxyacetic acid equivalent 40.8% by weight or 4 pounds per gallon



**AMCHEM PRODUCTS, INC.**

First Name in Herbicide Research  
 AMBLER, PA. ■ Clinton, Iowa ■ St. Joseph, Mo. ■ Fremont, Calif.  
 EPA Reg. No. 264-62-ZA      EPA Est. 264-PA-1

**ADVANTAGES OF  
WEEDAR 2,4,5-T**

WEEDAR 2,4,5-T contains 2,4,5-T as an amine salt. Does not have the volatility problem associated with some ester formulations.

WEEDAR 2,4 5-T is effective on root suckering species resistant to 2,4-D.

WEEDAR 2,4,5-T may be used in areas of high temperature, or where volatility is a problem.

WEEDAR 2,4,5-T is a formulation proved effective for controlling the following species of woody plants:

- |                    |                     |                         |
|--------------------|---------------------|-------------------------|
| <u>ALDER</u>       | <u>ELM</u>          | <u>SASSAFRAS</u>        |
| <u>ASPEN</u>       | <u>HICKORY</u>      | <u>SPICEBUSH</u>        |
| <u>BIRCH</u>       | <u>HONEYSUCKLE</u>  | <u>SUMAC</u>            |
| <u>BLACKBERRY</u>  | <u>LOCUST</u>       | <u>SWEET GUM</u>        |
| <u>BOXELDER</u>    | <u>OAK</u>          | <u>WILD CHERRY</u>      |
| <u>BRAMBLES</u>    | <u>OSAGE ORANGE</u> | <u>WILD GRAPE</u>       |
| <u>CHESTNUT</u>    | <u>POISON IVY</u>   | <u>VIRGINIA CREEPER</u> |
| <u>CHOKECHERRY</u> | <u>POISON OAK</u>   | <u>WILLOW</u>           |
| <u>ELDERBERRY</u>  | <u>RASPBERRY</u>    | <u>WITCH HAZEL</u>      |

(and many other species)

**CAUTION**

Harmful if swallowed. Avoid contact with skin, eyes or clothing.  
 Avoid spray drift to susceptible plants such as cotton, beans, tomatoes, tobacco, and ornamentals as plant injury may result. Coarse sprays are less likely to drift. Do not use same spray equipment for other purposes.  
 Under very high temperatures of use, vapors from this product may injure susceptible plants in the immediate vicinity. Do not store near fertilizers, seeds, insecticides, or fungicides. Do not use in a greenhouse. Do not use around the home, in parks or recreational areas.  
 Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of waste. Do not contaminate water used for irrigation or domestic purposes.  
 Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.  
 Rinse spray equipment and containers and dispose of liquid wastes in a pit in non-crop lands away from water supplies and not accessible to livestock. Do not reuse empty containers. Destroy in a safe place.  
**Do not use this product for purposes other than those recommended on this label.**

**WARRANTY**

Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. Amchem, therefore, makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees, in purchasing this product, to assume the risks and in the event of damages arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of Amchem's liability.