

WEEDONE®

GRANULAR



ACCEPTED

AUG 15 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, the following pesticide is registered under EPA Reg. No. 264-83

CAUTION: Keep out of reach of children.
See Back Panel for Additional Cautions

Now Improved with SLAVEX

WEEDONE GRANULAR LAWN WEED CONTROL will kill weeds such as dandelion, plantain, chickweed, spotted spurge, clover and many others without injuring turf grasses.

Follow directions on label for use. Follow label for manufacturer's instructions.

WEEDONE GRANULAR LAWN WEED CONTROL

WEEDONE GRANULAR LAWN WEED CONTROL



ACCEPTED
 AUG 16 1975
 U.S. Environmental Protection Agency
 Office of Pesticide Programs
 Washington, D.C. 20460
 EPA Reg. No. 264-83

CAUTION: Read label carefully. See Back Panel for Additional Cautions

Now Improved with SILVEX

WEEDONE GRANULAR LAWN WEED CONTROL will kill weeds such as dandelion, plantain, chickweed, spotted spurge, clover and many others without injuring turf grasses. This material should be applied with a lawn spreader. Follow spreader manufacturer's directions for application of dry weed killers.

This package contains enough for 1 application to 3,000 square feet of lawn. Use WEEDONE GRANULAR LAWN WEED CONTROL as purchased, do not mix with water or otherwise dilute it.

Treats 3,000 sq. ft. of lawn area

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic acid*	1.6%
2,4-Dichlorophenoxyacetic acid, dimethylamine salt*	1.93%
SILVEX [2,4,5-Trichlorophenoxypropionic acid]**	0.80%
SILVEX [2,4,5-Trichlorophenoxypropionic acid, dimethylamine salt]**	0.94%
INERT INGREDIENTS:	94.72%


*Total 2,4-Dichlorophenoxyacetic acid equivalent 3.22%
 **Total SILVEX [2,4,5-Trichlorophenoxypropionic acid equivalent] 1.61%

AMCHEM PRODUCTS, INC.
 First Name: Herbicide Research
 AMBLER, PA ■ Clinton, IA ■ St. Joseph, MO ■ Fremont, CA

E.P.A. Reg. No. 264-P3AA E.P.A. Est. 264-PA-1

WEEDONE®

GRANULAR



BROADLEAF PLANTAIN



SPOTTED SPURGE



CLOVER



NARROWLEAF PLANTAIN

To control dandelion, broadleaved and narrowleaved plantain, common and smooth-stem chickweed, spotted spurge, clover, and many other broad-leaf weeds without injury to turf grasses, spread WEEDONE GRANULAR LAWN WEED CONTROL at the rate of 2 1/2 pounds per 1500 square feet of lawn. The contents of this 4 1/2 pound bag is enough for a uniform over-all treatment of 3,000 square feet of lawn.

Mow your lawn a few days before treatment so grass is not over 1 1/2" tall when treated. Wait at least two days after a treatment before mowing.

New lawns should have become established and been cut at least twice before treatment. If in doubt, make a trial application to a limited area.

Allow at least 3 weeks after treating with WEEDONE GRANULAR LAWN WEED CONTROL before reseeding. Water the area thoroughly if rain does not fall during this period.

Bentgrass and some southern or warm season turf grasses such as some strains of Bermuda and Zoysia, St. Augustine, Bahia, Centipede


WEEDONE GRANULAR LAWN WEED CONTROL



BROADLEAF PLANTAIN



SPOTTED SPURGE



CLOVER



NARROWLEAF PLANTAIN

To control dandelion, broadleaved and narrowleaved plantain, common and nutcracker chickweed, spotted spurge, clover, and many other broadleaf weeds without injury to turf grasses, spread WEEDONE GRANULAR LAWN WEED CONTROL at the rate of 2 1/2 pounds per 1500 square feet of lawn. The contents of this 4 1/2 pound bag is enough for a uniform over-all treatment of 3,000 square feet of lawn.

WHEN TO USE

WEEDONE GRANULAR LAWN WEED CONTROL can be applied whenever the weeds are noticed, but it is most effective in the early fall or in spring when warm weather and adequate soil moisture encourage rapid weed growth. Under ordinary conditions, weeds will start to resist in a few days. Effectiveness is increased if application is made to damp or wet foliage. Treatments applied during dry or very cool weather act more slowly. Dandelions can be treated at any time, but respond best to late summer or early fall treatments.

Repeat the treatment in two weeks if heavy rain falls within 24 hours. A few weeds (including veronica, violets, sheep sorrel and nutcracker weed) resist chemical treatment and probably will not die.

Another reason for favoring early fall or spring treatments is that turf grasses are growing strongly at these times and will fill in many spots where weeds have died.

Proper fertilization is essential for improving and maintaining a good turf. Within 1 or 2 weeks after treatment with WEEDONE GRANULAR LAWN WEED CONTROL, apply fertilizer as recommended by your local dealer.

HOW TO USE

Proper application is essential for best results. So is the correct amount of WEEDONE GRANULAR LAWN WEED CONTROL. It is applied to the area specified and that it is applied uniformly. Check out spreader's output by treating a small measured area. Crisscross from end to end and then from side to side until the prescribed amount of material is distributed evenly. Adjust the spreader setting as necessary.

When treating an entire lawn, mark off the area to be treated and cover it by crisscrossing in the same manner. Overlap wheel tracks by 2 to 3 inches to avoid missed strips. If grass is wet, check to see opening occasionally to be sure material is flowing freely.

Be sure to shut off spreader when stopping or turning around.

SUGGESTED SPREADER SETTING FOR A CRISSCROSS APPLICATION.

	Model	Setting
CENTRAL	20"	# 6
HERITAGE HOUSE	20"	# 4 1/2
LAWN BEAUTY	24"	# 6
SCOTT'S	24"	# 3 3/4
SEARS	20"	# 6

Adjust the spreader setting if necessary to cover specified area.

Mow your lawn a few days before treatment so grass is not over 1 1/2" tall when treated. Wait at least two days after a treatment before mowing.

New lawns should have become established and been cut at least twice before treatment. If in doubt, make a trial application to a limited area.

Allow at least 3 weeks after treating with WEEDONE GRANULAR LAWN WEED CONTROL before reseeding. Water the area thoroughly if rain does not fall during this period.

Bermgrass and some southern or warm season turf grasses such as newer strains of Bermuda and Zoysia, St. Augustine, Bahia, Carpetgrass and Centipede may be injured by this product. Use only if weed problem is severe. Treat with caution in the spring or fall only, not during hot summer months. Apply half the recommended rate by using the same spreader setting, but going over the area only once. Repeat the treatment in 2 weeks.

CAUTION

AVOID CONTACT WITH SKIN, EYES OR CLOTHING

Avoid drift by applying only when the air is still. Accidental contact could injure or kill desirable ornamental plantings or vegetables. WEEDONE GRANULAR LAWN WEED CONTROL should not be used in vegetable or flower gardens, or on broadleaf ground covers or dandelion.

Do not store near fertilizers, seed, insecticides or fungicides, or fish in a greenhouse.

Do not contaminate water used for irrigation or domestic purposes.

Do not reuse container. Destroy in a safe place. Do not burn.

WARRANTY

Amchem warrants that composition of this product conforms to the chemical description given in the ingredient statement and that the product is suited for the purpose described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks even though label directions are followed. Amchem therefore makes no other express or implied warranty, and no agent of Amchem is authorized to do so. Buyer agrees in purchasing this product to assume the risks and in the event of damages arising from a breach of the warranty to accept refund of the purchase price of the product as full discharge of Amchem's liability.

Camerman's Note

Poor Copy



AMCHEM PRODUCTS, INC.

First Name in Herbicide Research

AMBLER, PA ■ Clinton, IA ■ St Joseph, MO ■ Fremont, CA

102269

Form No. 2609-D-6-75



TUF-WELD®

E.P.7-75

WEEDONE GRANULAR LAWN WEED CONTROL