

2-1-74

## GENERAL INFORMATION AND DIRECTIONS

AMIZINE acts slowly but thoroughly. Weeds start to turn white or brown within a week and are usually dead in about three weeks. Small weeds growing actively are killed more rapidly than weeds in a dormant or dried condition.

Apply AMIZINE to small weeds in the spring or early summer, or before weeds have matured and gone to seed. It is not necessary to spray bare ground before weed growth starts in the spring.

For best results on tall weeds, cut or mow to ground line and remove. Treat one or two weeks later when regrowth appears.

In arid areas, AMIZINE should be watered into the ground if no rain falls within a week after treatment.

Where johnsongrass, bermudagrass or other weeds are not controlled by the first application, treat again with the same rate when regrowth is 6 to 12 inches tall (about 2 months).

**Mixing sprays:** Add the required amount of AMIZINE to a small portion of water and mix into a paste. Then with constant stirring, add the rest of the water necessary for application. This product forms a suspension in water, not a solution. Provide continuous agitation in the spray tank.

**Spray equipment:** AMIZINE may be applied with a watering can, a portable pressure sprayer, a knapsack sprayer, or a motor driven pump-type sprayer. Screens in nozzles, as well as those in the suction lines, should be no finer than 50 mesh size. A Tee-Jet 8002 or its equivalent fan type of nozzle has been used successfully.

AMIZINE does not corrode spray equipment, is non-flammable, and does not stain metal, stone or concrete. To clean AMIZINE from spray equipment, flush several times with clear water after each use.

## AMOUNTS TO USE

General vegetation control:  
around buildings, driveways, storage areas, fence lines, parking lots, drive-in-theaters.

AMIZINE	WATER	AREA
1 cup	2 gallons	500 sq. ft. (10'x50')
1 pound	5 gallons	2,000 sq. ft. (40'x50')
5 pounds	25 gallons	10,000 sq. ft. (100'x100')
20 pounds	100 gallons	1 acre

Some of the weeds controlled by recommended rates of AMIZINE:

alfalfa	lambquarters	sow thistle
bluegrass	mustard	plantain
Canada thistle	nightshade	poison ivy
crabgrass	pigweed	purslane
dandelion	quackgrass	velvet grass
foxtail	ragweed	wild oats
kochia	smartweed	witchgrass
	chrysanthemum weed	

Weed and grass control in bearing or non-bearing apple orchards:

Mix 7 pounds of AMIZINE in 100 gallons of water for each acre of ground to be treated. Apply AMIZINE in the spring, after weeds emerge, but before apple fruit starts to form.

Wet weeds thoroughly, treating a band at least 3 feet wide at the base of each tree. Do not allow spray to contact stems, foliage or trunks of these trees. Use a coarse spray with pump pressure adjusted to less than 40 psi.

Use about 2 1/2 gallons of spray per 1,000 square feet. If treating a swath down each side of a row of trees, use AMIZINE in proportion to the area actually sprayed. Example: tree rows are 40 ft apart and you spray a 10 ft. swath down each side of the row, you are spraying half the area of your orchard (20 ft. sprayed out of 40 ft. rows).



# AMIZINE

**AMCHEM PRODUCTS, Inc.**  
Originals of 240, 245T and Am-Trip Weed Killers

AMBLER, PA. • Clinton, Iowa • St. Joseph, Mo. • Fremont, Calif.

