

ACTIVE INGREDIENTS:
Alpha Naphthyl Acetamide 1%
INERT INGREDIENTS: 99.6%


CAUTION: May cause skin irritation. Avoid prolonged contact.

NET WEIGHT



1 LB.

264-129 **FOR PROFESSIONAL USE**



CAUTION—KEEP OUT OF REACH OF CHILDREN. MAY CAUSE SKIN IRRITATION. AVOID PROLONGED CONTACT. DO NOT TAKE INTERNALLY.

WARRANTY

The use of this material being beyond our knowledge and control, we make no warranty, express or implied, as to the effects of such use, whether or not in accordance with directions.

U.S. Patents: 2,258,291 and 2,258,292
Printed in U.S.A.

Select sturdy cuttings from healthy plants. Most cuttings root best when cut just below a node or the point of attachment of a leaf.

Moisten the end of the stem slightly, then dip base of cutting in ROOTONE No. 10. Tap off any excess powder.

Insert cuttings in moist sand, vermiculite, sphagnum or peat moss, or other satisfactory rooting medium.

Do not allow cuttings to become dry or wilted from the time they are cut until they are rooted. . . . 3 to 5 weeks or longer, depending on plant species.

Among the plants that will root are rhododendron, American and English holly, juniper.

USDA Reg. No. 264-129