ACTIVE INGREDIENTS:
Identifications Napthaleneacetic acid
INERT INGREDIENTS:

3 OZS. NET WT.

CAUTION: Keep out of reach of children. Harmful if swallowed.

The hormone-like starter for levelier flewers and finer shrubs. Reduces wift and less when transplanting flower seedlings, young shrubs and trees.

USDA, Reg. No. 264-130

For seedling flowers, dissolve ¼ teaspoon of TRANS-PLANTONE in 21/2 gallons of water, or 5 teaspoons of TRANSPLANTONE in 50 gailons of water. Use 1 cup of solution per plant. Pour around the roots of each plant before covering with soil.

For treating young woody shrubs or trees, mix 1 teaspoon of TRANSPLANTONE in 1 gallon of water. Use 1 quart of solution for most shrubs, more for larger shrubs or trees. Pour around the roots or ball of earth before covering with soil.

Use additional water as necessary to bring soil around each transplant to the desired moisture.

Use on: Annual flowers such as petunias, marigolds, zinnias, ageratum. Perennial flowers such as carnations, enrysanthemums, daisies, delphinium. Shrubs such as Taxus (yew), arborvitae, juniper, nemlock, privet, box-W000, 10565.

CAUTION: Do not reuse container. Destroy in a safe place. Form No. 2600---11-68PL Printed in U.S.A.