

DIRECTIONS

BASAL BARK SPRAY

Use this method for control of woody plants on utility rights-of-way. Spray the basal bark of trees and shrubs with a spray nozzle. The spray should be applied to the basal bark of the tree or shrub. The spray should be applied to the basal bark of the tree or shrub. The spray should be applied to the basal bark of the tree or shrub.

Maximum application of DINOX SUPER-6 should be 10 gallons per acre. Apply with a spray nozzle. The spray should be applied to the basal bark of the tree or shrub. The spray should be applied to the basal bark of the tree or shrub.

STUMP TREATMENT

This method can be used for control of woody plants on utility rights-of-way. Spray the end of the stump with a spray nozzle. The spray should be applied to the end of the stump. The spray should be applied to the end of the stump.

Use this method after the stump has been cut. Spray the end of the stump with a spray nozzle. The spray should be applied to the end of the stump. The spray should be applied to the end of the stump.

Maximum application of DINOX SUPER-6 should be 10 gallons per acre. Apply with a spray nozzle. The spray should be applied to the end of the stump.

FRILL TREATMENT

This method can be used for control of woody plants on utility rights-of-way. Frill the bark of the tree or shrub with a spray nozzle. The spray should be applied to the frilled bark of the tree or shrub. The spray should be applied to the frilled bark of the tree or shrub.

Maximum application of DINOX SUPER-6 should be 10 gallons per acre. Apply with a spray nozzle. The spray should be applied to the frilled bark of the tree or shrub.

TREE INJECTION

This method can be used for control of woody plants on utility rights-of-way. Inject the herbicide into the tree or shrub. The herbicide should be injected into the tree or shrub. The herbicide should be injected into the tree or shrub.

Use this method for control of woody plants on utility rights-of-way. Spray the basal bark of trees and shrubs with a spray nozzle. The spray should be applied to the basal bark of the tree or shrub. The spray should be applied to the basal bark of the tree or shrub.

Maximum application of DINOX SUPER-6 should be 10 gallons per acre. Apply with a spray nozzle. The spray should be applied to the basal bark of the tree or shrub.

DORMANT CANE BROADCAST

Use this method for control of woody plants on utility rights-of-way. Broadcast the herbicide on dormant canes. The herbicide should be broadcast on dormant canes. The herbicide should be broadcast on dormant canes.

Maximum application of DINOX SUPER-6 should be 10 gallons per acre. Apply with a spray nozzle. The herbicide should be broadcast on dormant canes.

AERIAL SPRAY

Use this method for control of woody plants on utility rights-of-way. Spray the basal bark of trees and shrubs with a spray nozzle. The spray should be applied to the basal bark of the tree or shrub. The spray should be applied to the basal bark of the tree or shrub.

Maximum application of DINOX SUPER-6 should be 10 gallons per acre. Apply with a spray nozzle. The spray should be applied to the basal bark of the tree or shrub.

Use this method for control of woody plants on utility rights-of-way. Spray the basal bark of trees and shrubs with a spray nozzle. The spray should be applied to the basal bark of the tree or shrub. The spray should be applied to the basal bark of the tree or shrub.

DINOX SUPER-6 WOODY PLANT HERBICIDE

For control of woody plant species on utility rights and other industrial areas.

CAUTION: Keep out of the reach of children. Cautions on right panel.

ACTIVE INGREDIENTS:

**2,4-Dichlorophenoxyacetic acid, butoxy ether

**2,4,5-Trichlorophenoxyacetic acid, butoxy ether

INERT INGREDIENTS:

*2,4-Dichlorophenoxyacetic acid equivalent 29.3 pounds per gallon.

**2,4,5-Trichlorophenoxyacetic acid equivalent or 3 pounds per gallon.



AMCHEM PRO

First Name in Herbicide Research
AMBLER PA ■ Clinton, Iowa ■ St. Joseph, Mo.
EPA Reg. No. 264-132

