

AMITROL

LIQUID HERBICIDE

Harmful if swallowed. Avoid contact with skin, eyes or clothing.

NOTE: Do not allow spray solution to remain in a sprayer any longer than necessary. Flush all spray equipment with water immediately after use to prevent corrosion of metal parts.

Do not use this product for any purpose that might harm you or others. If exposed to temperatures of 100°F or below, the container must be vented.

Amitrole (3-amino-1,2,4 triazole)*

* Equivalent to 2 lbs per gallon.



First Name in Herbicide Research

AMBLER, PA ■ Clinton, IA ■ St. Joseph, MO ■ Fremont, CA

EPA Rev. No. 254-135-AA

[illegible][illegible][illegible]

KNAPSACK SPRAYERS - Carry the tank through a variety of leaves and stems, and can be used for spraying all types of plants for control of water. One gallon of spray solution is enough to cover 1000 sq. ft. of leaf surface. The sprayer is used in mountain states and other dry areas for spraying **AMTRAC** to control water and soil, and for spraying.

POWER SPRAYERS Apply through the spray gun into water, on the surface of the water.

		Gallons AMTROL-T	Gallons WATER	WHEN TO APPLY
Preventive	1000 gph	1	100	Apply before the pump starts to pump and before the pump has been running for 10 minutes. A 1000 gph pump will pump 1000 gallons of water in 10 minutes.
Remedial	1000 gph	1	100	Apply after the pump has been running for 10 minutes and before the pump has been running for 10 minutes.

AMOUNTS PER ACRE			WHEN TO TREAT
GROUND SPRAYING		AERIAL SPRAYING	
1. COTTON	1.00 LBS. 10% DDT	1.00 LBS. 10% DDT	Apply when the first sign of infestation is observed.
2. COTTON	1.00 LBS. 10% DDT	1.00 LBS. 10% DDT	Apply when the first sign of infestation is observed.
3. COTTON	1.00 LBS. 10% DDT	1.00 LBS. 10% DDT	Apply when the first sign of infestation is observed.
4. COTTON	1.00 LBS. 10% DDT	1.00 LBS. 10% DDT	Apply when the first sign of infestation is observed.
5. COTTON	1.00 LBS. 10% DDT	1.00 LBS. 10% DDT	Apply when the first sign of infestation is observed.
6. COTTON	1.00 LBS. 10% DDT	1.00 LBS. 10% DDT	Apply when the first sign of infestation is observed.
7. COTTON	1.00 LBS. 10% DDT	1.00 LBS. 10% DDT	Apply when the first sign of infestation is observed.
8. COTTON	1.00 LBS. 10% DDT	1.00 LBS. 10% DDT	Apply when the first sign of infestation is observed.
9. COTTON	1.00 LBS. 10% DDT	1.00 LBS. 10% DDT	Apply when the first sign of infestation is observed.
10. COTTON	1.00 LBS. 10% DDT	1.00 LBS. 10% DDT	Apply when the first sign of infestation is observed.

WARRANTY

As a result, the authors conclude that the use of the *in vitro* model is not appropriate for the study of the effects of the chemical on the *in vivo* system. The authors also suggest that the use of the *in vitro* model is not appropriate for the study of the effects of the chemical on the *in vivo* system.