

DIRECTIONS FOR USE

Apply BUTYRAC ester broadleaf herbicide in spring or fall, when seedling legumes have 2 to 4 trifoliate leaves and when weeds are less than 3 inches high. Applications made later to more mature weeds will give less satisfactory control. Weeds such as dock and fanweed that form rosettes should be treated when these rosettes are less than 3 inches across.

Apply BUTYRAC ester broadleaf herbicide in 5 or more gallons of water per acre using air or ground equipment. The total volume of spray applied should be enough for good coverage, but not so much that spray will run off weed leaves.

For irrigated crops, apply BUTYRAC ester broadleaf herbicide as soon as possible after irrigation. Delay the next irrigation for at least 7 to 10 days after spraying to avoid washing the chemical into the root zone.

Do not use BUTYRAC ester broadleaf herbicide as a pre-emergence spray. Do not use on a sweet clover or other legumes not mentioned.

WEED CONTROL IN ALFALFA AND BIRDSFOOT TREFOIL

Weeds Controlled:	Amount
Lambsquarters, pigweed, kochia, nightshade, Russian thistle, prickly (Chinese) lettuce, groundsel, stinkweed, fanweed, smartweed, annual ragweed, bull thistle, filaree, shepherd's purse, dock, fiddleneck	BUTYRAC ester broadleaf herbicide per acre
Fiddleneck with more than 2-3 leaves will only be suppressed	2 pints to 4 pints

Use this higher rate only if weeds have grown to height of 4 to 6 inches before spraying. Poorer weed control and crop stunting can be expected when using this higher rate on older weeds.

CAUTION: Do not graze treated crops or feed hay from treated crops to livestock within 60 days after application.

Band Treatments:

Use BUTYRAC ester broadleaf herbicide in proportion to the area actually sprayed. For example, if a 12 inch band is sprayed on 36 inch rows, the amount used per banded acre will be 1/3 of the amount used per broadcast acre. Calibrate sprayers before making band applications. Keep boom at the same height while calibrating and spraying.

MIXING DIRECTIONS

Add one-half the required amount of water to the spray tank, then add this product with agitation, and finally the balance of the water with continued agitation. The material forms an emulsion in water—not a solution. This tends to separate on standing. Provide agitation to prevent such separation.

Do not add any wetting agents or detergents to the spray solution.

CAUTION: A sprayer or other container previously used to apply 2,4-D, MCPA, Silvex or 2,4,5-T must be thoroughly cleaned with alkali and water before using for this product. Even a small amount of one of these chemicals mixed with this product can cause serious injury to legumes. Contact your supplier for special instructions on cleaning out spray equipment.

LIMITED WARRANTY AND DISCLAIMER

This product, when used in accordance with the directions on the label, is warranted to perform as represented on the label. The product is not a substitute for good agricultural practices. The user assumes all responsibility for the proper use of this product. The manufacturer disclaims any liability for damage to crops or livestock resulting from the use of this product. The manufacturer disclaims any liability for damage to crops or livestock resulting from the use of this product. The manufacturer disclaims any liability for damage to crops or livestock resulting from the use of this product.



BROADLEAF HERBICIDE

Controls Broadleaved Weeds in Alfalfa and Birdsfoot Trefoil

ACTIVE INGREDIENT:

*4-(2,4-Dichlorophenoxy) butyric acid, butoxyethanol ester 31.4%

INERT INGREDIENTS:

*4-(2,4-Dichlorophenoxy) butyric acid equivalent 24.6% by weight or 2 pounds per gallon 65.6%

EPA Reg. No. 264-145-2A

EPA Est. 264-110-1

CAUTION: Keep out of reach of children. See other cautions on side panel.

IN CASE OF EMERGENCY
PHONE COLLECT (24 HOURS A DAY)
IN U.S.A. (304) 744-3487

UNION CARBIDE AGRICULTURAL PRODUCTS COMPANY, INC.
P.O. Box 1014, T. A. Amherst, N.Y.
Research Triangle Park, N.C. 27709

H15407

CONTENTS: 5 Gallons (18.92 liters)

PRE
CAL
ENV
PRO
STO

in proportion to the
reach band is sprayed
and acre will be 1/3 of
rate sprayers before
the same height while

[illegible]

H15407
CONTENTS: 5 Gallons (18.92 liters)

[illegible]