

Naphthylacetamide

2-Methyl-1-Naphthylacetic Acid

2-Methyl-1-Naphthylacetic Acid

2-Methyl-1-Naphthylacetic Acid

3-Methyl-1-Naphthylacetic Acid

3-Methyl-1-

264-151

CAUTION—Keep out of reach of children. Harmful if swallowed. May cause skin irritation. Do not reuse container. Destroy when empty. Avoid prolonged contact. Do not take internally.

forms to the chemical description given in the ingredient statement and the product is suited for the purposes described when used according to directions. Because of the broad range of conditions which may be encountered with the use of this product, it is impossible to eliminate all risks, even though label directions are followed. Amchem therefore makes no other express or implied warranty, and no agent of Amchem is authorized to do so.

Select sturdy cuttings from healthy plants. Remove any flower buds or leaves from one or two nodes at the base of the cutting.

Dip the cutting into ROOTONE about '4 inch deeper than it will be set in the rooting medium. Tap off any excess powder and insert cutting in a hole in the rooting medium with at least one node covered. Make holes wide enough so that no ROOTONE is scraped off during planting. Firm medium around cutting to avoid air pockets.

Do not allow cuttings to become dry or wilted from the time they are cut until they are rooted. This is 3 to 5 weeks or longer, depending on plant species.

Among the plants that will root are azalea, barberry, boxwood, chionanthus, dogwood, euonymus, itex (holly), juniper, pachysandra, pyracantha, rhododendron, rose, spiraea, lilac, taxus (yew) and viburnum.

USDA Reg. No. 264-151

Form No. 2592-10-68G

Printed in U.S.A.