

1189

PM23

264-164

26 FEB 1992

Ms. Karen Shearer
Rhône-Poulenc
P. O. Box 12614
2 T.W. Alexander Dr
Research Triangle NC 27709

Dear Ms. Shearer:

Subject: Butyroc 175 Broadleaf Herbicide
EPA Registration No. 264-164
Re: Label Amendment
Your Letter Dated January 13, 1992

The labeling referred to above submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable provided that under the Environmental Hazards Section of the label replace the statement "do not apply directly to water of wetlands to read "do not apply directly to water in areas where surface water is present or to intertidal areas below the mean high water mark." Correct the foot note under Tank Mixtures of Butyroc in Soybeans to read 4 and 5 instead of 6 and 7. Please submit five (5) copies of your final printed labels incorporating this amendment. A stamped copy of labeling is enclosed for your records.

Sincerely

Joanne L Miller
Product Manager (25)
Fungicide-Herbicide Branch
Registration Division (H-70650)

Enclosure

BEST AVAILABLE COPY

CONCURRENCES							
SYMBOL	H75D5C						
SURNAME	Miller						
DATE	12/25/92						

BUTYRAC® 175 Broadleaf Herbicide

Controls Broadleaf Weeds in Soybeans and Peanuts

ACTIVE INGREDIENT: 4-(2,4-Dichlorophenoxy)butyric acid, dimethylamine salt* 23.0%
INERT INGREDIENT: 77.0%
*4-(2,4-Dichlorophenoxy)butyric acid equivalent 19.5% by weight or 1.75 pounds per gallon.

EPA Reg. No. 264-164

EPA Est. No. 264-MO-01

**KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO**

For **MEDICAL** and **TRANSPORTATION Emergencies ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-800-334-9745

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: If patient is conscious and alert, give 2-3 glasses of water to drink. Do not induce vomiting. Get medical attention.

IF ON SKIN: Immediately wash skin with plenty of soap and water, if available, while removing contaminated clothing and shoes. Wash clothing separately before reuse. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration. Administer oxygen if necessary. Get medical attention.

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for at least 15 minutes. Get medical attention, preferably an ophthalmologist.

NOTE TO PHYSICIAN

No specific antidote is available. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

PRECAUTIONARY STATEMENTS

DANGER

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin or clothing. Avoid breathing spray mist. When handling this product, wear chemical resistant gloves and goggles or face shield. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters.

MIXING AND LOADING: Most cases of ground water contamination involving phenoxy herbicides such as 2,4-DB have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-DB pesticides at such sites to prevent contamination of ground water supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent ground water contamination.

Do not apply BUTYRAC® 175 Broadleaf Herbicide directly to or otherwise permit it to come in contact with cotton, okra, grapes, tomatoes, fruit trees, vegetables, flowers, or other desirable crop or ornamental plants. Do not permit spray mist to drift onto susceptible plants since very small quantities of the BUTYRAC® 175 Broadleaf Herbicide can cause severe injury during the growing or dormant periods. Use coarse sprays to minimize drift. Do not apply with nozzles that produce fine spray droplets.

Drift from ground application may be reduced by: (1) keeping the spray boom as near to the crop as possible in order to obtain complete coverage; (2) by applying 10 gallons or more of spray per acre; (3) by using no more than 20 pounds of pressure at the nozzle tips; and (4) by not spraying when wind exceeds 5 miles per hour.

Drift from aerial application may be reduced by: (1) applying as near to the target as possible to obtain adequate coverage; (2) by applying 5 or more gallons of spray per acre; (3) by using 20 pounds pressure or less at the nozzle tips; (4) by using nozzles which produce a coarse spray pattern; (5) by spraying when there is no possibility for a temperature inversion at time of spraying.

Applications by aircraft, ground rig and hand sprayers should be carried out only when there is no hazard from spray drift.

26 FEB 1992

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

BEST AVAILABLE COPY

264-164

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Read entire label before using this product.

STORAGE AND DISPOSAL

STORAGE

Do not contaminate water, food or feed by storage or disposal. Store in original container in a dry secured storage area. Keep container tightly closed when not in use.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL CAUTIONS AND RESTRICTIONS

Do not apply BUTYRAC® 175 Broadleaf Herbicide through any type of irrigation system.

Do not use in or near a greenhouse.

APPLICATION PROCEDURES

BUTYRAC® 175 can be applied to registered use areas by ground, aerial and sprinkler irrigation equipment. The following provides recommended methods of application for each crop.

GROUND APPLICATION

Use a standard herbicide boom sprayer that provides uniform and accurate application. Sprayer should be equipped with screens no finer than 50 mesh in the nozzle tips and in-line strainers.

Select a spray volume and delivery system that will ensure thorough and uniform spray coverage. For optimum spray distribution and thorough coverage use of flat fan nozzles (maximum tip size 8008) with a minimum spray pressure of 30 psi at the nozzle tips are recommended. Other nozzle types that produce coarse spray droplets may not provide adequate coverage of the weeds to ensure optimum control. Raindrop nozzles are not recommended as weed control with BUTYRAC® may be reduced. In general a minimum spray volume of 10 gallons per acre (GPA) is recommended for optimum spray coverage. When using higher speed equipment a maximum speed of 10 mph is suggested if field conditions cause excessive boom movement during application and subsequent poor spray coverage. Ground applications made when dry, dusty field conditions exist may provide reduced weed control in wheel track areas.

AERIAL APPLICATION

Use orifice discs, cores and nozzle types and arrangements that will provide for optimum spray distribution and maximum coverage. In general a minimum spray volume of 5 GPA and a maximum pressure of 40 psi are recommended.

MIXING INSTRUCTIONS

BUTYRAC® ALONE: Fill the spray tank 1/2 to 3/4 full with clean water. Begin agitation and add the recommended amount of BUTYRAC® 175. Add water to the spray tank to the desired level. Maintain sufficient agitation to ensure a uniform spray mixture during application.

TANK MIXTURES: BUTYRAC® 175 can be applied in tank mixture with other herbicides registered for use on approved crops. Refer to the specific crop section for rate recommendations and other restrictions. To apply BUTYRAC® 175 in mixture with another product, fill the spray tank 1/2 to 3/4 full with clean water and begin agitation. If tankmixing with wettable powder, soluble powder flowable or dry flowable products, add the powder or flowable product first. After the other herbicide is thoroughly mixed with water, add the recommended amount of BUTYRAC® 175 and add water to the spray tank to the desired level. If tankmixing with other product types, add the BUTYRAC® 175 first before adding the other product. Always mix one product in water thoroughly before adding another product or compatibility problems may occur. Never mix two products together without first mixing in water.

Maintain sufficient agitation while mixing and during application to ensure a uniform spray mixture. If spray mixture is allowed to remain without agitation for short periods of time, be sure to agitate until uniformly mixed before application.

COMPATIBILITY

Evaluate tank mixtures not listed on this label for compatibility and crop safety on a small area before applying to the entire field.

BUTYRAC® 175 may form an insoluble precipitate in very hard water. If you expect to mix BUTYRAC® 175 with very hard water, test compatibility by mixing a small amount of BUTYRAC® 175 in the proposed dilution ratios, shake and observe. A compatibility agent approved for use on growing crops such as UNITE or E-Z MIX may be tested to reduce precipitation. Whenever hard water is used to dilute BUTYRAC® 175, spray immediately and do not allow spray mixture to sit overnight.

BEST AVAILABLE COPY

GENERAL INFORMATION

Spray tank residues of 2,4-D or MCPA mixed with BUTYRAC® 175 Broadleaf Herbicide can cause serious crop or ornamental plant injury. A sprayer previously used to apply these chemicals must be thoroughly cleaned with alkali and water before applying BUTYRAC® 175. Be sure sprayer is clean before applying BUTYRAC® 175 Broadleaf Herbicide.

Local conditions may affect the use of herbicides. Consult your State Agricultural Experiment Station, Farm Advisors, or Extension Weed Specialists for advice in selecting treatment from this label to best fit local conditions.

INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY DO NOT USE THIS PRODUCT.

Crop varieties vary in response to 2,4-DB and some are easily injured. Apply BUTYRAC® 175 only to varieties known to be tolerant to 2,4-DB. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-DB, consult your seed company, state Agricultural Extension Service or qualified crop consultant for advice.

Be sure that use of this product conforms to all applicable laws, rules, and regulations. Certain states have restrictions pertaining to application distances from susceptible crops. The applicator should become familiar with these laws, rules, or regulations and follow them exactly.

GENERAL WEED LIST

BUTYRAC® 175 broadleaf herbicide will control or suppress the following weeds depending on weed size.

COMMON NAME	MAXIMUM SIZE BOTANICAL NAME	CONTROLLED*
pigweed (carelessweed)	<i>Amaranthus spp.</i>	3 inches
morningglory	<i>Ipomoea spp.</i>	3 inches
cocklebur	<i>Xanthium spp.</i>	3 inches
velvetleaf	<i>Abutilon theophrasti</i>	1 inch
Virginia copperleaf	<i>Acalypha virginica</i>	1 inch
prickly sida (teaweed)	<i>Sida spinosa</i>	1 inch
common ragweed	<i>Ambrosia artemisiifolia</i>	1 inch
jimsonweed	<i>Datura stramonium</i>	1.5 inches
devilsclaw	<i>Proboscidea louisianica</i>	2 inches
thersal croton	<i>Croton lindheimeri</i>	1 inch
lambquarters	<i>Chenopodium album</i>	1 inch
wild mustard	<i>Sinapis arvensis</i>	1 inch
field pennycress	<i>Thlaspi arvense</i>	1 inch

*Growth of larger weeds will be only suppressed.

SOYBEANS

BUTYRAC® 175 Recommendations

PRODUCT	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
		CROP	WEEDS
Preplant through pre-emergence BUTYRAC® 175 + Non-ionic surfactant	0.8 - 1.0 pint/A + 0.5% V/V	Apply to soybeans before planting or before crop emergence.	For control of emerged cocklebur, annual morningglories and other susceptible broadleaf weeds, apply when weeds are small and actively growing (see GENERAL WEED LIST). BUTYRAC® 175 may not give complete control of large overwintered mustards. Best results will be achieved by adding a non-ionic surfactant at a 0.5% V/V to the spray tank when making a preplant or preemergence application of BUTYRAC® 175.

BEST RESULTS OBTAINED

5 of 9

SOYBEANS

BUTYRAC® 175 Recommendations

PRODUCT	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
		CROP	WEEDS
Postemergence broadcast (over the top) BUTYRAC® 175	0.8 - 1.0 pint/A	Apply to soybeans grown in the Southern states only from 7 to 10 days before bloom up to mid-bloom when soybeans are about knee-high and growing actively. Soybean foliage should be dark green indicating that nodulation and nitrification are under way. Post emergence broadcast application at these rates prior to or after this application timing is not recommended as reduced flowering and yield may result. DO NOT APPLY BUTYRAC® 175 postemergence broadcast to soybeans grown in the midwest states of IA, IL, IN, KS, KY (except the Purchase area), MO (except the MO bootheel), MI, MN, NE, ND, OH, SD and WI.	For control of emerged cocklebur, annual morningglories and other susceptible broadleaf weeds, apply when weeds are small and actively growing (see GENERAL WEED LIST).
Postemergence directed band. BUTYRAC® 175	0.8 - 1.8 pints/A	Apply when soybeans are 8 or more inches tall with sprayer nozzles mounted to insure proper placement of spray on only the lower 1/3 of the soybean plants. Do not allow spray to contact growing terminals of beans as excessive crop injury will result. <i>Do not mount nozzles on booms with drop pipes or on cultivators without gauge wheels. Use flat fan type nozzles, 8001 or larger or the equivalent with nozzle pressure less than 30 psi and at least 10 gallons of spray volume per acre</i>	To control emerged cocklebur and annual morningglories up to 3 inches tall, apply 0.7 to 0.9 pint/A per broadcast acre as a directed band treatment. To control other susceptible broadleaf weeds up to 1 inch tall, apply 1.4 to 1.6 pints per broadcast acre as a directed band treatment. Apply no more than 2 higher rate applications per season to reduce the potential for crop stunting.

TANK MIXTURES OF BUTYRAC® 175 BROADLEAF HERBICIDE AND OTHER HERBICIDES IN SOYBEANS

Applying tank mixtures of BUTYRAC® 175 Broadleaf Herbicide preplant/preemergence or postemergence with other soybean herbicides as directed below can reduce competition from early weed populations and can improve weed control or control mid-to-late-season weed populations, thus minimizing the likelihood of yield reduction in soybean fields with heavy broadleaf weed infestations. However, treating soybeans under stress (as from drought or disease such as Phytophthora root rot) or in any other manner not directed on this label, can cause crop injury and yield reduction. The advantage of treating weeds with an appropriate tank mixture should be balanced against any potential for loss. Base your decision on an evaluation of each factor. Follow the local recommendations of your state, Cooperative Extension Service, or other agricultural weed control authority. Always follow the recommendations of any product used in tank mixture with BUTYRAC® 175.

PRODUCT	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
		CROP	WEEDS
BUTYRAC® 175 + Gramoxone® Extra + Non-ionic surfactant	0.6 - 0.8 pint/A + 1.5 - 5 pints/A + 0.5% V/V	Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.	For improved control of emerged annual morningglory (ivy leaf, tall), common cocklebur, narrow leaf and other problem weeds, apply to small actively growing weeds. Refer to the Gramoxone® Extra label for full list of weed species controlled and specific application stage and rate recommendations.

BEST AVAILABLE COPY

TANK MIXTURES OF BUTYRAC® 175 BROADLEAF HERBICIDE AND OTHER HERBICIDES IN SOYBEANS

PRODUCT	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
		CROP	WEEDS
BUTYRAC® 175 + Prowl ² + Non-ionic surfactant	0.8 - 1.0 pint/A + 1.0 - 3.0 pints/A + 0.5% V/V	Apply up to 45 days prior to soybean planting. Do not apply a BUTYRAC® 175 + Prowl ² tank mixture at or after planting North of Interstate 80. Surface applications of Prowl ² tank mixtures North of Interstate 80 require at least 1 inch rainfall or mechanical incorporation prior to planting or crop injury may result. Do not apply this tank mixture after crop emergence.	For control of emerged cocklebur, annual morningglories and other susceptible broadleaf weeds, apply when broadleaf weeds are actively growing and small (see GENERAL WEED LIST). BUTYRAC® 175 mixtures may not give complete control of larger overwintered mustards. Best results will be achieved by adding a non-ionic surfactant to the spray tank when making a preplant application.
BUTYRAC® 175 + Roundup ³ or Honcho ³ + Non-ionic surfactant	0.6 - 0.8 pint/A + 1 - 1.5 pints/A + 0.5 - 1.0% V/V	Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.	For improved control of emerged annual morningglories (ivyleaf, tall), common cocklebur and other problem weeds, apply to small actively growing weeds. Refer to the Roundup ³ or Honcho labels for specific rate recommendations, application stage and weed species controlled.
BUTYRAC® 175 + Pursuit Plus ² + Non-ionic surfactant	0.6 - 0.8 pint/A + 2.5 pints/A + 0.25% V/V	Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.	For improved control of emerged mustards, field pennycress and other problem weeds, apply when broadleaf weeds are actively growing and small (see GENERAL WEED LIST).
BUTYRAC® 175 + Scepter ² or Scepter 70DG ² + Non-ionic surfactant	0.6 - 0.8 pint/A + 0.66 pint/A or 2.8 oz./A + 0.25% V/V	Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.	For improved control of emerged mustards, field pennycress and other problem weeds, apply when broadleaf weeds are actively growing and small (see GENERAL WEED LIST).
BUTYRAC® 175 + Squadron ² + Non-ionic surfactant	0.6 - 0.8 pint/A + 3 pints/A + 0.25% V/V	Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.	For improved control of emerged mustards, field pennycress and other problem weeds, apply when broadleaf weeds are actively growing and small (see GENERAL WEED LIST).

BEST AVAILABLE COPY

7-89

TANK MIXTURES OF BUTYRAC® 175 BROADLEAF HERBICIDE AND OTHER HERBICIDES IN SOYBEANS

PRODUCT	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
		CROP	WEEDS
BUTYRAC® 175 + Lorox®	0.9 pint/A + 1 pint/A	Apply to soybeans as a directed band treatment only when soybeans are at least 8 inches high. Do not spray higher than 3 inches on the soybean stem or unacceptable crop injury may result. Do not spray over the top of soybean plants.	For postemergence control of annual morningglories, sicklepod, teaweed, cocklebur and sesbania, apply when weeds do not exceed 4 inches in height. A second application may be necessary, but do not make more than 2 applications per season.
BUTYRAC® 175 + Sencor DF	1.0 pint/A + 0.33 - 0.66 lb/A	Apply to soybeans as a directed band treatment only when soybeans are at least 8 inches high, with spray or nozzles mounted to insure proper placement of spray on no more than the lower 1/3 of the soybean plants. Do not apply directly to soybean plants or serious crop injury will occur. Soybean leaves contacted by spray will be killed. Follow all variety restrictions on the full Sencor DF label.	For improved control of cocklebur, annual morningglories and other broadleaf weeds, apply before weeds are 3 inches tall. A non-ionic surfactant may be added to improve broadleaf weed control.

RESTRICTIONS AND LIMITATIONS FOR USE ON SOYBEANS

- Beans stressed by drought or other influences should not be sprayed.
- Do not use this product on soybeans that show symptoms of disease such as Phytophthora root rot.
- Do not graze or feed soybean hay within 60 days after application of BUTYRAC® 175 tank mix application.
- Do not harvest beans within 60 days after spray application.
- Do not treat soybeans with a tank mixture of BUTYRAC® 175 Broadleaf Herbicide and SEVIN® Carbaryl Insecticide as severe injury may result.
- When preplant through preemergence treatment is followed with a BUTYRAC® 175 postemergence application, the cumulative rate should not exceed 1.6 pints per acre per season.
- Follow all restrictions and limitations of any product used in tank mixture with BUTYRAC® 175.
- Do not use BUTYRAC® 175 alone or in tank mixture as a preplant through preemergence application to soybeans in California.

PEANUTS

BUTYRAC® 175 Recommendations

PRODUCT	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
		CROP	WEEDS
BUTYRAC® 175	0.9 - 1.1 pints/A	Apply to peanuts 2 to 12 weeks after planting in the states of AL, AR, FL, GA, LA, NC, SC, TN and VA. In NC, SC and VA, do not apply later than 45 days before harvest.	For control of annual morningglories, cocklebur and other broadleaf weeds, apply when weeds are small and actively growing (see GENERAL WEED LIST). A second application may be made for late germinating cocklebur and morningglories.
	0.9 - 1.8 pints/A	Apply to peanuts 2 to 12 weeks after planting in the states of OK, TX and NM only. Do not apply later than 100 days after planting or 30 days before harvest.	For control of annual morningglories and cocklebur, apply 0.8 - 1.0 pint/A of BUTYRAC® 175 when the weeds are actively growing and before they are 3 inches tall. For optimum control of other susceptible broadleaf weeds (see GENERAL WEED LIST), apply BUTYRAC® 175 at 1.6 pints/A. For optimum prickly sida suppression, make a second application 14 days later.

RESTRICTIONS AND LIMITATIONS FOR USE ON PEANUTS

Do not feed treated peanut vines or peanut hay to livestock.

Do not apply BUTYRAC® 175 Broadleaf Herbicide if peanut plants are under stress from drought as injury may occur.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants that this product conforms to the chemical description on the label; that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.

¹ Gramoxone and Reflex are registered trademarks of ICI Americas, Inc.

² Prowl, Pursuit Plus, Scepter, Scepter 70DG and Squadron are registered trademarks of American Cyanamid Company.

³ Roundup and Honcho are registered trademarks of Monsanto Company.

⁴ Lorox is a registered trademark of E.I. DuPont de Nemours Co., Inc.

⁵ Sencor DF is a registered trademark of Bayer AG, Germany.

Rhône-Poulenc Ag Company
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709

BUTYRAC and SEVIN are registered trademarks of Rhône-Poulenc.
BUTYRAC® 175 Broadleaf Herbicide (PENDING) Submitted 1/10/92