

FENAMINE
264-172

GENERAL DIRECTIONS

Apply FENAMINE to small weeds in the spring or early summer, or before weeds have matured and gone to seed. It is not necessary to spray bare ground before weed growth starts in the spring. Weeds treated after they have developed strong stems may remain standing for the rest of the season even though they are dead. These standing dead weeds are unsightly and a fire hazard. Therefore, for best results on tall weeds, cut or mow to ground line and remove. Treat one or two weeks later when regrowth appears. FENAMINE acts slowly but thoroughly. Weeds start to turn white or brown within a week and are usually dead in about three weeks. Small weeds growing actively are killed more rapidly than weeds in a dormant or dried condition. When johnsongrass or other resistant weeds are not controlled by the first application, treat again with the same rate when regrowth is 6 to 12 inches tall (about 2 months).

Shake container before using

AMOUNTS TO USE—CONVENTIONAL SPRAYERS

General Recommendations: For mixed annual and perennial grasses and broadleaf weeds where annual rainfall is usually 15 to 20 inches and above, apply 9 gallons FENAMINE per acre in enough water to wet weeds thoroughly. At least 100 gallons per acre is suggested.

Special Situations: For annual grasses and broadleaf weeds where annual rainfall is usually below 15 inches, apply 4½ gallons FENAMINE per acre in enough water to wet weeds thoroughly. At least 100 gallons water per acre is suggested.

MIXING SPRAYS

Half fill the spray tank with water. While agitating, add the required FENAMINE. Then add the rest of the water. This product forms a suspension when mixed with water. Use continuous agitation while spraying. To clean FENAMINE from spray equipment, flush several times with clear water after each use.

FENAMINE may be applied with a watering can, a portable pressure sprayer, a knapsack sprayer or a motor driven pump-type sprayer. Screens in nozzles, and suction lines should be no finer than 50 mesh size. Use a Tee-Jet 8002 or larger fan type of nozzle or its equivalent.

SOME OF THE WEEDS CONTROLLED BY RECOMMENDED RATES OF FENAMINE

alfalfa	downy brome	pigweed	ryegrass
bluegrass	(chest)	plantain	smartweed
Canada thistle	filaree	poison ivy	sowthistle
chrysanthemum	foxtail	puncture vine	tarweed
weed (mugwort)	kochia	purslane	velvetgrass
crabgrass	lambsquarters	quackgrass	wild oats
dandelion	mustard	ragweed	witchgrass
	nightshade	Russian thistle	

KEEP FROM FREEZING

To control weeds in winter, apply FENAMINE to weeds before they freeze. Do not apply to frozen weeds. The weed must be green and growing when FENAMINE is applied. Apply FENAMINE to weeds before they freeze. Do not apply to frozen weeds. The weed must be green and growing when FENAMINE is applied.